

# Electrogenicity of the Na<sup>+</sup>-ATPase from the marine microalga *Tetraselmis (Platymonas) viridis* and associated H<sup>+</sup> countertransport

Yurii V. Balnokin\*, Larissa G. Popova, Igor M. Andreev

Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, Botanicheskaya 35, Moscow 127276, Russia

Received 20 October 1999

Edited by Vladimir Skulachev

**Abstract** Sodium accumulation by the Na<sup>+</sup>-ATPase in the plasma membrane (PM) vesicles isolated from the marine alga *Tetraselmis (Platymonas) viridis* was shown to be accompanied by  $\Delta\psi$  generation across the vesicle membrane (positive inside) and H<sup>+</sup> efflux from the vesicle lumen. Na<sup>+</sup> accumulation was assayed with <sup>22</sup>Na<sup>+</sup>;  $\Delta\psi$  generation was detected by recording absorption changes of oxonol VI; H<sup>+</sup> efflux was monitored as an increase in fluorescence intensity of the pH indicator pyranine loaded into the vesicles. Both ATP-dependent Na<sup>+</sup> uptake and H<sup>+</sup> ejection were increased by the H<sup>+</sup> ionophore carbonyl cyanide *m*-chlorophenylhydrazone (CICCP) while  $\Delta\psi$  was collapsed. The lipophilic anion tetraphenylboron ion (TPB<sup>−</sup>) inhibited H<sup>+</sup> ejection from the vesicles and abolished  $\Delta\psi$ . Based on the effects of CICCP and TPB<sup>−</sup> on H<sup>+</sup> ejection and  $\Delta\psi$  generation, the conclusion was drawn that H<sup>+</sup> countertransport observed during Na<sup>+</sup>-ATPase operation is a secondary event energized by the electric potential which is generated in the course of Na<sup>+</sup> translocation across the vesicle membrane. Increasing Na<sup>+</sup> concentrations stimulated H<sup>+</sup> efflux and caused the decrease in the  $\Delta\psi$  observed, thus indicating that Na<sup>+</sup> is likely a factor controlling H<sup>+</sup> permeability of the vesicle membrane.

© 1999 Federation of European Biochemical Societies.

**Key words:** Marine alga; Na<sup>+</sup>-ATPase; Na<sup>+</sup>/H<sup>+</sup> countertransport; Na<sup>+</sup> pump; Plasma membrane; *Tetraselmis (Platymonas) viridis*

## 1. Introduction

The functioning of the primary Na<sup>+</sup> pump in the plasma membrane (PM) of the marine green microalga *Tetraselmis (Platymonas) viridis* has been shown by us earlier [1,2]. The inverted PM vesicles isolated from this organism were able to transport <sup>22</sup>Na<sup>+</sup> from outside into the vesicle lumen upon

MgATP addition. The process occurred in the presence of the protonophore carbonyl cyanide *m*-chlorophenylhydrazone (CICCP) and amiloride, an inhibitor of the Na<sup>+</sup>/H<sup>+</sup> antiporters, thus indicating that ATP-dependent sodium uptake by the PM vesicles was performed by a proton-motive-force-independent fashion. The ATP-driven sodium uptake was sensitive to orthovanadate, and pH optimum of the process was 7.8–8.0. Recently, in the PM preparations isolated from *T. viridis*, a polypeptide with apparent molecular mass 100 kDa being phosphorylated in the presence of [ $\gamma$ -<sup>32</sup>P]ATP was demonstrated by means of acid PAGE in combination with autoradiography [3]. The 100 kDa polypeptide phosphorylation was shown to require the presence of sodium ions in the reaction mixture. The phosphorylated polypeptide was sensitive to hydroxylamine treatment, thus indicating that an acyl-phosphate bond was formed during the phosphorylation reaction. The results of the above-mentioned experiments demonstrated that the primary Na<sup>+</sup> pump of *T. viridis* is a Na<sup>+</sup>-transporting ATPase forming the phospho-intermediate in the course of its catalytic cycle and, therefore, belonging to the family of the p-type ATPases.

An important problem related to the mechanism of ATP-dependent Na<sup>+</sup> transport is whether a counterion is utilized by the Na<sup>+</sup>-ATPase from *T. viridis*. Many cation-translocating p-type ATPases are known to operate by cation exchange, for example Na<sup>+</sup>,K<sup>+</sup>-ATPase and H<sup>+</sup>,K<sup>+</sup>-ATPase from mammalian cells [4], Ca<sup>2+</sup>,H<sup>+</sup>-ATPase from sarcoplasmic reticulum [5], Ca<sup>2+</sup>-ATPase from plant plasma membrane that catalyzes Ca<sup>2+</sup>/H<sup>+</sup> exchange as well [6]. Na<sup>+</sup>-ATPase from the brown microalga *Heterosigma akashiwo* was suggested to be a Na<sup>+</sup>,K<sup>+</sup>-exchanging ATPase as judged by (i) synergistic stimulation of the PM ATPase activity by Na<sup>+</sup> and K<sup>+</sup> ions, and (ii) dephosphorylation of the phosphorylated intermediate of the ATPase by K<sup>+</sup> ions [7,8]. Since K<sup>+</sup> inhibited Na<sup>+</sup> translocation performed by the Na<sup>+</sup>-ATPase from *T. viridis* [2] and there was no synergistic stimulation of the PM ATPase activity by K<sup>+</sup> and Na<sup>+</sup> [9], it is hardly possible to ascribe the role of the counterion to K<sup>+</sup> in the present case. Another possibility is that H<sup>+</sup> serves as a counterion in primary Na<sup>+</sup> transport in the *T. viridis* plasma membrane.

In this work we demonstrate on isolated PM vesicles from *T. viridis* that ATP-dependent <sup>22</sup>Na<sup>+</sup> uptake performed by the Na<sup>+</sup>-ATPase is accompanied by  $\Delta\psi$  generation across the vesicle membrane and alkalization of the vesicle lumen, i.e. H<sup>+</sup> efflux from the vesicles. The data obtained allowed us to conclude that the Na<sup>+</sup>-ATPase from *T. viridis* PM operates as an electrogenic uniporter. H<sup>+</sup> is a counterion for Na<sup>+</sup> and movement of H<sup>+</sup> out of the vesicle lumen is energized by  $\Delta\psi$  generated in the course of Na<sup>+</sup> translocation by Na<sup>+</sup>-ATPase.

\*Corresponding author. Fax: (7)-095-482 16 85.  
E-mail: balnokin@ippras.ru

**Abbreviations:** BTP, 1,3-bis-(tris(hydroxymethyl)methylamino)propane; CICCP, carbonyl cyanide *m*-chlorophenylhydrazone; EGTA, ethylene glycol-bis(β-aminoethyl ether)-*N,N,N',N'*-tetraacetic acid; HEPES, *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid; MES, 2-[*N*-morpholino]ethanesulfonic acid; oxonol VI, bis-(3-propyl-5-oxo-isoxazol-4-yl)pentamethine oxonol; PM, plasma membrane; pyranine, 8-hydroxy-1,3,6-pyrene trisulfonate; TPB<sup>−</sup>, tetraphenylboron ion;  $\Delta\psi$ , membrane electric potential difference

## 2. Materials and methods

### 2.1. Plant material and preparation of PM vesicles

*Tetraselmis (Platymonas) viridis* Rouch. cells were cultured in artificial sea water containing 0.45 M NaCl as described elsewhere [2].

The PM vesicles were prepared according to the method described earlier [9] with some minor modifications [2]. The final membrane pellet was suspended in medium containing 0.5 M mannitol, 10 mM 1,3-bis-(tris(hydroxymethyl)methylamino)propane-*N*-2-hydroxyethyl-piperazine-*N'*-2-ethanesulfonic acid (BTP-HEPES), pH 7.8, 2 mM MgSO<sub>4</sub>, 1 mM dithiothreitol, 0.4 mM ethylene glycol-bis(β-aminoethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA), and 10 μg/ml phenylmethylsulfonyl fluoride.

### 2.2. Na<sup>+</sup> uptake

Sodium uptake by the PM vesicles was monitored by a tracer technique using <sup>22</sup>Na<sup>+</sup> as described in [1]. The assay mixture contained 0.5 M mannitol, 20 mM BTP-HEPES, pH 7.8, 25 mM NO<sub>3</sub><sup>-</sup>-BTP, 1 mM MgSO<sub>4</sub>, and ca. 600 μg of membrane protein. The reaction was initiated by successive addition of 10 mM <sup>22</sup>NaCl up to 0.5 MBq and 2 mM ATP (Tris salt). After defined time intervals, 60 μl aliquots of the suspension were taken, and vesicles were separated from the medium by filtering through a Synpore nitrocellulose filter with a pore size of 0.6 μm. Filters were washed three times with 1 ml of the assay solution free of the label and ATP, and radioactivity retained on the filters was counted with a scintillation counter.

### 2.3. Detection of transmembrane electric potential

Formation of transmembrane electric potential accompanying the Na<sup>+</sup> pump operation was detected by continuously recording the differential absorption change (621–582 nm) of the ΔΨ indicator oxonol VI [10] with Hitachi 557, a dual wavelength spectrophotometer. The assay was performed in the reaction medium containing

0.5 M sucrose, 20 mM BTP-MES, pH 7.8, 1 mM MgSO<sub>4</sub>, 5 mM Na<sub>2</sub>SO<sub>4</sub>, 1 mM EGTA, 3 μM oxonol VI, and ca. 100 μg of membrane protein in 2 ml. After 15 min incubation of the vesicles in the medium, 2 mM ATP (Tris salt) was added to initiate the ATP-driven ΔΨ generation.

### 2.4. Detection of H<sup>+</sup> ejection from the vesicles

The formation of the interior-alkaline pH gradient across the vesicle membranes in the course of Na<sup>+</sup> pump operation (luminal alkalization) was detected by monitoring the changes in fluorescence intensity of the pH indicator pyranine loaded into the vesicles. To load pyranine into the vesicles, isolation of the PMs was carried out in the presence of 100 μM pyranine added to the solutions used. To eliminate pyranine outside the vesicles, 0.5 ml final vesicle suspension was passed through a 1 × 10 cm column with G-50 Sephadex.

The fluorescence of pyranine loaded into the PM vesicles was recorded with a Hitachi 850 fluorescence spectrophotometer set at 450 nm (excitation) and 510 nm (emission). The assay was performed in 1 ml of the reaction medium containing 0.5 M mannitol, 20 mM BTP-HEPES, pH 7.8, 50 mM NO<sub>3</sub><sup>-</sup>-BTP, 1 mM MgSO<sub>4</sub>, and ca. 40 μg of membrane protein. After 15 min incubation of the vesicles in the medium, the reaction was initiated by successive addition of 10 mM Na<sub>2</sub>SO<sub>4</sub> and 2 mM ATP (Tris salt).

### 2.5. Protein determination

Protein content was determined by the micromethod of Simpson and Sonne [11] with bovine serum albumin as a standard.

## 3. Results

The PM vesicles isolated from *T. viridis* accumulated <sup>22</sup>Na<sup>+</sup> in the presence of <sup>22</sup>NaCl (Fig. 1A). Sodium uptake was sig-




Fig. 1. <sup>22</sup>Na<sup>+</sup> translocation into the PM vesicles isolated from *T. viridis* (A) and associated processes: luminal alkalization (B) and generation of membrane potential (C). Additions made: 10 mM Na<sub>2</sub>SO<sub>4</sub>, 2 mM ATP (Tris salt), 12 μM CICCP. The conditions of the experiments are described in Section 2.




Fig. 2. Effects of vanadate and  $\text{TPB}^-$  on luminal alkalization (A) and membrane potential generation (B) in the PM vesicles from *T. viridis*. Additions made: 10 mM  $\text{Na}_2\text{SO}_4$ , 2 mM ATP (Tris salt), 500  $\mu\text{M}$  vanadate, 100  $\mu\text{M}$   $\text{TPB}^-$ . The conditions of the experiments were as described in Section 2, except for traces 3 and 4 in A, where vanadate (trace 3) and  $\text{TPB}^-$  (trace 4) were present in the reaction mixture initially, before additions of  $\text{Na}_2\text{SO}_4$  and ATP.

nificantly accelerated in response to ATP addition when the reaction medium was supplied with  $\text{Mg}^{2+}$  ions and the reaction was performed in the weakly alkaline medium. The protonophore CCCP added to the medium following ATP enhanced  $\text{Na}^+$  uptake. These results are consistent with our previous data on sodium transport in *T. viridis* PM vesicles [2] and show that ATP-linked  $\text{Na}^+$  translocation across the PM is mediated by a primary  $\text{Na}^+$  pump ( $\text{Na}^+$ -ATPase) in this organism.

The experiments carried out on the PM vesicles loaded with the pH indicator pyranine demonstrated that  $\text{Na}^+$  uptake into the vesicles is accompanied by alkalization of the vesicle lumen indicating  $\text{H}^+$  efflux from the vesicles to the outer medium (Fig. 1B, trace 1). Addition of  $\text{Na}^+$  ions to the vesicle suspension initiated  $\text{H}^+$  efflux from the vesicle interior. This  $\text{H}^+$  efflux coupled with  $\text{Na}^+$  uptake obviously reflects the functioning of the  $\text{Na}^+/\text{H}^+$  antiporter in the vesicular membrane. ATP accelerated  $\text{H}^+$  efflux to a great extent, implying the involvement of the ATPase reaction in the  $\text{H}^+$  translocation.  $\text{Na}^+$  ions were found to be a prerequisite of ATP-dependent luminal alkalization. In the absence of  $\text{Na}^+$ , ATP added to the reaction mixture resulted in a slight  $\text{H}^+$  uptake by the vesicles, thus indicating some residual activity of the  $\text{H}^+$  pump at pH 7.8 (Fig. 1B, trace 2). Proton-translocating ATPase with optimum activity at pH 6.5–7.0 was demonstrated in *T. viridis* PM earlier [1,12]. Subsequent addition of  $\text{Na}^+$  to the suspension turned  $\text{H}^+$  flux to the opposite direction and led to luminal alkalization so that intravesicular pH reached a much higher value than the initial one observed before ATP addition. In this case,  $\text{Na}^+$  ions not only led to dissipation of the  $\Delta\text{pH}$  (acid inside) generated by the  $\text{H}^+$  pump, but also initiated operation of the  $\text{Na}^+$  pump

accompanied by  $\text{H}^+$  translocation from the vesicle lumen to the outside.

The protonophore CCCP stimulated ATP-dependent luminal alkalization just as observed for the ATP-dependent  $\text{Na}^+$  uptake. Stimulatory effects of CCCP on ATP-dependent sodium uptake and luminal alkalization (Fig. 1A,B) indicate that an overall excess of positive charge inhibiting subsequent transfer of  $\text{Na}^+$  ions to the vesicle interior is produced




Fig. 3. Effects of increasing  $\text{Na}^+$  concentrations on ATP-dependent luminal alkalization ( $\Delta\text{pH}$ ) and membrane potential generation ( $\Delta\Psi$ ) in the PM vesicles from *T. viridis*.

in the vesicle lumen due to the  $\text{Na}^+$ -ATPase operation. It may be assumed that CICCIP-facilitated  $\text{H}^+$  diffusion out of the vesicle lumen converts electric potential difference (positive inside) into  $\Delta\text{pH}$  (alkali inside) and thus stimulates ATP-dependent  $\text{Na}^+$  uptake. To confirm the electrogenic nature of  $\text{H}^+$  transfer out of the vesicles, the effect of lipophilic anion  $\text{TPB}^-$  on vesicle alkalinization was tested.  $\text{TPB}^-$ , known to induce negative electric potential at the vesicular membrane, inhibited the ATP-dependent luminal alkalinization when added following ATP or presented initially in the reaction mixture (Fig. 2A, traces 2 and 4). These results suggest that the ATP-dependent luminal alkalinization is mediated by the  $\Delta\Psi$  (positive inside) generated across the vesicle membrane during  $\text{Na}^+$  translocation by  $\text{Na}^+$ -ATPase. It should be noted that  $\text{TPB}^-$  did not affect  $\text{Na}^+$ -induced  $\text{H}^+$  efflux from the vesicles observed before ATP addition (Fig. 2A, trace 4).

The monitoring of electric potential difference across the vesicle membrane with oxonol VI, a  $\Delta\Psi$  indicator, demonstrated that ATP added to the vesicles suspended in the reaction mixture permitting  $\text{Na}^+$  uptake resulted in  $\Delta\Psi$  generation (positive inside vesicle lumen) (Fig. 1C). CICCIP collapsed  $\Delta\Psi$  generated upon ATP addition (Fig. 1C) as well as  $\text{TPB}^-$  (Fig. 2B, trace 1).

Like ATP-dependent  $\text{Na}^+$  uptake by the PM vesicles from *T. viridis* [2], ATP-dependent luminal alkalinization and  $\Delta\Psi$  generation across the vesicle membrane are related to p-type ATPase functioning as indicated by the sensitivity of both processes to orthovanadate. Orthovanadate inhibited luminal alkalinization (Fig. 2A, trace 1) and  $\Delta\Psi$  generation (Fig. 2B, trace 2), although higher concentrations of orthovanadate were necessary than usually used for inhibition of plant PM ATPases [13]. The inhibition of ATP-dependent luminal alkalinization was observed when orthovanadate was introduced into the reaction mixture in the course of  $\text{H}^+$  translocation (Fig. 2A, trace 1) as well as when it was added initially, prior to  $\text{Na}^+$  and ATP (Fig. 2A, trace 3).

Fig. 3 shows the effects of various  $\text{Na}^+$  concentrations on the ATP-dependent luminal alkalinization and  $\Delta\Psi$  formation. The initial rate and final level of luminal alkalinization increased as the  $\text{Na}^+$  concentration increased. The dependence of the initial rate of luminal alkalinization on  $\text{Na}^+$  concentrations demonstrates a saturable kinetics with an apparent  $K_m$  of about 5 mM. This value is appreciated as the sodium concentration at which a semi-maximal rate of intravesicular alkalinization is achieved. Opposite to ATP-dependent luminal alkalinization, the  $\Delta\Psi$  generated upon ATP addition declined as the  $\text{Na}^+$  concentration increased. This result suggests the rise in  $\text{H}^+$  permeability of the vesicle membrane as  $\text{Na}^+$  concentrations increased and, thus, stimulated conversion of  $\Delta\Psi$  (positive inside) into  $\Delta\text{pH}$  (alkaline inside). ATP-dependent  $\Delta\Psi$  generation observed in the absence of  $\text{Na}^+$  in the reaction mixture (Fig. 3) is likely mediated by a residual activity of the  $\text{H}^+$  pump.

#### 4. Discussion

The results of the present work demonstrate clearly that (i)  $\text{Na}^+$  translocation across *T. viridis* PM by  $\text{Na}^+$ -ATPase is an electrogenic process, and (ii)  $\text{H}^+$  is a counterion for  $\text{Na}^+$ . Electrogenicity of the  $\text{Na}^+$ -ATPase from *T. viridis* PM has been demonstrated in the experiments with the  $\Delta\Psi$  indicator, oxonol VI, by direct recording of the electric potential differ-

ence generated across the vesicle membrane in response to ATP addition in the presence of  $\text{Na}^+$  ions.  $\text{H}^+$  countertransport from the vesicle lumen to the exterior was detected by recording the ATP-dependent luminal alkalinization in the experiments with the pH indicator pyranine entrapped in the PM vesicles.

We found that the observed ATP-dependent processes, such as  $\text{Na}^+$  uptake,  $\Delta\Psi$  generation and luminal alkalinization, are interrelated and reflect the operation of the electrogenic  $\text{Na}^+$ -ATPase in *T. viridis* PM. The following facts allowed us to draw such a conclusion:

1. the requirement for  $\text{Na}^+$  ions, substrate of the  $\text{Na}^+$ -ATPase for the ATP-dependent luminal alkalinization (Fig. 1B, trace 2);
2. the sensitivity of both ATP-dependent  $\Delta\Psi$  generation and luminal alkalinization to orthovanadate (Fig. 2), the inhibitor of the *T. viridis* PM  $\text{Na}^+$ -ATPase [2,3];
3. the similar kinetics of  $\text{Na}^+$  uptake and luminal alkalinization (Fig. 1A,B);
4. the occurrence of all three processes at the same pH 7.8–8.0.

The abolition of the  $\Delta\Psi$  (positive inside) generated across the vesicular membrane upon ATP addition (Fig. 2B) and the inhibition of ATP-dependent luminal alkalinization (Fig. 2A, traces 2,4) by the lipophilic anion  $\text{TPB}^-$  not only confirm the electrogenicity of the  $\text{Na}^+$ -ATPase but also suggest that ATP-dependent  $\text{H}^+$  efflux from the vesicle lumen to the outside is a secondary event with respect to the electrogenic  $\text{Na}^+$  uptake. The  $\Delta\Psi$ , positive inside, generated in the course of ATP-dependent  $\text{Na}^+$  translocation produces a moving force for  $\text{H}^+$  transport to the exterior.  $\text{TPB}^-$  diminishing the  $\Delta\Psi$  across the vesicle membrane abolishes the driving force for  $\text{H}^+$  translocation.

The effects of the protonophore CICCIP on the  $\text{Na}^+$  and  $\text{H}^+$  transport and, accordingly, on the electrogenesis in the PM vesicles from *T. viridis* are in agreement with the effects exerted by  $\text{TPB}^-$ . CICCIP stimulated ATP-dependent  $\text{Na}^+$  uptake by the vesicles (Fig. 1A) and luminal alkalinization (Fig. 1B), and led to dissipation of the ATP-induced  $\Delta\Psi$  generated on the vesicular membrane (Fig. 1C). These findings can also be explained by assuming that the  $\text{Na}^+$ -ATPase of *T. viridis* is an electrogenic uniporter, and generation of  $\Delta\Psi$ , positive inside vesicles, is necessary for  $\text{H}^+$  efflux to the exterior. A low  $\text{H}^+$  permeability of the PM most likely limits the  $\text{H}^+$  translocation rate and, hence, the  $\text{Na}^+$  influx into vesicles. Addition of CICCIP results in facilitation of  $\text{H}^+$  transport and, thereby, induces conversion of  $\Delta\Psi$ , positive inside, into  $\Delta\text{pH}$ , alkaline inside, thus favoring the operation of the electrogenic  $\text{Na}^+$  pump and acceleration of  $\text{Na}^+$  uptake.

The dependences of the ATP-induced luminal alkalinization and  $\Delta\Psi$  generation on  $\text{Na}^+$  concentration (Fig. 3) show that  $\text{Na}^+$  not only is the substrate for the  $\text{Na}^+$ -ATPase but also appears to be a regulator of  $\text{H}^+$  efflux from the vesicles, i.e.  $\text{Na}^+$  is a factor controlling  $\text{H}^+$  permeability of the vesicle membranes. This conclusion follows from the observed reciprocal correlation between the  $\Delta\Psi$  and  $\Delta\text{pH}$  generated during  $\text{Na}^+$  pump operation at increasing  $\text{Na}^+$  concentration. At a low  $\text{Na}^+$  concentration, when the  $\text{H}^+$  permeability of PM is low, the  $\Delta\mu\text{H}$  is presented mainly in the form of  $\Delta\Psi$ . Increasing  $\text{Na}^+$  concentrations provide enhancement of  $\text{H}^+$  conduc-

tivity, as may be suggested, by opening some specific  $H^+$  channels, thus promoting conversion of  $\Delta\Psi$  into  $\Delta pH$ .

The obtained results do not agree with the model of  $Na^+/H^+$  antiport catalyzed by the  $Na^+$ -ATPase directly, but provide strong evidence that this enzyme is an electrogenic uniporter which translocates only  $Na^+$  ions while  $H^+$  movement is electrophoretically driven in response to generation of  $\Delta\Psi$  across the vesicle membrane.

Another  $Na^+$ -transporting system displayed in the experiments on isolated PM vesicles from *T. viridis* is the  $Na^+/H^+$  antiporter. Addition of  $Na^+$  to the vesicle suspension in the absence of ATP resulted in  $Na^+$  uptake and luminal alkalization (Fig. 1A,B). In this case,  $Na^+$ -induced luminal alkalization was not affected by the lipophilic anion  $TPB^-$ , indicating that the  $Na^+/H^+$  antiporter in the PM of *T. viridis* is an electroneutral system exchanging  $Na^+$  and  $H^+$  with a 1:1 stoichiometry.

In conclusion, one point should be commented upon concerning the effects of  $NO_3^-$  ions used as a component of the reaction mixtures in our experiments on sodium uptake and luminal alkalization.  $NO_3^-$  is an anion that generally penetrates easily across biological membranes. It is often used as a charge-compensating anion to abolish  $\Delta\Psi$  (positive inside) on the vesicular membranes during the measurements of ATP-driven  $H^+$ -pumping activity accomplished with optical pH indicators. Hence, one should expect  $NO_3^-$  ions will cause a decay of  $\Delta\Psi$  generated on the vesicular membrane from *T. viridis* as well. As a result of dissipation of the membrane potential difference, stimulation of ATP-dependent sodium uptake and inhibition of associated  $\Delta\Psi$ -driven luminal alkalization should be predicted. Indeed,  $NO_3^-$  ions stimulated ATP-dependent sodium uptake by the PM vesicles from *T. viridis* [2], but, unexpectedly, did not inhibit the intravesicular alkalization under investigation. On the contrary, they

stimulated the latter process (not shown). Therefore, we suggest that the stimulatory effect of  $NO_3^-$  ions on ATP-driven  $Na^+$  uptake by the PM vesicles from *T. viridis* is not related directly to the action of nitrate as a permeant anion but rather is conditioned by increased  $H^+$  conductivity of the membrane in the presence of nitrate and/or direct activation of the  $Na^+$ -ATPase by this ion.

**Acknowledgements:** This work was supported by the Russian Foundation for Basic Research (Grant 98-04-49433).

## References

- [1] Balnokin, Y.V. and Popova, L.G. (1994) FEBS Lett. 343, 61–64.
- [2] Balnokin, Y., Popova, L. and Gimmler, H. (1997) J. Plant Physiol. 150, 264–270.
- [3] Popova, L., Balnokin, Y., Dietz, K.-J. and Gimmler, H. (1998) FEBS Lett. 426, 161–164.
- [4] Stein, W.D. (1986) Transport and Diffusion across Cell Membranes, pp. 477–571, Academic Press, San Diego, CA.
- [5] Yu, X., Carrol, S., Rigaud, J.-L. and Inesi, G. (1993) Biophys. J. 64, 1232–1242.
- [6] Rasi-Caldogno, F., Pugliarella, M.C. and De Mishelis, M.I. (1987) Plant Physiol. 83, 994–1000.
- [7] Wada, M., Satoh, S.H., Kasamo, K. and Fujii, T. (1989) Plant Cell Physiol. 30, 923–928.
- [8] Shono, M., Wada, M. and Fujii, T. (1995) Plant Physiol. 108, 1615–1621.
- [9] Balnokin, Y.V., Popova, L.G. and Myasoedov, N.A. (1993) Plant Physiol. Biochem. 31, 159–168.
- [10] Apell, H.-J. and Bersch, B. (1987) Biochim. Biophys. Acta 903, 480–494.
- [11] Simpson, I.A. and Sonne, O. (1982) Anal. Biochem. 119, 424–427.
- [12] Popova, L.G. and Balnokin, Y.V. (1992) FEBS Lett. 309, 333–336.
- [13] Gallagher, S.R. and Leonard, R.T. (1982) Plant Physiol. 70, 1335–1340.