

# A REGULATORY MECHANISM FOR CO<sub>2</sub> ASSIMILATION IN PLANT PHOTOSYNTHESIS: ACTIVATION OF RIBULOSE-1,5-DIPHOSPHATE CARBOXYLASE BY FRUCTOSE 6-PHOSPHATE AND DEACTIVATION BY FRUCTOSE 1,6-DIPHOSPHATE


Bob B. BUCHANAN and Peter SCHÜRMANN

Department of Cell Physiology, University of California, Berkeley, California 94720, USA

Received 20 April 1972

One of the puzzling features of carbon assimilation in photosynthesis is the low affinity for CO<sub>2</sub> of

ribulose 1,5-diphosphate (RuDP) carboxylase in the initial CO<sub>2</sub> incorporation reaction (eq. 1).



Purified RuDP carboxylase shows a  $K_m$  for bicarbonate of about  $2 \times 10^{-2}$  M [1–4] and isolated intact chloroplasts carrying out complete photosynthesis show an apparent  $K_m$  for bicarbonate of about  $0.05 \times 10^{-2}$  M [5, 6]. We now report evidence for a new mechanism which increases the affinity of RuDP carboxylase for CO<sub>2</sub> (supplied as bicarbonate) *in vitro*




Fig. 1. Effect of fructose 6-phosphate and fructose 1,6-diphosphate on the activity of RuDP carboxylase. The reaction mixture contained the following (added in the order indicated): Tricine buffer, pH 7.5, 200 mM; EDTA, 0.06 mM; MgCl<sub>2</sub>, 1 mM; reduced glutathione, 5 mM; RuDP carboxylase (purified to homogeneity by a procedure based on that of Paulsen and Lane [3]), 20  $\mu$ g; fructose 6-phosphate and fructose 1,6-diphosphate, concentrations as shown; NaH<sup>14</sup>C<sub>2</sub>O<sub>3</sub> ( $5 \times 10^{-5}$  cpm  $\mu$ mole), 1 mM; and RuDP, 0.3 mM. Volume, 0.5 ml; temperature, 25°; reaction time, 10 min. The reaction, carried out in scintillation vials, was stopped with 0.05 ml 6 N HCl; samples were evaporated to dryness and the <sup>14</sup>C-phosphoglycerate (PGA) formed was determined in a scintillation counter. The control treatment gave 690 cpm PGA formed. PGA was identified as the reaction product in each of the treatments by thin-layer electrophoresis-chromatography [11].




Fig. 2. Effect of [MgCl<sub>2</sub>] on RuDP carboxylase activation by fructose 6-phosphate and deactivation by fructose 1,6-diphosphate. Except for varying the MgCl<sub>2</sub> concentration and adding the indicated fructose 6-phosphate and fructose diphosphate at 0.5 mM, conditions were as described for fig. 1.




Fig. 3. Effect of fructose 6-phosphate and fructose 1,6-diphosphate on the  $K_m$  of RuDP carboxylase for  $\text{HCO}_3^-$ . Except for varying the  $\text{HCO}_3^-$  concentration and adding the indicated fructose 6-phosphate and fructose diphosphate at 0.5 mM, conditions were as described for fig. 1.

and which may govern the activity of the enzyme *in vivo*.

Activity of homogeneous preparations of the carboxylase was increased up to 4-fold by 1 mM fructose 6-phosphate (F6P) (fig. 1). The F6P-activated carboxylase was fully deactivated by the addition of fructose 1,6-diphosphate (FDP). The activation by F6P (and deactivation by FDP) was independent of

the concentration of enzyme or RuDP and was observed at  $\text{MgCl}_2$  concentrations ranging from 0.5 to 10 mM (fig. 2). The sharpest activation was obtained at 1 mM  $\text{MgCl}_2$ .

Activation by F6P was markedly influenced by the concentration of bicarbonate: it was most pronounced at low concentrations and was maximal at 1 mM or less bicarbonate. 1 mM bicarbonate was also found to be optimal for complete photosynthesis by isolated intact chloroplasts.

Evidence that F6P acts by increasing the affinity of RuDP carboxylase for  $\text{CO}_2$  (added as bicarbonate) is shown in fig. 3. F6P lowered the  $K_m$  of the carboxylase for bicarbonate from  $2.5 \times 10^{-2}$  M to  $0.4 \times 10^{-2}$  M. The addition of FDP to F6P gave the expected return to the original  $K_m$  for bicarbonate.

These results indicate that chloroplast RuDP carboxylase is indeed subject to regulatory control (cf. [7, 8]) but that at physiological levels of  $\text{CO}_2$  the activity of the enzyme may be controlled by the relative concentrations of F6P and FDP. The levels of F6P and FDP may in turn be regulated by the ferredoxin-dependent FDPase system of chloroplasts that is actuated by light [9, 10]. A regulation




Fig. 4. Role of light in the regulation of fructose diphosphatase (FDPase) and RuDP carboxylase in spinach chloroplast.

mechanism for RuDP carboxylase for which there is now evidence in chloroplasts is summarized in fig. 4.

### Acknowledgements

We gratefully acknowledge the advice and support of Professor D.I. Arnon. This investigation was supported in part by Grant GB-30494X from the National Science Foundation to D.I.A.

### References

- [1] A. Weissbach, B.L. Horecker and J. Hurwitz, *J. Biol. Chem.* 218 (1956) 795.
- [2] E. Racker, *Arch. Biochem. Biophys.* 69 (1957) 300.
- [3] J.M. Paulsen and M.D. Lane, *Biochemistry* 5 (1966) 2350.
- [4] N. Kawashima and S.G. Wildman, *Ann. Rev. Plant Physiol.* 21 (1970) 325.
- [5] R.G. Jensen and J.A. Bassham, *Proc. Natl. Acad. Sci. U.S.* 56 (1966) 1095.
- [6] M. Gibbs, E. Latzko, R.G. Everson and W. Cockburn, in: *Harvesting The Sun*, eds. A. San Pietro, F.A. Green and T.J. Army (Academic Press, New York and London, 1967) p. 111.
- [7] T.A. Pedersen, M. Kirk and J.A. Bassham, *Physiol. Plant* 19 (1966) 219.
- [8] T. Sugiyama, N. Nakayama and T. Akazawa, *Arch. Biochem. Biophys.* 126 (1968) 737.
- [9] B.B. Buchanan, P.P. Kalberer and D.I. Arnon, *Biochem. Biophys. Res. Commun.* 29 (1967) 74.
- [10] B.B. Buchanan, P. Schürmann and P.P. Kalberer, *J. Biol. Chem.* 246 (1971) 5952.
- [11] P. Schürmann, B.B. Buchanan and D.I. Arnon, *Biochim. Biophys. Acta* 267 (1972) 111.