

One- and two-dimensional 90.5-MHz ^{13}C -NMR spectroscopy of the *N*-linked triantennary oligosaccharide units of calf fetuin

Elisha Berman* and Peter Bendel

Department of Organic Chemistry, The Weizmann Institute of Science, Rehovot 76100 and School of Chemistry, Tel Aviv University, Ramat Aviv, Israel

Received 13 May 1986; revised version received 9 June 1986

Complete assignments of all anomeric resonances in the proton and carbon spectra of the *N*-linked oligosaccharide units of fetuin were made using one- and two-dimensional NMR spectroscopy. We are able to confirm the presence of microheterogeneity in the *N*-acetylneuraminic acid linkages to the galactose residues and the presence of a unique triantennary structure which carries a side chain: NeuAc α (1–3)Gal β (1–3)GlcNAc β (1–4). Anomeric carbon chemical shifts changes resulting from long-range conformational effects were observed.

Triantennary oligosaccharide two-dimensional NMR spectroscopy ^{13}C -NMR Fetuin
Structural microheterogeneity (Calf)

1. INTRODUCTION

Calf fetuin is a glycoprotein found in fetal calf serum [1], with three of its six carbohydrate side chains known to be *N*-linked to an asparagine residue. Two different isomeric side chain structures were proposed for these oligosaccharides [2,3]; but recently it was shown that only one of the proposed triantennary structures is correct [4–6]. Moreover, evidence has been accumulated for the presence of β (1–3) galactose residue directly linked to *N*-acetylglucosamine [5–7], as well as a ratio of ~1:1 between the NeuAc α (2–3) and NeuAc α (2–6) residues [5].

A study of the isolated fetuin triantennary glycopeptides using high magnetic field ^{13}C -NMR

and 2D-NMR spectroscopy is reported. The results provide an additional proof for the presence of Gal β (1–3)GlcNAc β (1–4) antenna in calf fetuin and correlates between the ^{13}C -NMR and ^1H -NMR chemical shifts of the anomeric resonances through heteronuclear shift-correlated NMR spectroscopy [8].

2. MATERIALS AND METHODS

An *N*-linked glycopeptide fraction of calf fetuin and its neuraminidase-treated glycopeptide fraction were isolated and purified from fetuin as described [2,3,5].

The 90.5-MHz ^{13}C -NMR (360 MHz for ^1H -NMR) spectra were recorded on a Bruker AM-360 spectrometer at ambient probe temperature. The fetuin glycopeptide sample was prepared as a 70 mM solution in D_2O at neutral pH (10 mm NMR tube). The asialo sample, from which the sialic acid residues were removed [5], was prepared in a similar manner (50 mM solution) in a 5 mm NMR tube. The heteronuclear shift-correlated 2D-

* Incumbent of The Charles H. Revson Foundation Career Development Chair

Abbreviations: 2D, two-dimensional; Man, D-mannopyranose; Gal, D-galactopyranose; GlcNAc, *N*-acetylglucosamine; NeuAc, *N*-acetylneuraminic acid




Fig. 1. 2D chemical shift-correlated NMR contour plot (5.3–4.1 ppm for proton and 109–95.5 ppm for carbon) of the *N*-linked glycopeptides derived from calf fetuin. Residue numbering is given in the scheme shown at the bottom of table 1.

relaxation rates and overlap of closely spaced resonances arising from the heterogeneity in the substitution by the sialic acid residues [5]. No carbon-proton connectivity, other than the connectivity for one of the α -mannose residues [9–11], was observed in the anomeric proton chemical shift region (>5.1 ppm) which correspond to α -galactose residue [12,13]. Thus it was concluded that all the galactose residues must have a β -anomeric configuration. In addition, two proton connectivities were observed for the anomeric carbon resonance of the $\beta(1-4)$ -linked *N*-acetylglucosamine residue. Although it may arise from some long-range substituent chemical shift effects, it is more likely to be a result of different galactose substitutions [i.e. $\beta(1-4)$ vs $\beta(1-3)$]. The problem was resolved when the ^{13}C -NMR spectrum of the asialofetuin glycopeptide was examined, where any long-range effects due to different sialic acid substitutions were removed from the spectrum (fig. 2).

The 90.5-MHz ^{13}C -NMR spectrum of the anomeric carbons region of the fetuin triantennary units from which the α -linked sialic acid was removed enzymatically [5] is shown in fig. 2. The assignments of the anomeric carbon resonances are based on an extensive chemical shift comparisons with model compounds and the analysis of the ^{13}C -NMR spectra of the corresponding sialyl and agalactosyl derivatives [5]. Unlike the lower




Fig. 2. 90.5-MHz ^{13}C -NMR spectrum, showing the anomeric region of the asialylated *N*-linked glycopeptides derived from calf fetuin. Residue numbering corresponds to the scheme shown at the bottom of table 1.

magnetic field study [5], the improved spectral resolution enabled the observation of four anomeric carbon resonances which are associated with β -galactose residues. The smaller and most downfield resonance, at 104.82 ppm, was assigned previously to a $\text{Gal}\beta(1-3)$ residue, (Gal-8^*) [5]. The other, closely spaced, resonances at 104.30, 104.27 and a shoulder peak at 104.23 ppm, were assigned to residues 6, 6' and 8 (see scheme in table 1). Integration of the carbon resonances appearing in the anomeric region, excluding the GlcNAc-1 residue, has indicated the presence of 10 sugar residues, as expected.

Two of the three resonances corresponding to Gal-6 , $-6'$ and -8 , are of equal intensity while the intensity of the third resonance is reduced by an amount approximately equal to the signal intensity of the peak assigned to Gal-8^* . Therefore, one unique antenna in the triantennary structure must be associated with the $\beta(1-3)$ -linked galactose. The carbon resonance of GlcNAc-7 is composed of two components, one at 102.84 ($\sim 1/3$ carbon) and the other at 102.75 ppm ($\sim 2/3$ carbon), so that it must be the GlcNAc-7 residue which is substituted by $\text{Gal}\beta(1-3)$ as previously proposed [5]. Therefore, on average, one out of the three *N*-linked triantennary structures in calf fetuin carries this type of side chain.

4. CONCLUSIONS

The heteronuclear shift-correlated experiment confirmed the ^{13}C -NMR anomeric chemical shift assignments for the proposed triantennary structure of calf fetuin [5] via the correlation to the well-established proton chemical shift assignments [9–13]. This is the first report which directly correlates between these chemical shifts for the anomeric part of a triantennary oligosaccharide structure. It was used to establish the presence of a $\text{Gal}\beta(1-3)$ residue in the *N*-linked side chain of calf fetuin.

Some long-range chemical shift effects were observed in the ^{13}C -NMR spectrum (fig.2), which are most likely conformational in origin. The two carbon peaks corresponding to GlcNAc-5 and $-5'$ in the sialylated glycopeptide (not shown) yielded a sharp resonance, 100.81 ppm, following the removal of the sialic residues (fig.2). In general, most of the anomeric resonances corresponding to the inner core *N*-acetylglucosamine and mannose residues undergo measurable chemical shift changes upon desialylation as observed for smaller model oligosaccharides [14,15].

REFERENCES

- [1] Graham, E.R.B. (1972) in: *Glycoproteins Part A* (Gottschalk, A. ed.) vol.5A, pp.717–731, Elsevier, Amsterdam, New York.
- [2] Nilsson, B., Norden, N.E. and Svensson, S. (1979) *J. Biol. Chem.* 254, 4545–4553.
- [3] Baenziger, J.U. and Fiete, D. (1979) *J. Biol. Chem.* 254, 789–795.
- [4] Krusius, T. and Finne, J. (1978) *Carbohydr. Res.* 90, 203–214.
- [5] Berman, E. (1986) *Carbohydr. Res.* 150, in press.
- [6] Van Halbeek, H., Rijkse, I., Van Beek, W.P., Kamerling, J.P. and Vliegthart, J.F.G. (1985) *Proc. VIII Int. Symp. Glycoconjugates* 1, 120–121.
- [7] Graham, E.R.B. (1961) *Aust. J. Sci.* 24, 140–141.
- [8] Bax, A., Egan, W. and Kovac, P. (1984) *J. Carbohydr. Chem.* 3, 593–611.
- [9] Dorland, L., Haverkamp, J., Vliegthart, J.F.G., Strecker, G., Michalski, J.-C., Fournet, B., Spik, G. and Montreuil, J. (1978) *Eur. J. Biochem.* 87, 323–329.
- [10] Endo, M., Suzuki, K., Schmid, K., Fournet, B., Karamanos, Y., Montreuil, J., Dorland, L., Van Halbeek, H. and Vliegthart, J.F.G. (1982) *J. Biol. Chem.* 257, 8755–8760.
- [11] Bhattacharyya, S.N., Lynn, W.S., Dabrowski, J., Trouner, K. and Hull, W.E. (1984) *Arch. Biochem. Biophys.* 231, 72–85.
- [12] Falk, K.-E., Joval, P.-A. and Winyard, P. (1982) *Acta Chem. Scand.* B36, 558–560.
- [13] Dorland, L., Van Halbeek, H. and Vliegthart, J.F.G. (1984) *Biochem. Biophys. Res. Commun.* 122, 859–866.
- [14] Berman, E. (1983) *Carbohydr. Res.* 118, 9–20.
- [15] Berman, E. (1984) *Biochemistry* 23, 3754–3759.