

Identification of the structures of multiple subspecies of protein kinase C expressed in rat brain

Ushio Kikkawa, Yoshitaka Ono⁺, Kouji Ogita, Tomoko Fujii⁺, Yoshinori Asaoka, Kazuo Sekiguchi, Yoshiyuki Kosaka, Koichi Igarashi⁺ and Yasutomi Nishizuka

Department of Biochemistry, Kobe University School of Medicine, Kobe 650 and ⁺Biotechnology Laboratories, Central Research Division, Takeda Chemical Industries, Osaka 532, Japan

Received 22 April 1987

Rat brain protein kinase C purified to apparent homogeneity [(1986) *Biochem. Biophys. Res. Commun.* 135, 636–643] was resolved into three distinct fractions, type I, II and III, upon chromatography on a hydroxyapatite column connected to high-performance liquid chromatography. Comparison of each fraction with the four subspecies of protein kinase C, that were separately expressed in COS cells transfected by the respective cDNAs, α , β I, β II and γ , identified the primary structures of these three fractions of protein kinase C. Type I corresponded to the enzyme encoded by the γ -sequence; type II was a mixture of the two subspecies determined by the β I- and β II-sequences; and type III had the structure encoded by the α -sequence. The structures and properties of these subspecies of protein kinase C were similar to each other.

Protein kinase C; cDNA; (COS cell)

1. INTRODUCTION

Protein kinase C is now generally accepted as playing crucial roles in cell surface signal transduction [1]. The enzyme was thought to be a single entity, but recent analysis of its cDNA clones indicates that in the mammalian brain there are multiple subspecies of this enzyme with closely related structures [2–9]. Comparison of their predicted amino acid sequences reveals that at least four subspecies of protein kinase C may exist in this tissue. Since the sequence analysis has been carried out independently in several laboratories [2–9], different nomenclatures have been proposed for the various cDNA clones as given in table 1. The three cDNA clones, α , β and γ , ob-

tained from the bovine and human brain libraries by Coussens et al. [3] are shown to be encoded by distinct chromosomes, whereas the two cDNA clones, type I and II, obtained from the rat brain library are derived from alternative splicing and differ from each other in the carboxyl-terminal region of only about 50 amino acid residues as described earlier [5]. Thus, the nomenclature of α , β I, β II and γ will be used hereafter for the four cDNA clones having very similar but different nucleotide sequences (table 1). An apparently homogeneous preparation of rat brain protein kinase C, on the other hand, is shown to be resolved further into three fractions, type I, II and III, upon chromatography on a hydroxyapatite column [10,11]. The fraction of type II is a mixture of the two subspecies of protein kinase C encoded by β I- and β II-sequences [11]. Extending these studies, we wish to describe briefly here the identification of the structures of type I and type III by

Correspondence address: U. Kikkawa, Department of Biochemistry, Kobe University School of Medicine, Kobe 650, Japan

Table 1
cDNA clones of protein kinase C

	cDNAs				References
Bovine (human) brain	α (672)	—	β (673)	γ (697?)	2,3
Rat brain	X ^a (?)	type I (671)	type II (673)	—	4,5
Rat brain	—	PKC III (?)	PKC II (673)	PKC I (697)	6
Rat brain	—	RP41 (?)	—	—	7 ^b
Rabbit brain	γ (672)	β (671)	α (673)	—	9
Nomenclature used here for cDNA clones	α	β I	β II	γ	
Subfractions of brain enzymes	type III	type II	type I		

^a Analysis of partial amino acid sequence of the fragments of protein kinase C predicted the existence of this clone

^b A cDNA clone designated as RP16, which apparently differs from all of the cDNA clones described above, was also reported

Numbers in parentheses indicate the length of deduced amino acid sequences of each clone

comparison with the subspecies expressed in COS cells transfected by each of the rat brain cDNA clones, α , β I, β II and γ .

2. MATERIALS AND METHODS

2.1. Assay of protein kinase C

The enzyme was assayed by measuring the incorporation of ³²P_i into calf thymus H₁ histone from [γ -³²P]ATP in the presence of 8 μ g/ml phosphatidylserine, 0.8 μ g/ml diolein, and 0.5 mM Ca²⁺ under the conditions specified in [12]. Basal activity was measured in the presence of 0.5 mM EGTA instead of phosphatidylserine, diolein and Ca²⁺.

2.2. Isolation of α , β I, β II and γ cDNA clones

Phage cDNA library was constructed using rat brain poly(A)⁺ RNA with λ gt10 vector system, and β I and β II cDNA clones were isolated as in [5]. Both α and γ cDNA clones of rat brain protein kinase C were obtained by sequential plaque

screening according to Maniatis et al. [13] using synthetic 26-base oligonucleotide probes, which corresponded to amino acid residues from 162 to 170 and 334 to 342 of bovine α cDNA [3], and those from 165 to 173 and 338 to 346 of rat γ cDNA [6], respectively. Although cDNA clones encoding the α type of bovine, human, and rabbit protein kinase C were isolated [2,9], the clone of this type has not yet been obtained from the rat tissue library. Using the oligonucleotide probes for the α type, a cDNA clone containing a 3.2 kb insert was obtained. This clone, designated λ CKR α 5, encodes the whole coding sequences of 672 amino acids together with the 5'- and 3'-noncoding regions. The coding sequence was very similar to the α cDNA clones of bovine and rabbit brain [2,9]. The γ cDNA clone designated λ CKR γ 1 of 3.1 kb includes the full coding sequence with non-coding regions on both sides. The nucleotide sequence of this clone was identical with that of PKC I described by Knopf et al. [6]. The detailed procedures of isolation and the precise comparison of

the nucleotide sequences of rat α , β I and β II and γ cDNAs will be described elsewhere.

2.3. Construction of expression plasmids

The expression plasmid was constructed from Okayama and Berg vectors [14], which was essentially similar to pcDL1 reported by Ebina et al. [15]. The *Eco*RI-*Eco*RI cDNA fragment of isolated clones was inserted into the *Eco*RI site of the expression plasmid as described [11].

2.4. Expression in COS cells

Fresh monolayers of COS 7 cells with Dulbecco's modified Eagle medium containing 5% fetal calf serum were transfected with the plasmid DNA using the calcium phosphate co-precipitation technique [16,17].

3. RESULTS AND DISCUSSION

3.1. Resolution of protein kinase C into three fractions

Protein kinase C was purified from the cytosol of rat brain by DE-52 (Whatman), threonine-Sepharose, and TSK phenyl-5PW (Toyo Soda) column chromatographies as described previously [12]. This enzyme was resolved further into three distinct fractions upon chromatography on a hydroxyapatite column connected to high-performance liquid chromatography, Pharmacia FPLC system. The TSK phenyl-5PW fraction of protein kinase C was diluted with an equal volume of 20 mM potassium phosphate buffer at pH 7.5, containing 0.5 mM EGTA, 0.5 mM EDTA, 10% glycerol, and 10 mM 2-mercaptoethanol (buffer A), and was applied to a packed hydroxyapatite column (0.78 \times 10 cm, Koken, Tokyo, type S) equilibrated with buffer A. Protein kinase C was then eluted by application of a linear concentration gradient of potassium phosphate buffer at pH 7.5 (20–250 mM) in 84 ml buffer A. Fractions of 1 ml each were collected. All procedures were carried out at 0–4°C. Three distinct fractions of protein kinase C were obtained; type I (fractions 24–28), type II (fractions 31–39), and type III (fractions 50–57) as shown in fig.1A. The three enzyme fractions did not appear to be simple artifacts during the purification procedures, and the elution profile was highly reproducible. Proteolytic modification may presumably be excluded, since the extraction




Fig.1. Hydroxyapatite column chromatography of several subspecies of protein kinase C. A purified preparation of rat brain protein kinase C and the enzymes newly expressed in the COS 7 cells were applied to hydroxyapatite column chromatography, and enzymatic activity was assayed as described in section 2. (A) Rat brain protein kinase C; (B–D) protein kinase C from the COS 7 cells transfected with α , β II, and γ cDNA, respectively; (E) protein kinase C from untransfected control COS 7 cells. (●—●) Protein kinase activity in the presence of phosphatidylserine, diolein, and CaCl_2 ; (○—○) protein kinase activity in the presence of EGTA instead of phosphatidylserine, diolein, and CaCl_2 ; (—) potassium phosphate.

buffer contained high concentrations of both EGTA and EDTA to remove Ca^{2+} . The enzymes in these fractions each showed a single band with roughly an identical molecular mass of 80 kDa upon SDS-polyacrylamide gel electrophoresis. The three enzymes reacted equally with a monoclonal

antibody raised to a common polypeptide sequence present in the four subspecies of protein kinase C. Type III enzyme normally appeared as a broad peak, and heterogeneity of this enzyme fraction is unknown at present.

3.2. Expression of four subspecies of protein kinase C

The cDNAs encoding α , β I and β II and γ subspecies of protein kinase C were each inserted in the expression plasmid, and transfected separately to COS 7 cells as described in section 2. The enzymes expressed in the COS 7 cells were partially purified under the conditions described below. The COS 7 cells were homogenized by sonication in 20 mM Tris-HCl at pH 7.5, containing 0.25 M sucrose, 10 mM EGTA, 2 mM EDTA and 20 μ g/ml leupeptin. The homogenate was centrifuged for 60 min at $100\,000 \times g$. The supernatant was diluted with 6 vols of 20 mM Tris-HCl at pH 7.5, containing 0.5 mM EGTA, 0.5 mM EDTA and 10 mM 2-mercaptoethanol (buffer B), and was applied to a Mono Q column (1 \times 10 cm, Pharmacia HR 10/10) equilibrated with buffer B. The enzyme was eluted by application of a linear concentration gradient of NaCl (0–0.6 M) in 164 ml buffer B. Fractions of 4 ml each were collected, and the fractions possessing protein kinase C activity were pooled and subjected to hydroxyapatite column chromatography as described above. The activity of protein kinase C from the COS 7 cells transfected by each cDNA was usually 3–10-times higher than that of the control cells. When equivalent amounts of the enzyme from the Mono Q column was subjected to hydroxyapatite column chromatography, several different patterns of the enzyme were obtained as shown in fig.1B–D. The COS 7 cells transfected with the plasmid containing α cDNA showed one major peak (fractions 47–60) (fig.1B), whereas the control enzyme preparation from untransfected COS 7 cells showed only a minor, broad peak at the same position (fig.1E). Fig.1C shows the enzyme pattern obtained from the β II cDNA-transfected cells. A new major peak appeared (fractions 29–38) that eluted at the position corresponding to type II protein kinase C. A similar result was obtained for the COS 7 cells transfected by β I cDNA, and both β I and β II enzymes expressed in COS 7 cells were indistinguishable from one another. In contrast,

when COS 7 cells were transfected with the plasmid containing γ cDNA, another new peak of protein kinase appeared which eluted at the position corresponding to the type I enzyme (fractions 24–30) (fig.1D). All protein kinases thus obtained showed typical characteristics of protein kinase C, being dependent on the simultaneous presence of Ca^{2+} , phospholipid, and diacylglycerol for full enzymatic activity. The results presented above together with those described in a preceding report [11] indicate that the three types of protein kinase C isolated from the rat brain cytosol, type I, II and III, closely resemble the structures of γ , β I and β II and α -sequences, respectively, that have been predicted by analysis of the cDNA clones. However, the possibility may not be excluded that additional species of protein kinase C exist in these three fractions that are separated by hydroxyapatite column chromatography. There may also be some posttranslational modification of the enzyme, since the amino-terminal of the enzyme is not detected by Edman degradation. The amino acid sequence analysis of each species of protein kinase C is needed to establish the exact relationship between the gene and protein molecule. More details of the properties and comparison of these different fractions of protein kinase C will be described elsewhere.

ACKNOWLEDGEMENTS

We thank Drs Y. Sugino and A. Kakinuma for encouragement, and Ms S. Nishiyama for secretarial assistance. The research in the Department of Biochemistry, Kobe University School of Medicine was supported in part by research grants from the Scientific Research Fund of the Ministry of Education, Science and Culture, Japan; Muscular Dystrophy Association; Yamanouchi Foundation for Research on Metabolic Disorders; Merck Sharp & Dohme Research Laboratories; Biotechnology Laboratories of Takeda Chemical Industries; Ajinomoto Central Research Laboratories; and Meiji Institute of Health Sciences.

REFERENCES

- [1] Nishizuka, Y. (1986) Science 233, 305–312.

- [2] Parker, P.J., Coussens, L., Totty, N., Rhee, L., Young, S., Chen, E., Stabel, S., Waterfield, M.D. and Ullrich, A. (1986) *Science* 233, 853–859.
- [3] Coussens, L., Parker, P.J., Rhee, L., Yang-Feng, T.L., Chen, E., Waterfield, M.D., Francke, U. and Ullrich, A. (1986) *Science* 233, 859–866.
- [4] Ono, Y., Kurokawa, T., Kawahara, K., Nishimura, O., Marumoto, R., Igarashi, K., Sugino, Y., Kikkawa, U., Ogita, K. and Nishizuka, Y. (1986) *FEBS Lett.* 203, 111–115.
- [5] Ono, Y., Kurokawa, T., Fujii, T., Kawahara, K., Igarashi, K., Kikkawa, U., Ogita, K. and Nishizuka, Y. (1986) *FEBS Lett.* 206, 347–352.
- [6] Knopf, J.L., Lee, M.-H., Sultzman, L.A., Kriz, R.W., Loomis, C.R., Hewick, R.M. and Bell, R.M. (1986) *Cell* 46, 491–502.
- [7] Housey, G.M., O'Brian, C.A., Johnson, M.D., Kirschmeier, P. and Weinstein, I.B. (1987) *Proc. Natl. Acad. Sci. USA* 84, 1065–1069.
- [8] Makowske, M., Birnbaum, M.J., Ballester, R. and Rosen, O.M. (1986) *J. Biol. Chem.* 261, 13389–13392.
- [9] Ohno, S., Kawasaki, H., Imajoh, T., Suzuki, K., Inagaki, M., Yokokura, H., Sakoh, T. and Hidaka, H. (1987) *Nature* 325, 161–166.
- [10] Huang, K.-P., Nakabayashi, H. and Huang, F.L. (1986) *Proc. Natl. Acad. Sci. USA* 83, 8535–8539.
- [11] Ono, Y., Kikkawa, U., Ogita, K., Fujii, T., Kurokawa, T., Asaoka, Y., Sekiguchi, K., Ase, K., Igarashi, K. and Nishizuka, Y. (1987) *Science*, in press.
- [12] Kikkawa, U., Go, M., Koumoto, J. and Nishizuka, Y. (1986) *Biochem. Biophys. Res. Commun.* 135, 636–643.
- [13] Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) in: *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- [14] Okayama, H. and Berg, P. (1983) *Mol. Cell. Biol.* 3, 280–289.
- [15] Ebina, Y., Edery, M., Ellis, L., Standring, D., Beaudoin, J., Roth, R.A. and Rutter, W.J. (1985) *Proc. Natl. Acad. Sci. USA* 82, 8014–8018.
- [16] Reddy, E.P., Smith, M.J. and Srinivasan, A. (1983) *Proc. Natl. Acad. Sci. USA* 80, 3623–3627.
- [17] Graham, F.L. and Van der Eb, A.J. (1973) *Virology* 54, 536–539.