

Activation of chloride channels in the plasmalemma of *Nitella syncarpa* by inositol 1,4,5-trisphosphate

O.M. Zherelova

Institute of Biological Physics, Academy of Sciences of the USSR, Pushchino, Moscow Region 142292, USSR

Received 16 March 1989

The introduction of $1.5 \times 10^{-8} - 10^{-6}$ M inositol 1,4,5-trisphosphate (IP_3) into the bathing solution of *Nitella syncarpa* cells induced: (i) an increase in amplitude of the inward Ca^{2+} current; (ii) the appearance of an inward Cl^- current; (iii) a shift in the threshold for activation of the Cl^- channels on hyperpolarization of the membrane to more positive values. These facts suggest that IP_3 participates in the causing of changes in the intracellular free Ca^{2+} concentration and, hence, in the regulation of ionic channels in *Nitella syncarpa* plasmalemma.

Inositol 1,4,5-trisphosphate; Ca^{2+} ; Ionic channel; (Characeae, *Nitella syncarpa*)

1. INTRODUCTION

In recent years the role of membrane phospholipid exchange in the regulation of intracellular Ca^{2+} has been widely studied [1,2]. Of particular interest in this regard is the exchange of minor membrane lipids, viz. phosphoinositides [3,4]. The products of inositol 1,4-bisphosphate (IP_2) hydrolysis, IP_3 and diacylglycerol, are secondary messengers [2]. Diacylglycerol triggers phosphorylation of regulatory proteins by activation of protein kinase C, whereas IP_3 participates in the release of Ca^{2+} from intracellular stores thus initiating a cascade of calcium-regulated reactions in the cell. In particular, intracellular calcium acts in the regulation of various types of ionic channels in membrane electrogenesis [5-7].

In membranes from a variety of plant cells, phospholipid exchange components and protein kinase C have been detected [8-10]. Moreover, IP_3 has been shown to be involved in the release of Ca^{2+} accumulated in protoplasts and microsomes [11,12].

The present study was aimed at clarifying whether the product of phosphoinositide hydrolysis, IP_3 , participates in controlling activation of ionic channels in the plasmalemma of *Nitella syncarpa*.

2. MATERIALS AND METHODS

Experiments were performed on perfused tonoplast-free cells of *N. syncarpa* under voltage-clamp conditions on the plasmalemma. The technique has been detailed in [14]. For intracellular and extracellular perfusion the following respective solutions (mM) were used: (i) 1 EGTA, 1.5 MgCl_2 , 15 KCl, 10 Hepes-Tris, 0.5 ATP, 280 sucrose; $[\text{Ca}^{2+}]_{\text{ext}} 10^{-8}$ (pH 7.2); (ii) 1 CaCl_2 , 5 Hepes-Tris, 220 sucrose, pH 7.2.




Fig.1. Increase in amplitude of the inward Ca^{2+} current at 50 mV (a) potential and development of the inward Cl^- current (b) in response to injection of 1.5×10^{-8} M IP_3 into the bathing solution. 2×10^{-5} M ethacrinic acid blocked the Cl^- current.

Correspondence address: O.M. Zherelova, Institute of Biological Physics, Academy of Sciences of the USSR, Pushchino, Moscow Region 142292, USSR




Fig.2. Shift of activation threshold for Cl^- channels on hyperpolarization under the influence of 1.5×10^{-8} M IP_3 introduced into the bathing solution. 1, control; 2, after addition of 1.5×10^{-8} M IP_3 .

D-*myo*-Inositol 1,4,5-trisphosphate (Amersham) and ethacrynic acid (Sigma) were used.

3. RESULTS

Incubation of *Nitella* cells in an IP_3 -containing solution led to an increase in amplitude of the Ca^{2+} current (I_{Ca}) developed on activation of the channels by depolarization. After a definite time period (10–20 min, depending on the IP_3 concentration), spontaneous generation of an inward current took place whose amplitude became larger with time. This current was easily blocked by ethacrynic acid, as reported for the case of *N. obtusa* cells [16]. We identified the current to be of chloride. Typical recordings of inward potential-dependent Ca^{2+} currents and Cl^- current are shown in fig.1.

The second set of experiments was carried out with Cl^- currents developed on shifting the fixed potential on the membrane toward hyperpolariza-

tion [15]. Under standard experimental conditions ($[\text{Ca}^{2+}]_{\text{ext}} 10^{-8}$ M) the threshold for Cl^- current generation was observed to be at -240 mV. When cells were incubated in an IP_3 -containing solution, the threshold Cl^- current on hyperpolarization shifted markedly to positive values (fig.2).

4. DISCUSSION

As shown earlier [17], an increase in concentration of intracellular Ca^{2+} caused activation of potential-dependent Ca^{2+} channels in *N. obtusa* plasmalemma. The increase in I_{Ca} resulting from introduction of IP_3 into the bathing solution indicated that an increase occurred in intracellular Ca^{2+} level, most probably at the expense of the release from submembrane stores.

Intracellular Ca^{2+} also acts as a transmitter, capable of activating Cl^- channels in *N. obtusa* [7,14]. Ca^{2+} released from submembrane stores and the local increase in Ca^{2+} concentration in the perimembrane layer might be the factors causing the development of a Cl^- current, as has been shown for oocytes [18].

The change in the threshold for activation of Cl^- channels of *Nitella* plasmalemma during hyperpolarization for incubation of cells in an IP_3 solution suggests that the Cl^- channels in *Nitella* cells are under the control of intracellular Ca^{2+} .

Thus, the data obtained on the ability of IP_3 to produce an increase in I_{Ca} and to induce the appearance of a Cl^- current through the *Nitella* plasmalemma, as well as its effect of shifting the threshold for activation of hyperpolarization-




Fig.3. Reversible ethacrynic-acid-induced block of Cl^- currents on hyperpolarization (0 to -130 mV). The concentration of IP_3 was 10^{-6} M. \uparrow , addition of 2×10^{-5} M ethacrynic acid; \downarrow , washing from ethacrynic acid.

dependent Cl^- channels, constitute evidence in support of the contention that IP_3 promotes mobilization of intracellular Ca^{2+} .

The exchange of minor membrane lipids, namely phosphoinositides, probably plays an essential role in the regulation of ionic channels in the plasmalemma of Characeae.

Acknowledgements: The author expresses her gratitude to N. Agafonova for help in performing the present study and to M. Ilyina for technical assistance.

REFERENCES

- [1] Berridge, M.J. and Irvin, R.F. (1984) *Nature* 312, 315-321.
- [2] Berridge, M.J. (1986) *J. Exp. Biol.* 124, 323-325.
- [3] Kikkawa, U., Kaibuchi, K., Takai Y. and Nishizuka, Y. (1985) *Phospholipids and Cellular Regulation*, vol. II, 112-121.
- [4] Kucherenko, N.E. and Blum, Ya.B. (1987) *Ukr. Biokhim. Zh.* 58, 86-101.
- [5] Meech, R.W. (1978) *Annu. Rev. Biophys. Bioeng.* 7, 1-8.
- [6] Kazachenko, V.N. and Geletyuk, V.I. (1984b) *Biol. Membrany* 1, 1253-1265.
- [7] Lunevsky, V.L., Zherelova, O.M., Vostrikov, I.Y. and Berestovsky, G.N. (1983) *J. Membrane Biol.* 72, 43-58.
- [8] Wheeler, J.J. and Boss, W.F. (1987) *Plant Physiol.* 85, 389-392.
- [9] Heim, S., Bauleke, A., Wylegalla, C. and Wagner, K.G. (1987) *Plant Sci.* 49, 159-165.
- [10] Morse, M.J., Crain, R.C. and Satter, R.L. (1987) *Plant Physiol.* 83, 640-644.
- [11] Rincon, M. and Boss, W.F. (1987) *Plant Physiol.* 83, 395-398.
- [12] Drobak, B.K. and Ferguson, I.B. (1985) *Biochem. Biophys. Res. Commun.* 130, 1241-1246.
- [13] Schumaker, K.S. and Sze, H. (1987) *J. Biol. Chem.* 262, 3944-3946.
- [14] Kataev, A.A., Zherelova, O.M. and Berestovsky, G.N. (1984) *Gen. Physiol. Biophys.* 3, 447-462.
- [15] Tyerman, S.D., Findlay, G.P. and Paterson, G.J. (1986) *Membrane Biol.* 89, 139-152.
- [16] Zherelova, O.M., Berestovsky, G.N. and Natochin, Yu.V. (1984) *Biol. Membrany* 1, 728-733.
- [17] Zherelova, O.M., Kataev, A.A. Berestovsky, G.N. (1987) *Biophysica* 32, 348-350.
- [18] Parker, I. and Miledi, F.R.S. (1987) *Proc. R. Soc. Lond.* B231, 27-36.
- [19] Chesnoy-Marchais, D. (1983) *J. Physiol.* 342, 277-308.