

Functional involvement of calcium in the homologous up-regulation of the 1,25-dihydroxyvitamin D₃ receptor in osteoblast-like cells

J.P.T.M. van Leeuwen, J.C. Birkenhäger, C.J. Buurman, J.P. Schilte and H.A.P. Pols

Department of Internal Medicine III, Erasmus University Medical School, Rotterdam, The Netherlands

Received 3 July 1990; revised version received 16 July 1990

In several cell types 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) causes up-regulation of its receptor. The present study demonstrates that in the osteoblast-like cell line UMR 106 this up-regulation is inhibited by two different calcium channel blockers (nitrendipine, verapamil). Also with chelating extracellular calcium (EGTA) and by inhibition of calcium release from intracellular stores (TMB-8) comparable results were obtained. These findings indicate that calcium is functionally involved in this cellular response to the steroid hormone 1,25(OH)₂D₃. Moreover, data obtained with EGTA show that the 1,25(OH)₂D₃ receptor level is closely regulated by the extracellular calcium concentration.

1,25-Dihydroxyvitamin D₃; Calcium; 1,25(OH)₂D₃ receptor up-regulation

1. INTRODUCTION

1,25-Dihydroxyvitamin D₃ (1,25(OH)₂D₃) is an important regulator of calcium homeostasis and bone metabolism. It has been shown that 1,25(OH)₂D₃ causes homologous up-regulation of its receptor in the osteoblast-like cell lines ROS 17/2 [1] and UMR 106 [2], and in fibroblasts and cloned kidney cells [3].

The occupied 1,25(OH)₂D₃ receptor is thought to act direct at the genome. However, in various cellular systems 1,25(OH)₂D₃ has been shown to stimulate calcium uptake independent of de novo RNA and protein synthesis [4–6], and genome independent effects on membrane potential have been reported [7]. Also, recently it has been demonstrated that 1,25(OH)₂D₃ causes a rapid (less than 30 s) rise of the intracellular ionized calcium concentration ([Ca²⁺]_i) in isolated mouse osteoblasts [8]. The 1,25(OH)₂D₃-induced rise of the [Ca²⁺]_i has been shown to be inhibited by the calcium channel blockers nifedipine and verapamil, by chelating extracellular calcium with EGTA and to be reduced by blocking release from intracellular stores by 8-(diethylamino)octyl 3,4,5-trimethoxybenzoate HCl (TMB-8) [8]. In the osteoblast-like cell line ROS 17/2.8 1,25(OH)₂D₃ modulates dihydropyridine-sensitive L-type calcium channels [9]. Moreover, in keratinocytes and rat enterocytes 1,25(OH)₂D₃ increases the generation of inositol 1,4,5-trisphosphate indicating an effect on calcium release from intracellular stores [10,11].

Based on these data it is conceivable that calcium acts as an intracellular messenger in the regulation of cellular responses by 1,25(OH)₂D₃. However, as yet a functional role for calcium has to be demonstrated.

In the present study we have investigated whether calcium is functionally involved in 1,25(OH)₂D₃-induced up-regulation of its receptor in the osteoblast-like cell line UMR 106.

2. MATERIALS AND METHODS

2.1. Materials

Verapamil and EGTA were obtained from Sigma, St. Louis, USA, TMB-8 from Aldrich Chemical Co., Bruxelles, Belgium. Nitrendipine was a generous gift from Dr B. Garthoff of Bayer AG, Wuppertal, FRG. [23,24-³H]1,25(OH)₂D₃ (90 Ci/mmol) was purchased from Amersham International, Amersham, UK, and non-radioactive 1,25(OH)₂D₃ was generously provided by LEO Pharmaceuticals BV, Weesp, The Netherlands.

2.2. Culture and treatment of the cells

UMR 106 cells were cultured as described previously [12]. The cells were incubated for 4 h with or without 1,25(OH)₂D₃ in the absence or presence of nitrendipine, verapamil, EGTA, or TMB-8.

2.3. Preparation of cell extracts and 1,25(OH)₂D₃ binding assay

For single point assays, conditions were employed which were previously shown to provide valid estimates of total VDR content in cytosolic extracts [2]. The protein concentration was measured according to the method of Bradford [13].

2.4. Data analysis

Data presented are means ± SD of triplicate determinations of at least 2 different experiments, i.e. at least 6 replicates. Interactions between 1,25(OH)₂D₃ and calcium antagonists were evaluated using an analysis of variance for 2-way factorial design [14]. Other statistical analyses were done by Student's *t*-test.

Correspondence address: J.P.T.M. van Leeuwen, Department of Internal Medicine III, Erasmus University Medical School, PO Box 1738, 3000 DR Rotterdam, The Netherlands

Table I

Effect of chelating extracellular calcium with EGTA and blocking calcium release from intracellular stores by TMB-8 on homologous up-regulation of VDR. Data expressed as fmol [^3H]1,25(OH) $_2\text{D}_3$ bound/mg protein.

[1,25(OH) $_2\text{D}_3$]	Control	1.5 mM EGTA	2 mM EGTA	TMB-8
0	25.4 \pm 1.3	23.5 \pm 0.2	45.0 \pm 1.6*	19.7 \pm 0.8**
10 $^{-10}$ M	40.6 \pm 1.8*	40.4 \pm 1.6	45.8 \pm 0.9 ∇	29.7 \pm 1.0 ∇
10 $^{-8}$ M	106.8 \pm 3.4*	105.9 \pm 4.3	82.0 \pm 2.1 \blacksquare	64.3 \pm 1.1 \blacksquare

* $P < 0.001$, ** $P, 0.05$ with respect to basal VDR content (25.4 \pm 1.3). $\nabla P < 0.05$, $\nabla \nabla P < 0.002$ and $\blacksquare P < 0.001$ calculated as the significance of interaction between 1,25(OH) $_2\text{D}_3$ and EGTA

3. RESULTS AND DISCUSSION

To our knowledge this is the first report showing a direct functional involvement of calcium as intracellular signal in a cellular response to 1,25(OH) $_2\text{D}_3$.

Basal 1,25(OH) $_2\text{D}_3$ binding was significantly reduced by the calcium channel blockers nitrendipine (5×10^{-5} M, $P < 0.05$) and verapamil (10^{-4} M, $P < 0.001$) [12]. With 1.5 mM EGTA no effect was observed whereas 2 mM EGTA induced an increase in VDR level (Table I). Recently it was reported that expression of murine epidermal differentiation markers is tightly regulated by restricted extracellular calcium concentrations [15]. In our study basal medium calcium concentration was 1.42 mM and in the presence of 1.5 and 2 mM EGTA 0.28 and 0.05 mM, respectively. These concentrations are in the same range as those reported to be important for the expression of epidermal differentiation markers [15]. Therefore it is likely that basal VDR level is also tightly regulated by extracellular calcium concentrations. Moreover, preliminary data obtained in our laboratory show an increase of VDR mRNA by 2 mM but not by 1.5 mM EGTA (manuscript in preparation).

As depicted in Figs. 1 and 2, VDR up-regulation by 10^{-10} M 1,25(OH) $_2\text{D}_3$ is significantly reduced by 10^{-5} and 5×10^{-5} M nitrendipine (18 and 48% inhibition)

and by 10^{-5} and 10^{-4} M verapamil (24 and 50% inhibition). In contrast, 10^{-8} M 1,25(OH) $_2\text{D}_3$ -induced VDR up-regulation is not affected by 10^{-5} M nitrendipine and 10^{-5} M verapamil and only 30% and 40% reduction was observed with 5×10^{-5} M nitrendipine and 10^{-4} M verapamil, respectively (Figs. 1 and 2). Together with the results obtained with EGTA (Table I), these data indicate that calcium is functionally involved in homologous up-regulation of VDR in UMR 106 cells. Recently we demonstrated that calcium is also involved in heterologous up-regulation of VDR [12].

A role for intracellular calcium stores in the action of 1,25(OH) $_2\text{D}_3$ is indicated by the data obtained with 0.1 mM TMB-8 (Table I). VDR up-regulation by 10^{-10} and 10^{-8} M 1,25(OH) $_2\text{D}_3$ was inhibited by 34 and 45%, respectively. This is in agreement with the reports showing effects of 1,25(OH) $_2\text{D}_3$ on the phosphoinositide metabolism [10,11].

Both verapamil and nitrendipine inhibited 10^{-10} M 1,25(OH) $_2\text{D}_3$ -induced VDR up-regulation at a lower concentration and with a greater magnitude than the up-regulation by 10^{-8} M 1,25(OH) $_2\text{D}_3$ (Figs. 1 and 2). It is unlikely that this is due to the fact that the increase of the $[\text{Ca}^{2+}]_i$ by 10^{-8} M is more pronounced than the increase by 10^{-10} M 1,25(OH) $_2\text{D}_3$. As Lieberherr [8] reported that a maximum increase of $[\text{Ca}^{2+}]_i$ was




Fig. 1. Effect of nitrendipine on homologous up-regulation of VDR. 5×10^{-5} M nitrendipine significantly ($P < 0.05$) reduced basal 1,25(OH) $_2\text{D}_3$ binding. ** $P < 0.025$; * $P < 0.001$ calculated as the significance of interaction between 1,25(OH) $_2\text{D}_3$ and nitrendipine.




Fig. 2. Effect of verapamil on homologous up-regulation of VDR. 10^{-4} M verapamil significantly ($P < 0.001$) reduced basal 1,25(OH) $_2\text{D}_3$ binding. ** $P < 0.01$, * $P < 0.0025$ calculated as the significance of interaction between 1,25(OH) $_2\text{D}_3$ and verapamil.

observed at 10^{-10} M whereas 10^{-8} M, $1,25(\text{OH})_2\text{D}_3$ has only a minor effect on the $[\text{Ca}^{2+}]_i$. Moreover, Caffrey and Farach-Carson [9] reported agonistic (0.05–5 nM) and antagonistic (> 10 nM) effects on calcium currents. Based on the current data it is tempting to suggest that $1,25(\text{OH})_2\text{D}_3$ causes VDR up-regulation both dependent on and independent of an increase of the $[\text{Ca}^{2+}]_i$, and that the calcium dependency is more prominent at low than at high $1,25(\text{OH})_2\text{D}_3$ concentrations.

Although Bloor et al. [16] have reported detectable levels of $1,25(\text{OH})_2\text{D}_3$ in intestinal nuclei 5 min after administration it is, in view of the fast effect of $1,25(\text{OH})_2\text{D}_3$ on the $[\text{Ca}^{2+}]_i$, unlikely that the effect of $1,25(\text{OH})_2\text{D}_3$ on the $[\text{Ca}^{2+}]_i$ is exerted via the genome. Whether $1,25(\text{OH})_2\text{D}_3$ exerts its effects on $[\text{Ca}^{2+}]_i$ via a cell surface receptor is not yet clear, although recent data point to a cell surface receptor for $1,25(\text{OH})_2\text{D}_3$ [11] and a steroid receptor has been identified on the surface of *Xenopus* oocytes [17].

Taken together, the current study demonstrates (1) a role for the extracellular calcium concentration in the regulation of the VDR level, and (2) a functional messenger role for intracellular calcium in the action of $1,25(\text{OH})_2\text{D}_3$ in osteoblast-like cells and thereby provides new insights in the mechanism of action of this steroid hormone.

Acknowledgement: We thank Dr. J. Lindemans of The Sophia Hospital in Rotterdam for the measurement of the medium calcium concentration.

REFERENCES

- [1] Pan, L.C. and Price, P.A. (1987) *J. Biol. Chem.* 262, 4670–4675.
- [2] Pols, H.A.P., Birkenhäger, J.C., Schilte, J.P. and Visser, T.J. (1988) *Biochim. Biophys. Acta* 970, 122–129.
- [3] Costa, E.M., Hirst, M.A. and Feldman, D. (1985) *Endocrinology* 117, 2203–2210.
- [4] Bikle, D.D., Zolock, D.T., Morrissey, R.L. and Herman, R.H. (1978) *J. Biol. Chem.* 253, 484–488.
- [5] De Boland, A.R. and Boland, R.L. (1987) *Endocrinology* 120, 1858–1864.
- [6] Mezzetti, G., Monti, M.G., Pernecco Casolo, L., Piccinini, G. and Moruzzi, M.S. (1988) *Endocrinology* 122, 389–394.
- [7] Edelman, A., Thil, C.L., Garabedian, M., Anagnostopoulos, T. and Balsan, S. (1983) *Biochim. Biophys. Acta* 732, 300–303.
- [8] Lieberherr, M. (1987) *J. Biol. Chem.* 262, 13168–13173.
- [9] Caffrey, J.M. and Farach-Carson, M.C. (1989) *J. Biol. Chem.* 264, 20265–20274.
- [10] Tang, W., Ziboh, V.A., Isseroff, R.R. and Martinez, D. (1987) *J. Cell. Physiol.* 132, 131–136.
- [11] Lieberherr, M., Grosse, B., Duchambon, P. and Drüeke, T. (1989) *J. Biol. Chem.* 264, 20403–20406.
- [12] Van Leeuwen, J.P.T.M., Birkenhäger, J.C., Schilte, J.P., Buurman, C.J. and Pols, H.A.P. (1990) *Cell Calcium* 11, 281–289.
- [13] Bradford, M.M. (1976) *Anal. Biochem.* 72, 248–254.
- [14] Wallenstein, S., Zucker, C.L. and Fless, J.L. (1980) *Circ. Res.* 47, 1–9.
- [15] Yuspa, S.H., Kilkenny, A.E., Steinert, P.M. and Roop, D.R. (1989) *J. Cell Biol.* 109, 1207–1217.
- [16] Bloor, J.H., Dasmahapatra, A., Weiser, M.M. and Klohs, W.D. (1982) *Biochem. J.* 208, 567–575.
- [17] Sadler, S.E. and Maller, J.L. (1982) *J. Biol. Chem.* 257, 355–361.