

Translation enhancing properties of the 5'-leader of potato virus X genomic RNA

D.A. Zelenina^b, O.I. Kulaeva^a, E.V. Smirnyagina^a, A.G. Solovyev^a, N.A. Miroshnichenko^b,
O.N. Fedorkin^a, N.P. Rodionova^a, S.Yu. Morozov^a and J.G. Atabekov^a

^aDepartment of Virology, Moscow State University, Moscow 119899, USSR and ^bInstitute of Microbiology, Academy of Sciences of the USSR, Moscow 117811, USSR

Received 20 November 1991

The double-stranded DNA copy corresponding to the 5'-nontranslated $\alpha\beta$ -leader of potato virus X (PVX) genomic RNA (positions -3 to -85 according to AUG initiator) was chemically synthesized and fused to the transcription plasmids containing three different reporter genes: neomycinphosphotransferase type II (NPT II) gene, *Bacillus thuringiensis* coleopteran-specific toxic protein gene and β -glucuronidase (GUS) gene. Expression of the reporter genes in vitro and in plant protoplasts (in the case of GUS gene) reveals that the $\alpha\beta$ -leader of PVX RNA acts as a translation enhancer despite the presence of the upstream vector-derived sequence and irrespective of the length of the spacer sequence preceding the reporter genes.

Recombinant DNA; Translational enhancement; Plant virus

1. INTRODUCTION

The 5'-untranslated leaders of various mRNAs have been shown to enhance translation of foreign eukaryotic and prokaryotic genes in vivo and in vitro [1–5]. The 5'-leader of potato virus X (PVX) RNA (83 nt apart from cap structure) consists of two sequences referred to as α -sequence (41 nt 3' to cap) and β -sequence (42 nt upstream from the first AUG) [6]. We have tested $\alpha\beta$ -sequence for its translation-enhancing ability using the gene construct in which neomycin phosphotransferase I gene (NPT I) with deleted 5'-terminal region (including the AUG initiator) was under the control of the initiation codon of the PVX 5'-proximal 165K protein gene in T7-transcription plasmid. The $\alpha\beta$ -leader which contained no vector-derived sequences has been shown to enhance the translation of NPT I gene in different cell-free protein-synthesizing systems by 6–40-fold [6]. In comparative experiments [6] it has been found that the PVX $\alpha\beta$ -leader was almost equally effective as tobacco mosaic virus RNA leader, which was known to be the most efficient translational enhancer [1,7].

In the most versatile transcription vectors containing T7 or SP6 promoters for in vitro usage, or cauliflower mosaic virus (CaMV) 35S promoter for in vivo expression, the polylinker region is positioned downstream from the transcription initiation site. Routine genetic

engineering procedure for fusing of any regulatory element (like the $\alpha\beta$ -leader) and foreign gene usually results in insertion of the nucleotide spacers between the engineered elements. Therefore, the $\alpha\beta$ -carrying chimeric mRNAs constructed by routine procedure will typically contain the vector-derived 5'-nucleotides and/or spacer sequences. To determine the influence of the vector-derived 5'-terminal sequences and the spacer sequences located between the $\alpha\beta$ -leader and the AUG initiation codon, we have fused the $\alpha\beta$ -leader (namely, the sequence positioned -3 to -85 according to AUG-initiator of the 165K protein gene) to the T7-transcription plasmids containing different reporter genes.

Our results show that the $\alpha\beta$ -sequence significantly stimulates translation of chimeric mRNAs despite the presence of the vector-derived sequences upstream from $\alpha\beta$ -leader and the spacer sequences preceding the reporter genes.

2. MATERIALS AND METHODS

2.1. Construction of plasmids for the expression of the reporter genes

The following expression vectors were used: pTZ18R (Pharmacia-LKB) containing T7 promoter for in vitro transcription and pRT101 and pRT103 plasmids which are plant transient-expression vectors containing CaMV 35S promoter and multiple cloning site [8].

The plasmid pTZ-NEO was assembled by insertion of a *Bgl*II-*Bam*HI DNA fragment containing NPT II gene from pNEO (Pharmacia-LKB) into the *Bam*HI site of pTZ18R and selected further by sequence analysis of the recombinant plasmids isolated from the transformed *E. coli* strain XL-1B (Stratagene) cells.

The plasmid pTZ-TOX was assembled by insertion of a *Bam*HI-*Hind*III PCR-generated DNA fragment containing *Bacillus thuringiensis* coleopteran-specific toxic protein gene (Dobrzanskaya et al., unpublished data) into the *Bam*HI-*Hind*III-cut pTZ18R.

Abbreviations: NPT, neomycin phosphotransferase; PVX, potato virus X; GUS, β -glucuronidase.

Correspondence address: D.A. Zelenina, Institute of Microbiology, Academy of Sciences of the USSR, Moscow 117811, USSR.

28°C) in enzyme solution (2% Cellulysin (Calbiochem); 0.1% Macerase (Calbiochem); 10% mannitol, pH 5.5). Protoplasts were isolated by filtration through Miracloth, pelleted by centrifugation at $100 \times g$ for 65 s and then washed by two sedimentation cycles ($100 \times g$ for 65 s). The protoplasts were resuspended in 10% D-mannitol, pH 5.5.

A total of 0.4×10^6 protoplasts was suspended in approximately 5×10^5 ml of 10% mannitol and incubated at 0°C for 30 min. The protoplasts were mixed with pRTGUS or pRTGUS $\alpha\beta$ DNA (20–40 μ g/ml) and exposed to electric pulse of 1 ms at 750 V/cm. The electroporated protoplasts were incubated for 30 min at 0°C. After incubation protoplasts were suspended in 10% mannitol at room temperature, pelleted by centrifugation at $100 \times g$ for 70 s and resuspended in Aoki-Takebe media. After incubation for 20 h at 22°C the protoplasts were collected by centrifugation.

2.4. Determination of β -glucuronidase activity

Fluorometric assays of GUS activity were performed on protoplasts and in vitro-synthesized enzyme. Protoplasts harvested by centrifugation were used in GUS extraction buffer (50 mM sodium phosphate, pH 7.0, 10 mM β -mercaptoethanol, 10 mM Na₂EDTA, 0.1% sodium lauryl sarcosine, 0.1% Triton X-100). The samples were taken from the clarified extracts, mixed with 4-methyl umbelliferyl β -D-glucuronide (MUG) (final concentration 0.5 mM) and incubated at 37°C for 60 min. GUS activity was tested by using 100 μ l-reactions containing 1:10 dilutions of protoplast lysate. Activity assays for in vitro-synthesized GUS were conducted by using 1:100 dilutions of the cell-free system. Reaction samples after incubation with MUG were diluted with 0.4 M Na₂CO₃ up to 10 ml. Fluorescence (excitation 365 nm, emission 455 nm) of each sample was determined in a Hitachi Fluorescence Spectrophotometer 850.

3. RESULTS AND DISCUSSION

Uncapped synthetic RNAs derived from pTZ-NEO $\alpha\beta$, pTZ-NEO, pTZ-TOX $\alpha\beta$, pTZ-TOX, pTZ-




Fig. 3. Relative translation levels of the GUS-gene containing transcripts in rabbit reticulocyte lysate. The transcripts were synthesized using linearized plasmids pTX-GUS $\alpha\beta$ (alpha/beta), pTZ-G12 (control 1), pGEM-GUS (control 2) and pA103-GUS (A103) containing 53 nt leader sequence adjacent to the initiation codon of PVX coat protein gene (D. Zelenina et al., unpublished).

GUS $\alpha\beta$, pTZ-G12 and pGEM-GUS were translated in rabbit reticulocyte lysate. Polypeptide products were fractionated by polyacrylamide gel electrophoresis [9,10]. The results show that the translation of different reporter genes was considerably enhanced in a cell-free system in the presence of the 5'- $\alpha\beta$ -sequence (Fig. 2A, B and C).

To quantitate these data more accurately samples were withdrawn from translation probes of the transcripts from pTZ-GUS $\alpha\beta$, pTZ-G12 and pGEM-GUS and analyzed for GUS activity. Non-capped transcripts containing $\alpha\beta$ -sequence were significantly




Fig. 2. Cell-free translation of chimeric transcripts in rabbit reticulocyte lysate (autoradiograms of gradient 8–20% gels [12]): (A) Lane 1 = control NPT II transcript, 20 μ g/ml; lane 2 = $\alpha\beta$ -containing transcript, 20 μ g/ml; lane 3 = endogenous translation. (B) Lane 1 = control RNA transcript of *B. thuringiensis* TOX gene (20 and 40 μ g/ml, respectively); lanes 2 and 3 = $\alpha\beta$ -containing RNA transcript of *B. thuringiensis* TOX gene (20 and 40 μ g/ml, respectively). (C) Lanes 1 and 2 = control RNA transcript of GUS gene (46 nt leader) (20 and 40 μ g/ml, respectively); lanes 3 and 4 = $\alpha\beta$ -containing RNA transcript of GUS gene (20 and 40 μ g/ml, respectively); lanes 5 and 6 = control RNA transcript of GUS gene (12 nt leader) (20 and 40 μ g/ml, respectively).

more active than those containing the artificial non-physiological 5'-leaders (Fig. 3).

The difference in GUS activity between the transcripts from pTZ-GUS $\alpha\beta$ and control pTZ-G12 varied significantly in different rabbit reticulocyte lysate preparations. Differences between these two transcripts in lysates from different sources varied from 3-fold induction to 14-fold induction (data not shown).

Translation of the transcripts with the artificial 5'-leaders (from pTZ-G12 and pGEM-GUS) was highly dependent on the presence of a cap structure (m⁷GpppG) at the 5'-terminus. The effect of the $\alpha\beta$ -leader on the in vitro expression of GUS gene was quantitatively similar to the effect of mRNA capping (data not shown). It is worth mentioning that the capping of the $\alpha\beta$ -containing PVX-specific mRNAs resulted in further several-fold increase of in vitro translation [9].

To further investigate the translational enhancement activity of the $\alpha\beta$ -sequence, we have used protoplast transient-expression assay. The activity of GUS was detected in protoplasts electroporated with pRTGUS or pRTGUS $\alpha\beta$, but not with pRT103. Electroporation with pRTGUS $\alpha\beta$ resulted in approximately 3-fold higher GUS activity than that with pRTGUS (data not shown). This observation suggests that the $\alpha\beta$ -sequence substantially stimulates translation of naturally capped transcripts produced in plant cell nuclei.

The initiation codons in two control $\alpha\beta$ -lacking constructs (pTZ-G12 and pRTGUS) containing the artificial leaders (Fig. 1) were situated within the local nucleotide context ACCAUGG. This context was shown to be several-fold more effective than the environments in which pyrimidines replace the purines in position -3 [11]. It should be noted that the start codons of $\alpha\beta$ -

containing transcripts from pTZ-GUS $\alpha\beta$ and pRTGUS $\alpha\beta$ are preceded by pyrimidine in position 3 (Fig. 1), as in the case of natural PVX genomic leader [6]. Thus, the observed effect of enhancement by $\alpha\beta$ -sequence seems to be more pronounced taking into account nonfavorable AUG of $\alpha\beta$ -carrying messengers.

Our data demonstrate that a synthetic derivative of the 5'-untranslated sequence of PVX RNA confers translational enhancement on different reporter genes despite the fact that the length and the primary structure of these genes and nonviral spacers upstream from their initiation codons were significantly varied.

Acknowledgements: We thank Dr. R. Töpfer for providing us with the pRT 101 to 105 plasmids.

REFERENCES

- [1] Gallie, D.R., Sleat, D.E., Watts, J.W., Turner, P.C. and Wilson, T.M.A. (1987) *Nucleic Acids Res.* 15, 8693-8711.
- [2] Jobling, S.A. and Gehrke, L. (1987) *Nature* 325, 622-625.
- [3] Dolph, P.D., Racaniello, V., Villarmin, A., Palladino, F., Schneider, R. (1988) *J. Virol.* 62, 2059-2066.
- [4] Ruiz-Linares, A., Bouloy, M., Girard, M. and Cahour, A. (1989) *Nucleic Acids Res.* 17, 2463-2476.
- [5] Carrington, J.C. and Freed, D.D. (1990) *J. Virol.* 64, 1590-1597.
- [6] Smirnyagina, E.V., Morozov, S.Yu., Rodionova, N.P., Miroshnichenko, N.A., Solovyev, A.G., Fedorkin, O.N. and Atabekov, J.G. (1991) *Biochimie* 73, 998-1011.
- [7] Jobling, S.A., Cuthbert, C.M., Rogers, S.G., Fraley, R.T. and Gehrke, L. (1988) *Nucleic Acids Res.* 16, 4483-4498.
- [8] Töpfer, R., Matzeit, V., Gronenborn, B., Schell, J. and Steinbiss, H.-H. (1987) *Nucleic Acids Res.* 15, 5890.
- [9] Morozov, S.Y., Miroshnichenko, N.A., Zelenina, D.A., Fedorkin, O.N., Solovyev, A.G., Lukasheva, L.I. and Atabekov, J.G. (1990) *Biochimie* 72, 677-684.
- [10] Laemmli, U.K. (1970) *Nature* 227, 680-685.
- [11] Kozak, M. (1989) *Mol. Cell. Biol.* 9, 5073-5080.