

Reverse transcriptase-polymerase chain reaction detects induction of cardiac-like α myosin heavy chain mRNA in low frequency stimulated rabbit fast-twitch muscle

Heidemarie Peuker, Dirk Pette*

Fakultät für Biologie, Universität Konstanz, Postfach 5560-M641, D-78434 Konstanz, Germany

Received 8 May 1995; revised version received 15 May 1995

Abstract Using reverse transcriptase-polymerase chain reaction we quantified in rabbit skeletal muscles expression levels of the highly homologous cardiac α and β myosin heavy chain (α MHC, β MHC) mRNA isoforms. Masseter muscle displayed highest levels of a cardiac-like α MHC mRNA. This isoform was present at 20-fold lower amounts in slow soleus and at 200-fold lower levels in several fast-twitch muscles. Low-frequency stimulation periods exceeding 20 days drastically induced the α MHC mRNA in fast tibialis anterior. The α MHC mRNA was 140-fold elevated after 60 days when β MHC mRNA had increased 50-fold. Our results demonstrate the wide distribution of a cardiac-like α MHC mRNA in skeletal muscle and its marked induction during fast-to-slow transition as induced by low-frequency stimulation.

Key words: Cardiac-like α myosin heavy chain; Chronic low-frequency stimulation; Rabbit skeletal muscle; Reverse transcriptase-polymerase chain reaction

1. Introduction

Rabbit skeletal muscles are composed of three fast fiber types (IIB, IID, IIA) and the slow type I fibers, characterized by the fast-type myosin heavy chain (MHC) isoforms MHCIb, MHCIId, MHCIIa and the slow MHCI [1,2]. The latter isoform is thought to be identical with the cardiac β MHC [3], the major isoform in rabbit cardiac ventricle [4]. Contrary to the distribution of β MHC in cardiac ventricle and slow skeletal muscles, the expression of the cardiac α MHC isoform in non-cardiac muscles seems to be limited to special muscles and fibers. To date, the presence of cardiac-like α MHCs has been demonstrated in masticatory and extraocular muscles [5–8], in the diaphragm [9] and in intrafusal fibers [10–12]. A previous immunocytochemical study provided evidence that an α MHC-like isoform is expressed in fast-twitch rabbit muscle under the influence of chronic low frequency stimulation (CLFS) [9], transforming a fast-twitch into a slow-twitch muscle with sequential transitions in MHC isoform expression in the order of MHCIb \rightarrow MHCIId \rightarrow MHCIIa \rightarrow MHCI [13]. The present study addresses the question as to the assignment of α MHC expression in this sequence. For this purpose, we applied the

reverse transcriptase-polymerase chain reaction (RT-PCR) for the detection of a sequence specific to α MHC mRNA. Amplifying this sequence, as well as a sequence specific to the β MHC isoform, we studied the distribution of the two MHC isoforms in various muscles. The detection of α -skeletal actin mRNA served as a control of the method. We show the presence of an α MHC-like isoform at low concentrations in several skeletal muscles of the rabbit and demonstrate its pronounced induction during CLFS-induced muscle transformation.

2. Materials and methods

2.1. Animals, chronic stimulation, muscles

Adult male White New Zealand rabbits were used. Adductor magnus (ADM), diaphragm (DIA), extensor digitorum longus (EDL), gastrocnemius (GAS), masseter (MAS), psoas (PS), and soleus (SOL) muscles were excised and frozen in liquid nitrogen. Chronic low frequency stimulation (CLFS, 10 Hz, 12 hours daily) of tibialis anterior (TA) muscle was performed as described [14]. Three animals were used for each time point (0d, 6d, 14d, 21d, 30d, 60d).

2.2. Preparation of total RNA and oligonucleotide primers

Total RNA was prepared [15,16] and concentrations spectrophotometrically assessed. To amplify sequences specific to α -skeletal actin mRNA [17] and β MHC mRNA, primers were used [18], yielding PCR products of 367 nt and 173 nt length, respectively. For the detection of α MHC mRNA a primer pair was chosen from the 3'-terminal sequence (GenBank accession no. K01867 [19]) to amplify a 227 nt fragment extending from the translated into the untranslated region (Fig. 1). The primer sequences were (a) antisense: TTGCGGGTTAACAAGAGCGG, and (b) sense: TCAAGGCCTACAAGCGCCAG. For controlling the isoform specificity, an additional antisense primer CTGGGCGGATCAAGGCGTCA was chosen for the β MHC mRNA. This primer was derived from an unpublished sequence generously supplied by Dr. R. Zak, Chicago. It corresponds to an independently published sequence [20]. The 5'-ends of the antisense primers of all three sequences were labeled with digoxigenin to allow chemiluminescent detection of PCR products.

2.3. Reverse transcription-polymerase chain reaction

0.2 μ g of total RNA were reverse-transcribed according to [21] as previously described [17], using Avian Myoblastosis Virus reverse transcriptase and primer p(dT)₁₅ (Boehringer). In order to establish comparable conditions for cDNA synthesis the reaction mixture was: 1 μ l aliquots of the RT assay (corresponding to 10 ng of total RNA) were amplified separately for each of the three sequences [17]. The optimized MgCl₂ concentrations were 2 mM for α MHC and α -skeletal actin, and 3.5 mM for β MHC. For amplification the following conditions were chosen: denaturation at 94°C, annealing at 60°C and synthesis at 72°C. The number of cycles was adjusted to the exponential range of amplification [16]. It was 14 for α -skeletal actin and varied from 16 to 25 for the MHC sequences according to their expression levels. To allow quantitative analyses of gene expression, the amplification reactions were monitored using DNA standards. Therefore, PCR fragments were purified from primers, nucleotides and nonspecific reaction products using the QIAEX DNA gel extraction procedure (QIAGEN) after

*Corresponding author. Fax: (49) (7531) 88 39 40.

Abbreviations: ADM, adductor magnus; CLFS, chronic low frequency stimulation; DIA, diaphragm; EDL, extensor digitorum longus; GAS, gastrocnemius; MAS, masseter; MHC, myosin heavy chain; RT-PCR, reverse transcriptase-polymerase chain reaction; SOL, soleus.

electrophoretic separation of amplification assays on 1.5% agarose gels. The purified DNA was spectrophotometrically quantified, and known amounts of each sequence were amplified in parallel to cDNA aliquots derived from reverse-transcribed RNA. The calculated copy number of standard DNA fragments was 1.1×10^6 for actin and 1.1×10^4 , 1.1×10^5 and 1.1×10^6 for α MHC and β MHC.

2.4. Product analysis and quantitative evaluation

Identity and specificity of the PCR products were verified by restriction fragment analysis after 30 cycles. Digestion products were separated electrophoretically and visualized by ethidium bromide staining. For quantitative analysis, 2 μ l of each PCR assay were combined and separated electrophoretically. Product analysis by digoxigenin-based chemiluminescent detection was performed as described [16]. The photographically documented signal intensities were evaluated densitometrically. Quantitative analyses of gene expression was achieved by comparing the results obtained from RNA preparations with those obtained from the corresponding DNA standards. For the estimation of specific mRNA molecule numbers the efficiency of first-strand cDNA was calculated from data previously obtained by quantitative RT/PCR using cRNA standards [16].

3. Results and discussion

3.1. Amplification of a sequence specific to cardiac α MHC mRNA

The highest degree of sequence homology among the sarcomeric MHC isoforms most likely exists between α MHC and β MHC [22]. To distinguish the transcripts of these two isoforms by RT-PCR, the sequence specificity of the primers and the amplified DNA fragments had to be demonstrated. For the α MHC mRNA we chose a primer pair amplifying a 227 nt sequence from the 3'-terminal part extending from the translated into the untranslated region, characterized as highly variable between different isoforms [23] (Fig. 1). This sequence is identical with the recently cloned α MHC sequence EOM2, identified in extraocular muscles [7]. In agreement with previous findings [5,6], strong signal intensities were observed for this sequence in cardiac atrium and masseter muscle. A weaker signal was also detected in the ventricle (not shown), known to contain a developmentally regulated fraction of α MHC in combination with β MHC [24]. In addition, a strong signal was obtained with RNA preparations from low-frequency stimulated (60d) fast-twitch tibialis anterior muscle. The possibility that this signal resulted from unspecific cross reaction with the β MHC RNA was highly improbable because soleus muscle, rich in β MHC, yielded a strong signal for the β MHC sequence but only a faint signal for the α MHC (Table 1). Control assays in the absence of reverse transcriptase yielded no amplification products.

In order to verify isoform specificity, we compared the selected α MHC sequence with the corresponding part of the β MHC sequence (Fig. 1). The comparison indicated that the antisense primer is specific to the α MHC, whereas the sense primer is identical for both isoforms. To test whether the primer specificity was reliable under the applied conditions of amplification, we selected an additional β MHC-specific antisense primer to amplify a corresponding 210 nt fragment of the β MHC that should include an additional *AluI* restriction site (Fig. 1a). Based on this sequence data, the PCR fragment derived from the β MHC isoform could be distinguished from the corresponding α MHC sequence by four instead of three restriction fragments of predictable length. Fig. 1. shows that this was the case for the β MHC-specific fragments obtained from SOL

and stimulated TA muscles (Fig. 1b, lanes 1, 2 undigested and lanes 3, 4 digested) and also for the α MHC-specific fragments, amplified from MAS and stimulated TA muscles (Fig. 1b, lanes 5, 6 digested and lanes 7, 8 undigested). Moreover, the restriction fragment pattern of the β MHC and α MHC sequences were identical in normal and stimulated muscles. The completeness and purity of the digestion assay demonstrated the isoform specificity of the amplified PCR products. The specificity of the previously characterized 173 nt fragment [18] used for the detection of the β MHC transcripts was re-insured by the same approach. The corresponding α MHC sequence was distinct by a unique *PstI* restriction site, while the 173 nt β MHC fragment was not digested (not shown). The unequivocally established isoform specificity of the amplified α MHC and β MHC sequences was a prerequisite for their separate detection. Nevertheless, it seemed justified to assign this mRNA not as α MHC but as cardiac-like α MHC because its identification was based only on a restricted region of the molecule.

In our view, the RT-PCR technique is superior to other methods such as S1 nuclease mapping or in situ hybridization for the distinction of highly homologous mRNA isoforms. Obviously, the sequence specificity is more reliable with the use of short primers instead of long hybridization probes. Due to the high sequence homology between α MHC and β MHC it should be considered that previous data on the expression of β MHC




Fig. 1. Comparison of isoform specificity of the sequences specific to rabbit α MHC and β MHC mRNAs. (A) presents a sequence comparison of two cDNA clones, pMHC α 81 (bases 1–250) specific to α MHC, and pMHC450-S (bases 109–358) specific to β MHC. The sequences extend from the translated into the variable, untranslated region. 20-mer sense and antisense primers are marked. Arrows indicate *AluI* restriction sites. The *AluI*-restriction fragment analysis of the two amplified sequences, 227 nt specific to α MHC and 210 nt specific to β MHC, is documented by the ethidium bromide-stained gel shown in (B). Obtained fragment lengths were as expected. Lanes 1 and 2, 210 nt product from soleus and 60d stimulated tibialis anterior; lanes 3 and 4, corresponding digests. Lanes 5 and 6, 227 nt product from masseter and 60d stimulated tibialis anterior; lanes 7 and 8, corresponding digests. M, DNA molecular weight marker V (Boehringer Mannheim).




Fig. 2. Signal detection and quantitative evaluation of α MHC and β MHC PCR products detected by chemiluminescence. Signals specific to α -skeletal actin (367 nt), α MHC (227 nt), and β MHC (173 nt) were obtained from RNA preparations of tibialis muscle (TA) stimulated for different time periods at low frequency (10 Hz, 12 h daily). RNA samples from TA muscles were amplified together with DNA standard dilutions (ST) matching the corresponding signal intensities. For actin, number of cycles (14), as well as standard amounts (1.1×10^6 molecules) were the same for all measurements. Indications in the bottom lines refer to α MHC and β MHC only and give number of cycles (c.) and specify different standards: S α 1, S β 1, 1.1×10^4 molecules; S α 2, S β 2, 1.1×10^5 molecules; S α 3, S β 3, 1.1×10^6 molecules. Signal intensities were evaluated by densitometry, and template numbers were calculated according to the intensities of the corresponding standards.

(MHC1) obtained by these methods [20,25–27] may have related to both isoforms.

3.2. Electrostimulation-induced expression of α MHC mRNA detected by quantitative evaluation of RT-PCR

Our method used known amounts of purified PCR fragments as external DNA standards for monitoring the amplifications of different cDNAs by PCR and the subsequent chemiluminescent detection (Fig. 2). This approach permitted the quantitative analysis of gene expression of the different sequences with regard to possible variations in amplification efficiencies and allowed reliable evaluation of signal intensities. Using DNA standards, it was possible to determine initial first-strand cDNA template numbers introduced into PCR. Thus, reverse transcription was not included into the standardization procedure, and slight differences in the efficiency of first-strand cDNA synthesis for the different sequences could not be excluded. However, this may have affected only the comparison of data related to different sequences, but not the results for the same sequence. Based on previously determined molecule numbers of specific MHC mRNA isoforms with cRNA standards [16], the efficiencies of reverse transcription in the present study were determined to be in the range of 25 to 30%. Therefore, the applied method is suitable for determining relative expression levels, but it may also allow estimations of mRNA molecule numbers.

α -Skeletal actin has already been proven to be a suitable internal standard for determining relative mRNA expression levels of different MHC isoforms in muscle fiber fragments [17]. In a similar manner actin served as a control for RT-PCR

performed on total RNA to monitor the validity of RNA preparations and endogenous degradation events. In control TA muscle, α -skeletal actin first-strand cDNA amounted to $3.9 \pm 0.7 \times 10^8$ molecules/ μ g of total RNA. The present study showed only minor variations in its expression level (1.6-fold elevation), most likely due to increases in total RNA [28]. Therefore, actin mRNA content in normal TA muscle was used as reference for normalizing the α MHC and β MHC levels.

Fig. 2 demonstrates signal intensities of PCR products visualized by chemiluminescence. Results from muscles exposed to CLFS for increasing time periods (up to 60d) are shown for the three sequences, i.e. α -skeletal actin (367 nt), α MHC (227 nt) and β MHC (173 nt) in combination with corresponding amounts of standards. The increase in the expression of α MHC and β MHC is evident from the increase in the number of standard molecules (10^4 to 10^6) and the decrease in the number of

Table 1
Initial amounts of first-strand cDNA ($\times 10^6/\mu$ g total RNA, means \pm S.D., $n = 4$) present in PCR assays for α MHC and β MHC isoforms in various rabbit muscles

Muscle	α MHC	β MHC
Adductor magnus	1.8 ± 0.5	2.9 ± 1.1
Diaphragm	40.0 ± 10.4	129 ± 23
Extensor digitorum longus	6.2 ± 1.9	12.3 ± 1.9
Gastrocnemius	9.1 ± 2.7	31.7 ± 2.8
Masseter	265 ± 15	13.4 ± 6.4
Psoas (white portion)	0.4 ± 0.1	0.5 ± 0.2
Psoas (red portion)	1.1 ± 0.3	2.7 ± 1.1
Soleus	19.5 ± 3.9	279 ± 34
Tibialis anterior	1.4 ± 0.6	7.1 ± 2.2




Fig. 3. Time course of low frequency stimulation-induced increases in cardiac-like α MHC (filled triangle) and β MHC (filled circles) mRNA isoforms of rabbit tibialis anterior muscle. Data (means \pm S.D.) were calculated as described in legend to Fig. 2. The sum of the two isoforms is represented by the dotted line and open diamonds. Three animals were investigated for each time point. Four independent analyses were performed on each animal.

cycles (25 to 18) necessary to obtain similar signal intensities. Conversely, for actin the amount of standard and the number of cycles were the same up to 60d.

The values resulting from densitometry (Fig. 3) demonstrated pronounced increases in α MHC and β MHC mRNAs during prolonged CLFS. Up to 14d, both isoforms increased only slightly (α MHC 3.5-fold, β MHC 4.2-fold). At 21d, the level of the α MHC mRNA was 50-fold elevated, whereas that of the β MHC mRNA was only 6.5-fold augmented. At this time, the rise in α MHC seemed to precede that of β MHC mRNA. Thereafter, the increase in β MHC mRNA was steeper than that of the α MHC mRNA, reaching at 30d a level 30-fold elevated over control. This steep increase was in agreement with previous results obtained by S1 nuclease mapping [25]. During the same time period the level of α MHC mRNA was 53-fold augmented over control. Longer stimulation periods up to 60d led to further increases in α MHC mRNA at a similar rate, whereas the increase in β MHC mRNA became smaller. At 60d, the total increases in α MHC and β MHC mRNAs amounted to 140-fold and 50-fold, respectively. The attained levels (α MHC = $2.0 \pm 0.2 \times 10^8$, β MHC = $2.3 \pm 0.5 \times 10^8$ first-strand cDNA molecules/ μ g total RNA) were in the range of SOL and MAS, respectively (Table 1). The sum of both transcript levels has been included in Fig. 3 (dotted line) to indicate the time course that would have been measured using a probe reacting with both isoforms. Due to the stronger induction of α MHC, the α MHC/ β MHC ratio was shifted to higher values. It was 0.20 in the control muscle and amounted to 0.85 in 60d stimulated muscle. Therefore, the α MHC reached an expression level similar to that of β MHC. Obviously, α MHC mRNA expression in skeletal muscle, especially during fast-to-slow MHC isoform transition, has escaped detection up to now. Our results suggest, that the α MHC represents an important isoform during the last step of low-frequency-induced muscle transformation. Therefore, the α MHC can be assigned as an additional element in the sequential MHCIIb \rightarrow MHCIIId \rightarrow MHCIIa \rightarrow α MHC/ β MHC isoform transition. Whether its high level is transient or persists with ongoing stimulation can not be decided on the

basis of the present results. However, it may be noted that SOL, a typical slow muscle, displayed a significantly lower level of α MHC mRNA (Table 1) than 60d stimulated TA muscle.

3.3. Expression of α MHC mRNA in additional muscles

Recently the expression of α MHC mRNA and protein was shown for masticatory and extraocular muscles [5–8]. Moreover, α MHC is also present in muscle spindles [10–12]. However, this isoform was never shown to represent a sarcomeric compound in skeletal muscles other than masticatory and extraocular muscles. Its low value in normal TA may reflect its expression in intrafusal fibers. This assumption is in agreement with similar low expression levels detected in other typical fast-twitch muscles, such as ADM and PS (Table 1). Previous methods used for the assessment of α MHC mRNA in skeletal muscles may not have been sensitive enough to detect these very low expression levels [6]. EDL and GAS muscles display slightly higher levels of α MHC mRNA than TA muscle (Table 1). The higher content of β MHC mRNA in these two muscles possibly indicates that α MHC is associated with β MHC in extrafusal fibers. This assumption is corroborated by the twofold higher levels of α MHC mRNA in DIA than in SOL muscle (Table 1), and matched previous immunocytochemical findings on the presence of α MHC-positive fibers in adult rabbit DIA and low frequency stimulated muscle [9]. It will be interesting to investigate single fibers of transforming muscles in order to examine in more detail the sequence of α MHC and β MHC mRNA expression and their distribution with regard to pure and hybrid fibers.

In summary, we have identified a mRNA specific to a cardiac-like α MHC isoform present at low concentrations in normal skeletal muscles, as well as its pronounced induction during fast-to-slow transition of low frequency stimulated skeletal muscle. Posttranscriptional modifications of the α MHC gene transcript were recently demonstrated in rat heart to result from alternative splicing and three different polyadenylation sites [29]. As our data can not rule out possible diversities between the cardiac α MHC mRNA and the isoform under study, its identity remains to be established.

Acknowledgements: This study was supported by the Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 156.

References

- [1] Aigner, S., Gohlsch, B., Härmäläinen, N., Staron, R.S., Uber, A., Wehrle, U. and Pette, D. (1993) *Eur. J. Biochem.* 211, 367–372.
- [2] Pette, D. and Staron, R.S. (1993) *NIPS* 8, 153–157.
- [3] Lompré, A.-M., Nadal-Ginard, B. and Mahdavi, V. (1984) *J. Biol. Chem.* 259, 6437–6446.
- [4] Mahdavi, V., Periasamy, M. and Nadal-Ginard, B. (1982) *Nature* 297, 659–665.
- [5] d'Albis, A., Janmot, C., Mira, J.-C. and Couteaux, R. (1991) *BAM* 1, 23–34.
- [6] d'Albis, A., Anger, M. and Lompré, A.M. (1993) *FEBS Lett.* 324, 178–180.
- [7] Rushbrook, J.I., Weiss, C., Ko, K., Feuerman, M.H., Carleton, S., Ing, A. and Jacoby, J. (1994) *J. Muscle Res. Cell Motil.* 15, 505–515.
- [8] Bredman, J.J., Wessels, A., Weijis, W.A., Kofrage, J.A.M., Soffers, C.A.S. and Moorman, A.F.M. (1991) *Histochem. J.* 23, 160–170.
- [9] Härmäläinen, N. and Pette, D. (1994) *J. Muscle Res. Cell Motil.* 15, 189.
- [10] Pedrosa, F., Soukup, T. and Thornell, L.-E. (1990) *Histochemistry* 95, 105–113.

- [11] Pedrosa-Domellöf, F., Eriksson, P.O., Butler-Browne, G.S. and Thornell, L.E. (1992) *Experientia* 48, 491–494.
- [12] Wessels, A., Soffers, C.A.S., Vermeulen, J.L.M., Mijnders, T.A., Bredman, J.J. and Moorman, A.F.M. (1990) in: *Muscle and motility* (Maréchal, G. and Carraro, U., Eds.) pp. 71–77, Intercept, Andover, Hampshire.
- [13] Pette, D. and Vrbová, G. (1992) *Rev. Physiol. Biochem. Pharmacol.* 120, 116–202.
- [14] Schwarz, G., Leisner, E. and Pette, D. (1983) *Pflügers Arch. Eur. J. Physiol.* 398, 130–133.
- [15] Chomczynski, P. and Sacchi, N. (1987) *Anal. Biochem.* 162, 156–159.
- [16] Peuker, H. and Pette, D. (1993) *FEBS Lett.* 318, 253–258.
- [17] Peuker, H. and Pette, D. (1995) *Anal. Biochem.* 224, 443–446.
- [18] Uber, A. and Pette, D. (1993) *FEBS Lett.* 331, 193–197.
- [19] Sinha, A.M., Friedman, D.J., Nigro, J.M., Jakovcic, S., Rabinowitz, M. and Umeda, P.K. (1984) *J. Biol. Chem.* 259, 6674–6680.
- [20] Brownson, C., Little, P., Jarvis, J.C. and Salmons, S. (1992) *Muscle Nerve* 15, 694–700.
- [21] Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) *Molecular Cloning*, Cold Spring Harbor Laboratory.
- [22] McNally, E.M., Kraft, R., Bravo-Zehnder, M., Taylor, D.A. and Leinwand, L.A. (1989) *J. Mol. Biol.* 210, 665–671.
- [23] Saez, L. and Leinwand, L.A. (1986) *Nucleic Acids Res.* 14, 2951–2969.
- [24] Everett, A.W., Chizzonite, R.A., Clark, W.A. and Zak, R. (1983) in: *Perspectives in Cardiovascular Research*, Vol. 8 (Tarazi, R.C. and Dunbar, J.B., Eds.) pp. 83–92, Raven Press, New York.
- [25] Kirschbaum, B.J., Heilig, A., Härtner, K.-T. and Pette, D. (1989) *FEBS Lett.* 243, 123–126.
- [26] Aigner, S. and Pette, D. (1992) in: *Molecular biology of muscle*, Symp. Vol. 46 (El Haj, A.J., Ed.) pp. 311–317, Soc. Exp. Biol., Cambridge.
- [27] Jacobs-El, J., Ashley, W. and Russell, B. (1993) *Am. J. Physiol.* 34, C79–C84.
- [28] Kirschbaum, B.J., Simoneau, J.-A., Bär, A., Barton, P.J.R., Buckingham, M.E. and Pette, D. (1989) *Eur. J. Biochem.* 179, 23–29.
- [29] Sindhwani, R., Ismail-Beigi, F. and Leinwand, L.A. (1994) *J. Biol. Chem.* 269, 3272–3276.