

FEBS 15992

Erratum to: **Complement assembly of two fragments of the streptococcal protein G B1 domain in aqueous solution** (FEBS 15575)

[*FEBS Letters* 366 (1995) 99–103]*

Naohiro Kobayashi^a, Shinya Honda^b, Hirofumi Yoshii^a, Hatsuho Uedaira^b, Eisuke Munekata^a

^a*Institute of Applied Biochemistry, University of Tsukuba, Tsukuba 305, Japan*

^b*National Institute of Bioscience and Human Technology, Tsukuba 305, Japan*

Figures 5 and 6 were incorrectly positioned and should be interchanged without repositioning the figure legends.

*SSDI 0014-5793(95)00503-X

FEBS 15993

Corrigendum to: **Two distinct inwardly rectifying conductances are expressed in long term dibutyl-cyclic-AMP treated rat cultured cortical astrocytes** (FEBS 15660)

[*FEBS Letters* 367 (1995) 319–325]*

Stefano Ferroni^a, Cristina Marchini^a, Peter Schubert^b, Carmela Rapisarda^a

^a*Department of Human and General Physiology, University of Bologna, Via San Donato 19/2, 4017 Bologna, Italy*

^b*Max-Planck Institute for Psychiatry, Department of Neuromorphology, Martinsried, Germany*

2.3. *Electrophysiological recordings*

At line 9, the standard external solution is lacking of the KCl content. The composition of the used solution is (mM): 140 NaCl, 4 KCl, 2 CaCl₂, 2 MgCl₂, 5 TES, 5 EGTA, 5 glucose, buffered with NaOH to pH 7.3.

3. Results

At page 323, lines 6–7 (right column) the used sodiumbutyrate concentration is 250 μ M.

*SSDI 0014-5793(95)00588-9