

# Identification of the catalytic triad in the haloalkane dehalogenase from *Sphingomonas paucimobilis* UT26

Kamila Hynková<sup>a</sup>, Yuji Nagata<sup>b</sup>, Masamichi Takagi<sup>b</sup>, Jiří Damborský<sup>c,\*</sup>

<sup>a</sup>Department of Environmental Chemistry and Ecotoxicology, Masaryk University, Kotlarska 2, 611 37 Brno, Czech Republic

<sup>b</sup>Laboratory of Cellular Genetics, Department of Biotechnology, The University of Tokyo, Bunkyo-ku, Tokyo 113, Japan

<sup>c</sup>Laboratory of Biomolecular Structure and Dynamics, Masaryk University, Kotlarska 2, 611 37 Brno, Czech Republic

Received 14 December 1998

**Abstract** The haloalkane dehalogenase from *Sphingomonas paucimobilis* UT26 (LinB) is the enzyme involved in the  $\gamma$ -hexachlorocyclohexane degradation. This enzyme hydrolyses a broad range of halogenated aliphatic compounds via an alkyl-enzyme intermediate. LinB is believed to belong to the family of  $\alpha/\beta$ -hydrolases which employ a catalytic triad, i.e. nucleophile-histidine-acid, during the catalytic reaction. The position of the catalytic triad within the sequence of LinB was probed by a site-directed mutagenesis. The catalytic triad residues of the haloalkane dehalogenase LinB are proposed to be D108, H272 and E132. The topological location of the catalytic acid (E132) is after the  $\beta$ -strand six which corresponds to the location of catalytic acid in the pancreatic lipase, but not in the haloalkane dehalogenase of *Xanthobacter autotrophicus* GJ10 which contains the catalytic acid after the  $\beta$ -strand seven.

© 1999 Federation of European Biochemical Societies.

**Key words:**  $\alpha/\beta$ -Hydrolase; Active site mutant; Catalytic triad; Deamidation; Haloalkane dehalogenase; *Xanthobacter autotrophicus* GJ10

## 1. Introduction

Haloalkane dehalogenase LinB is the enzyme isolated from a  $\gamma$ -hexachlorocyclohexane degrading bacterial strain *Sphingomonas paucimobilis* UT26 [1]. This enzyme participates in the biochemical pathway enabling the microorganism to grow on  $\gamma$ -hexachlorocyclohexane as the sole carbon and energy source [2,3]. The LinB enzyme catalyses conversion of 1,3,4,6-tetrachloro-1,4-cyclohexadiene to 2,5-dichloro-2,5-cyclohexadiene-1,4-diol via 2,4,5-trichloro-2,5-cyclohexadiene-1-ol. In addition to cyclic dienes, LinB also converts a broad range of halogenated alkanes and alkenes to their corresponding alcohols [1]. The dehalogenation reaction is catalysed without the requirement of oxygen or any other cofactor.

The broadening substrate specificity of LinB could make this protein of interest for bioremediation purposes. For example, neither LinB nor any other dehalogenase known to date can efficiently catalyse the dehalogenation reaction of the important pesticides 1,2-dichloropropane and 2-chloropropane, which are consequently recalcitrant under aerobic conditions. The catalysis of 1,2-dichloropropane and 2-chloropropane by hydrolytic dehalogenases is possible from the point of view of the reaction mechanism as indicated by quantitative structure-biodegradability relationships studies [4,5]. Modification of the geometry and composition of an active

LinB site could possibly lead more efficient binding and catalysis of these chlorinated propanes.

Knowledge of the tertiary structure of a protein is the best starting point for rational re-design by site-directed mutagenesis. The 3D structure of LinB has not been experimentally determined as yet, but it was predicted by the method of homology modelling [6] based on its similarity to haloalkane dehalogenase from the bacterium *Xanthobacter autotrophicus* GJ10 (DhlA). The 3D structure of the haloalkane dehalogenase from *X. autotrophicus* GJ10 is known from X-ray crystallography [7]. Haloalkane dehalogenase LinB is expected to contain a main and cap domain and it presumably belongs to the family of  $\alpha/\beta$ -hydrolases [8]. All known  $\alpha/\beta$ -hydrolases have a highly conserved arrangement of their catalytic residues: nucleophile-histidine-acid.

In order to confirm the  $\alpha/\beta$ -hydrolase fold of LinB dehalogenase and to validate the homology model for further mutagenesis studies we introduced several point mutations into its structure in positions in which catalytic residues were expected. Here we describe the construction of these mutants and their characterisation.

## 2. Materials and methods

### 2.1. Materials

All chemicals were purchased from Sigma-Aldrich (Milwaukee, WI, USA). The enzymes used for DNA manipulations were obtained from Takara Shuzo Co. (Kyoto, Japan). The mutagenesis primers were obtained from Espec-oligo Service Co. (Tsukuba, Japan). The strain used in this study is *Escherichia coli* HB101 [T. Maniatis, E.F. Fritsch, and J. Sambrook (1982) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY].

### 2.2. Site-directed mutagenesis

Mutagenesis of LinB was performed by using the principle of a LA PCR in vitro mutagenesis kit (TaKaRa Shuzo Co., Kyoto, Japan), according to the provided protocol except for using KOD (King of DNA) polymerase (Toyobo Co., Osaka, Japan) whose fidelity is very high. All of the nucleotide sequences of mutants were confirmed by the dideoxy-chain termination method with an automated DNA sequencer (LI-COR model 4000L, Aloka Co., Tokyo, Japan).

The oligo nucleotides used are as follows: D108N (5'-GGC GGA CCC CCA GTT ATG CAC GAC CAG AAC-3'), D108A (5'-GGC GGA CCC CCA GGC ATG CAC GAC CAG AAC-3'), E132Q (5'-GGC GAT CGC TT~G~CAT ATA GGC-3') H272A (5'-CTC CTG GAT GAA A~GC~GGC GCC TGC GAC-3'), E244Q (5'-GGC TCC CGG CTG GGC GTT GAT-3').

### 2.3. Overexpression of LinB mutants in *E. coli*

To overproduce LinB mutants in *E. coli*, plasmids for overexpression were constructed from pAQN, which has the same structure as pAQI [9] except for differences in the aqualysin I-coding region. In these plasmids, *linB* mutants are transcribed by the *tac* promoter (*P<sub>tac</sub>*) under the control of *lacI<sup>q</sup>*.

*E. coli* HB101 containing these plasmids were cultured in 2 litres of

\*Correspondence author. Fax: (420) (5) 4112 9506.

E-mail: jiri@chemi.muni.cz

Table 1  
Michaelis-Menten constants with standard errors of wild-type haloalkane dehalogenase LinB and its single-point mutants

	Wt	D108A	D108N	E132Q	E244Q	H272A
$K_M$ (mM)	$0.14 \pm 0.007$	n.a.	$0.30 \pm 0.017$	n.a.	$0.32 \pm 0.022$	n.a.
$k_{cat}$ ( $s^{-1}$ )	$0.98 \pm 0.028$	0	$0.43 \pm 0.010$	0	$0.35 \pm 0.011$	0
$k_{cat}/K_M$	8.92	0	1.42	0	1.10	0

n.a., not applicable at 100 mM substrate concentration.

Luria broth (LB) without sodium chloride at 30°C. Cells were harvested (10 g of wet weight) after induction with 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG), washed in 50 mM potassium phosphate buffer (pH 7.5), and resuspended in the buffer (50 mM potassium phosphate buffer (pH 7.5) containing 1 mM 2-mercaptoethanol and 10% glycerol).

#### 2.4. Purification of LinB mutants

The cells were disrupted by sonication (Sonifier 250; Branson, Danbury, CT, USA). After centrifugation at  $100\,000 \times g$  for 1 h, the supernatant was used as a crude extract. It was further purified by a DEAE-Toyopearl 650M column (Tosho, Tokyo, Japan), and a Sephacryl S-200 column (Pharmacia, Uppsala, Sweden) with Bio Logic LP system (Bio-Rad). Fractions at which LinB wild-type protein was eluted were collected. A major one protein band was observed on SDS-polyacrylamide gel electrophoresis after two steps of purification.

The amount of protein was determined by the protein assay kit (Bio-Rad), with bovine serum albumin as a standard.

#### 2.5. Circular dichroism spectra

Circular dichroism (CD) spectra were measured with a JASCO J-720 spectropolarimeter at room temperature. The cells used were 1 and 0.1 cm light paths for wavelengths of 250–320 and 190–250 nm, respectively. 0.3–0.4 mg/ml of enzymes in 50 mM potassium phosphate buffer (pH 7.5) were used.

#### 2.6. Michaelis-Menten kinetics

Michaelis-Menten kinetic constants were determined by initial-velocity measurements. Gas chromatography was used for the determination of the substrate and product concentration. A dehalogenation reaction was performed in 25 ml Reacti-Flasks closed by Mininert Valves. 10 ml of glycine buffer (pH 8.6) was mixed with seven different substrate concentrations. The largest concentration of substrate in 10 ml glycine buffer served also as an abiotic control. The reaction mixture was equilibrated for 30 min in a shaking water bath at 37°C prior to the reaction initiation. Enzymatic reaction was initiated by adding 100  $\mu$ l of the enzyme preparation. The reaction progress was monitored by withdrawing 0.5 ml samples at 0, 20 and 30 min using a syringe needle to reduce evaporation of the substrate from the reaction mixture. The reaction mixture samples were mixed with 0.5 ml of methanol to terminate the reaction and directly applied on gas chro-

matography with a flame ionisation detector (Hewlett Packard 6890). The capillary column DB-FFAP 30 m  $\times$  0.25 mm  $\times$  0.25  $\mu$ l (J and W Scientific) was used for separation. Samples were injected by using a split technique (split ratio 50:1). The temperature program was isothermal and was dependent on the character of analysed compounds. Parallel determination of the substrate and product concentration in the same water-methanol sample was performed for construction of a calibration curve.  $K_M$  and  $k_{cat}$  values with their standard deviations were calculated by means of the program Leonora [10] by the method of least squares with relative weighting.

#### 2.7. Activity measurements

Dehalogenation reaction rates were determined for nine different substrates using the method of gas chromatography. A dehalogenation reaction was performed in 25 ml Reacti-Flasks closed by the Mininert Valves. 10 ml glycine buffer and 6.4  $\mu$ l of substrate was equilibrated at 37°C for 30 min in a shaking water bath prior to the reaction initiation. Enzymatic reaction was initiated by adding 100  $\mu$ l enzyme (the concentrations of wild-type enzyme and mutant proteins were the same, 0.33 mg/ml). The progress of the reaction was monitored at 10, 20 and 30 min using the same procedure as described in Section 2.6. The reaction rates were quantified by a slope of the relationship between the product concentration and time. Dehalogenation activity was expressed in percentages, where the rate of reaction of wild-type enzyme is 100% for each substrate.

### 3. Results

#### 3.1. Construction and purification of LinB mutants

The catalytic triad in DhIA consists of amino acid residues Asp-124, His-289 and Asp-260. The corresponding residues in the homology model of LinB are Asp-108, His-272 and Glu-132 [6]. The same residues were proposed to act as the catalytic triad in LinB by Krooshof and coworkers [11]. On the other hand, Fetzner and Lingens [12] suggested that it is the residue E244 which fulfils the role of acid in the catalytic triad of LinB. Five point mutants were prepared to test these proposals and to identify the catalytic triad in LinB dehaloge-




Fig. 1. Stereoview of the homology model of haloalkane dehalogenase LinB. The protein backbone and the side-chains of mutated residues are displayed.




Fig. 2. SDS-polyacrylamide gel electrophoresis of purified LinB wild type and its five mutants. Lanes: 1, molecular mass markers; 2, LinB (wt); 3, D108A; 4, D108N; 5, E244Q; 6, E132Q; 7, H272A; 8, molecular mass markers.

nase: D108N and D108A (putative nucleophile mutants), H272A (putative base mutant), E132Q and E244Q (putative catalytic acid mutant). The position of these residues in the 3D model of LinB is shown in Fig. 1.

The mutants were constructed by the method of site-directed mutagenesis and purified to homogeneity (Fig. 2). The integrity of each mutant structure was tested by circular dichroism spectroscopy. The CD spectra of the wild-type dehalogenase and its two mutants are shown in Fig. 3. Like the wild-type dehalogenase, point mutants had the double ellipticity minimum at 210 and 220 nm typical of  $\alpha$ -helical content. The CD spectra of the wild-type and five mutant proteins are essentially the same indicating that the backbone polypeptide chain of constructed mutants had a very similar conformation. It was concluded that the single point substitutions did not disturb the overall structure of the protein mutants.

### 3.2. Activities of mutant proteins

The purified wild-type and mutant enzymes were examined for their kinetic characteristics. 1-Chlorobutane was selected as a substrate because it easily undertakes the dehalogenation reaction by the wild-type enzyme and is often used as a reference compound for characterisation of the haloalkane dehalogenases [5]. No activity was observed for the mutant enzymes D108A, E132Q and H272A even at very high




Fig. 3. Circular dichroism spectra for wavelengths between 190 and 250 nm of wild-type haloalkane dehalogenase LinB and its five mutants (D108A, D108N, E244Q, H272A, E132Q).




Fig. 4. Plot of the product concentration versus time for dehalogenation of 1-chlorobutane by wild-type haloalkane dehalogenase LinB and its mutants D108N and E244Q. The product of this reaction is butanol. Slope of this linear relationship was used for quantification of the enzyme activities.

substrate concentration (100 mM), while the mutants D108N and E244Q showed significant activity Table 1. The activity of D108N was about 44% and E244Q 36% of the wild-type enzyme as deduced from the  $k_{cat}$  values.

The activity of D108N and E244Q enzymes was determined with the group of nine different substrates to test whether the reduction in activity observed is substrate-dependent. The reaction rates were quantified by a slope of the relationship between the product concentration and time (Fig. 4). The rates obtained with the mutants are expressed as the percentage activity of the wild-type enzyme in Table 2.

The mutant enzyme D108N expressed in an average of 58% of the activity of the wild-type enzyme and the mutant enzyme, while E244Q showed in an average of the 38% activity of the wild-type enzyme. The decrease in activity of the mutants E244Q and D108N was very similar for different substrates, with the standard deviation of 7% for D108N and 5% for E244Q. Activity reduction is independent of the substrates shape, size or number of their substituents.

## 4. Discussion

### 4.1. Identification of the catalytic triad of LinB

The objective of this work was to identify the catalytic residues in the haloalkane dehalogenase LinB and to confirm

Table 2  
Relative activities<sup>a</sup> of LinB mutants with different substrates

Substrate	D108N %	E244Q %
1-Chlorobutane	62	43
1,3-Dibromopropane	64	38
1,3-Dichloropropane	58	44
1-Bromopropane	43	34
1-Bromo-3-chloropropane	63	47
1,2-Dibromopropane	51	32
1-Bromo-2-methylpropane	58	31
1-Bromohexane	60	36
1-Bromocyclohexane	66	39
Average	58	38
S.D.	7	5

<sup>a</sup>Activities express the percentage of the rates obtained with the wild-type enzyme.




Fig. 5. Topological arrangement of the secondary structure elements in the structure of haloalkane dehalogenase. The different location of the catalytic triad residues are given by stars. (Empty stars: catalytic triad of LinB, full stars: catalytic triad of DhlA).

the structural fold of this enzyme. The catalytic triad of LinB was proposed from the model of its structure obtained from homology modelling. This model was based on the X-ray structure of the haloalkane dehalogenase from *X. autotrophicus* GJ10 (DhlA). Two out of three catalytic residues are conserved in both enzymes, Asp-108 in LinB corresponds to DhlA nucleophile Asp-124, while His-272 of LinB corresponds to DhlA histidine His-289 (Fig. 5). The amino acid sequence is strongly conserved in the regions surrounding these residues [11,13]. The position of the catalytic acid of LinB is less obvious, and two possible locations were consid-

ered, E132 and E244. Overall, five different point mutants of LinB have been constructed, purified and tested for activity: D108N, D108A, E132Q, E244Q and H272A.

The mutants D108A, E132Q and H272A were observed to be inactive in the dehalogenation experiments. These three amino acids are essential for the catalytic reaction and presumably form the catalytic triad of LinB. No activity was expected either with the mutant D108N. However, approximately 58% activity of the wild-type enzyme was observed with this mutant. The mutant enzyme D108N was most probably reactivated by deamidation yielding the wild-type enzyme. The mechanism of deamidation was previously described by Priest and coworkers who studied the D124N nucleophile mutant of the haloalkane dehalogenase DhlA [14]. Priest suggested that the deamidation is caused by a direct attack by the activated water molecule on the side-chain amide carbon of replaced asparagine. This highly nucleophilic water molecule is located close to the carboxyl moiety of the nucleophilic catalytic residues and in the wild-type enzyme is involved in the hydrolytic dehalogenation mechanism. Reduced activity indicates that not all enzyme mutants undertake the deamidation process.

The amino acid E244 was not essential for the LinB activity and apparently is not the third catalytic amino acid of the LinB enzyme. The deamidation reaction observed with the D108N mutant is not expected to occur in this mutant since no nucleophilic water molecule occurs in its surroundings. The significant reduction in the dehalogenation activity of the E244Q mutant, however, suggests that this residue still plays some important role in the function of LinB. Reduced activity, which is an average of 38% of the activity of the wild-type enzyme, is about the same for substrates with different chain-length, overall shape and the number of substituents.

#### 4.2. Topological position of the catalytic triad in LinB

The composition of the catalytic residues in DhlA and LinB is similar to that which is found in serine proteases, lipases, and several other  $\alpha/\beta$ -hydrolase fold enzymes [8,15].  $\alpha/\beta$ -Hydrolases generally show very little sequence similarity. However they share the same pattern in the composition of secondary elements and the same arrangement of the catalytic




Fig. 6. Proposed reaction mechanism of haloalkane dehalogenase LinB. A: Nucleophilic attack of A108 on the  $C_{\alpha}$  of the alkyl halide, leading to formation of the covalent alkyl-enzyme intermediate. B: H272-assisted hydrolysis of the intermediate.

residues. The order of the catalytic triad in the primary sequence of these proteins is nucleophile-histidine-acid. Without exception, the nucleophile is located at the end of the fifth  $\beta$ -strand and the histidine residue belongs to a loop that follows  $\beta$ -strand eight of the  $\beta$ -sheet. In all but one  $\alpha/\beta$ -hydrolase, the catalytic acid comes from a reverse turn following strand seven [8,16]. Schrag and coworkers noted that the acidic residue in pancreatic lipase 'migrated' to a bend one strand closer to the nucleophile and is positioned after the  $\beta$ -strand six [17]. The authors speculated that the direction of this migration was from strand seven to strand six rather than the opposite direction. This proposal was based on the fact that the enzymes with the acid after the  $\beta$ -strand six are mammalian compared to the enzymes with the acid after the  $\beta$ -strand six which are bacterial, fungal, plant and fish enzymes. The catalytic acid of the haloalkane dehalogenase from *Sphingomonas paucimobilis* UT26 is located after the  $\beta$ -strand six (Fig. 5). There are at least two other bacterial hydrolases, i.e. haloalkane dehalogenase of *Rhodococcus rhodochrous* NCIMB 13064 [18], and putative haloalkane dehalogenase of *Mycobacterium tuberculosis* H37Rv [13] which has a catalytic acid located after the  $\beta$ -strand six. The fact that bacterial enzymes exist also with the catalytic acid located after strand six creates an uncertainty about the direction of migration.

#### 4.3. Proposal of catalytic mechanism of LinB dehalogenase

The catalytic triad of LinB was identified within this study. It is proposed, that the LinB catalyses hydrolytic dehalogenation via the same reaction mechanism as Dh1A [15,19,20]. The putative reaction mechanism of LinB is schematically shown in Fig. 6. A108 is expected to act as a nucleophile causing release of the halide ion and formation of an intermediate ester. A hydrolytic water molecule, which may be located near the nucleophile, cleaves the covalent ester intermediate. This water molecule is activated by H272 acting as a base catalyst prior to hydrolysis. E132 keeps H272 in proper orientation and stabilises a positive charge that develops on histidine imidazole ring during the reaction. Catalytic residues all together act as an electronic pump by stabilising reaction intermediates. In future research we will focus on identification of the other amino acids which are located inside the enzyme active site and could be important in the catalytic reaction and specificity of the haloalkane dehalogenase LinB.

**Acknowledgements:** This project was financially supported by the Czech Grant Agency (Postdoc 203/97/P149), the Czech Ministry of Education (ME276/1998) and by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan. A part of this work was performed in the facilities of the Biotechnology Research Center, The University of Tokyo. Financial support is gratefully acknowledged.

#### References

- [1] Nagata, Y., Miyauchi, K., Damborsky, J., Manova, K., Ansorgova, A. and Takagi, M. (1997) Appl. Environ. Microbiol. 63, 3707–3710.
- [2] Nagata, Y., Nariya, T., Ohtomo, R., Fukuda, M., Yano, K. and Takagi, M. (1993) J. Bacteriol. 175, 6403–6410.
- [3] Nagasawa, S., Kikuchi, R., Nagata, Y., Takagi, M. and Matsuo, M. (1993) Chemosphere 26, 1719–1728.
- [4] Damborsky, J., Manova, K. and Kutý, M. (1996) in: Biodegradability Prediction (Peijnenburg, W.J.G.M. and Damborsky, J., Eds.), pp. 75–92, Kluwer Academic, Dordrecht.
- [5] Damborsky, J., Berglund, A., Kutý, M., Ansorgova, A., Nagata, Y. and Sjöström, M. (1998) Quant. Struct. Act. Rel. 17, 450–458.
- [6] Damborsky, J. and Koca, J. (1999) submitted for publication.
- [7] Verschuere, K.H.G., Seljee, F., Rozeboom, H.J., Kalk, K.H. and Dijkstra, B.W. (1993) Nature 363, 693–698.
- [8] Ollis, D.L. et al. (1992) Protein Eng. 5, 197–211.
- [9] Terada, I., Kwon, S.-T., Miyata, Y., Matsuzawa, H. and Ohta, T. (1990) J. Biol. Chem. 265, 6576–6581.
- [10] Cornish-Bowden, A. (1995) Analysis of Enzyme Kinetic Data, Oxford University Press, Oxford.
- [11] Krooshof, G.H., Kwant, E.M., Damborsky, J., Koca, J. and Janssen, D.B. (1997) Biochemistry 36, 9571–9580.
- [12] Fetzner, S. and Lingens, F. (1994) Microbiol. Rev. 58, 641–685.
- [13] Damborsky, J. (1998) Pure Appl. Chem. 70, 1375–1383.
- [14] Pries, F., Kingma, J. and Janssen, D.B. (1995) FEBS Lett. 358, 171–174.
- [15] Pries, F., Kingma, J., Pentega, M., VanPouderoyen, G., Jeronimus-Stratingh, C.M., Bruins, A.P. and Janssen, D.B. (1994) Biochemistry 33, 1242–1247.
- [16] Smith, L.C., Faustinella, F. and Chan, L. (1992) Curr. Opin. Struct. Biol. 2, 490–496.
- [17] Schrag, J.D., Winkler, F.K. and Cygler, M. (1992) J. Biol. Chem. 267, 4300–4303.
- [18] Kulakova, A.N., Larkin, M.J. and Kulakov, L.A. (1997) Microbiology 143, 109–115.
- [19] Pries, F., Kingma, J., Krooshof, G.H., Jeronimus-Stratingh, C.M., Bruins, A.P. and Janssen, D.B. (1995) J. Biol. Chem. 270, 10405–10411.
- [20] Kennes, C., Pries, F., Krooshof, G.H., Bokma, E., Kingma, J. and Janssen, D.B. (1995) Eur. J. Biochem. 228, 403–407.