

Biochemical and sensorial evaluation of intact and boned broiler breast meat tenderness during ageing

Pauline R. Kriese^a, Adriana L. Soares^a, Paulo D. Guarnieri^b, Sandra H. Prudencio^a,
Elza I. Ida^a, Massami Shimokomaki^{a,*}

^a Graduate Programme in Food Science, Department of Food Science and Technology, Agricultural Sciences Centre, Londrina State University, P.O. Box 6001, CEP 86051-990 Londrina, PR, Brazil

^b Graduate Program in Food Science, Faculty of Pharmaceutical Sciences, São Paulo University, São Paulo, Brazil

Received 13 November 2006; received in revised form 16 January 2007; accepted 2 March 2007

Abstract

Meat tenderness is the main characteristic demanded by consumers and is affected by *rigor mortis* development and proteolysis activities, both of which occur during carcass refrigeration. In this work, we demonstrate that broiler breast fillet tenderness can be further increased and its extension depends on whether or not meat is excised from the carcass. Post-harvest samples taken from 0 to 72 h after slaughtering and kept refrigerated at 2 ± 2 °C were evaluated for tenderness by myofibrillar fragmentation index determination, shear force analysis and sensorial testing. The 24 h post-harvested intact samples were 30.6% more tender than excised samples and 41.7% more tender than control samples ($p \leq 0.05$). The myofibrillar fragments index was 13.2% higher in intact samples than in deboned fillet ($p \leq 0.05$) and a sensory test showed that the 24 h intact samples were of major acceptability. Our results demonstrated that tenderness was best achieved with intact breast fillet samples stored at 2 ± 2 °C for 24 h.

© 2007 Published by Elsevier Ltd.

Keywords: Myofibrillar fragmentation index; Shear force values; Acceptance test; Calpain system; Meat aging

1. Introduction

Consumer satisfaction with meat tenderness will determine whether the purchase is repeated in the future (Morgan et al., 1991). Therefore, it is critical for food industries and meat processors to consistently produce a tender meat product that meets or exceeds consumer expectations. In order to achieve consistency in tenderness it is necessary to standardize the procedures of *ante-* and *post-mortem* handling of carcasses. Several factors explain this variation in meat tenderness, including the amount of intramuscular fat, water-holding capacity and actomyosin complex (Avery, Sims, Warkup, & Bailey, 1996). Collagen and its crosslinking are involved as the broilers advance in age

(Coro, Youssef, & Shimokomaki, 2002) and other factors are the changes that occur during *post-mortem* storage, such as aging, type of rigor, sarcomere length, skeletal restraint and proteolytic activity (Pearson, 1987). Koohmaraie (1996) reported that prevention of sarcomere length shortening could prevent meat toughening. In poultry, cold shortening is not normally a cause of toughening in normal industrial processing, provided that the breast meat is not prematurely excised or otherwise altered (Papinaho & Fletcher, 1996). Papinaho and Fletcher (1996) described temperature-induced shortening in broiler breast muscle sampled as intact and excised muscle. *Post-mortem* tenderization by proteolysis has been extensively studied and the calpain system has been known to initiate the myofibrillar framework enzymatic digestion (Goll, Boehm, Geesink, & Thompson, 1997). Koohmaraie (1996) pointed out that breakdown of the key proteins that maintain the sarcomere structure causes its weakening. Hence alterations in meat

* Corresponding author. Tel.: +55 43 33714565; fax: +55 43 33714080.
E-mail address: mshimo@uel.br (M. Shimokomaki).

tenderness and the Z-line region degradation are among the main changes occurring during *post-mortem* tenderization (Taylor, Gessink, Thompson, Koohmaraie, & Goll, 1995). Earlier work has shown that early boning of breast fillet can yield a tough meat (Sams & Janky, 1986; Smith, Fletcher, & Papa, 1992) and tenderness can be improved by prolonging the time between slaughter and deboning (Notthcutt, Buhr, Young, Lyon, & Ware, 2001). Sams and co-workers have reported the relationship between proteolysis activity and shear force values during cold storage (McKee, Hirschler, & Sams, 1997; Sams, 1999).

The objective of this work was to investigate the tenderness of broiler breast meat through myofibrillar fragmentation index, shear force measurement and sensory acceptance of intact and excised samples during refrigeration at an industrial processing plant.

2. Materials and methods

2.1. Sample preparation

Sixty six chickens, of Cobb lineage of 42 days, were slaughtered according to the standard industrial plant practice. This consists essentially of a sequence of electrical stunning, bleeding, defeathering, evisceration, carcass water cooling, deboning and refrigeration (Guarnieri et al., 2004; Northcutt, 2001).

2.2. Refrigeration techniques

Meat samples were divided into two experimental groups as follows: intact breast ($n = 6$) and excised breast ($n = 6$) meat (*Pectoralis major m.*) were stored in plastic boxes at 2 ± 2 °C for 0, 8, 12, 24, 48 and 72 h *post-mortem*. *Post-mortem* samples ($n = 6$), taken immediately after slaughter (0 h), were treated as controls, irrespective of being deboned or not. For temperature decline rate, measurement samples were taken at 0, 1, 2, 3, 4, 5, 6, 7, 8 and 24 h of storage time. Samples were taken for myofibrillar fragmentation index (MFI) determination and shear force measurement at 0, 8, 12, 24, 48 and 72 h *post-mortem*.

2.3. Myofibrillar fragmentation index (MFI)

MFI was determined as an indirect measurement of calpain activity according to Culler, Parrish, Smith, and Cross (1978). Four grams muscle samples, free of external fat and visible connective tissue, were homogenized for 30 s in 20 ml of MFI buffer (100 mM KCl, 20 mM potassium phosphate, 1 mM EDTA, 1 mM MgCl₂ and 1 mM NaN₃ at pH 7.0). The homogenate was centrifuged at 10×10^3 rpm for 15 min at 2 °C, the supernatant discarded and pellet resuspended in 20 ml of the MFI buffer and centrifuged at 10×10^3 rpm for 15 min at 2 °C. The supernatant was discarded and the pellet suspended in 10 ml of the same MFI buffer. The myofibril suspension was poured through a strainer to remove connective tissue, and then assayed for

protein concentration using the Biuret method. Aliquots of the suspensions were diluted in the MFI buffer to a final protein concentration of 0.5 mg/ml. The diluted protein suspension was poured into a cuvette and the absorbance at 540 nm was immediately measured with a spectrophotometer. The MFI was expressed as $A_{540\text{nm}} \times 200$.

2.4. Shear force measurement

Samples were packed in plastic bags and submitted for cooking in a water bath until the internal temperature reached the value of 75 °C. After refrigeration at 2 ± 2 °C for 12 h, samples were cut into 1 cm³, and analysed on a texturometer TATX-2i. The results were expressed in Newtons.

2.5. Sensory analysis

Six intact and six excised breast samples, taken at 0, 24 and 72 h *post-mortem* refrigerated at 2 ± 2 °C, were submitted for sensory analysis. Broiler breast fillet was vacuum-packed and submitted to the cooking process in an oven at 80 °C until the internal temperature reached the value of 75 °C. After cooling to 28 °C, samples were randomly presented to 28 subjects of a sensorial group in order to evaluate their acceptance. A structure hedonic scale of 9 points (1 = dislike very much to 9 = like very much) was used (Meilgaard, Civille, & Carr, 1999).

2.6. Statistical analysis

The experiment was entirely randomised and experimental treatments were intact and excised meat samples, evaluated at the refrigeration periods of 0, 8, 12, 24, 48 and 72 h *post-mortem*. Statistical analysis was carried out using the Statistica Program version 5.0 (Oklahoma, 1995). The Student test *t* was also applied to compare both treatments, intact and excised samples, in each period in relation to MFI and meat tenderness. The sensorial experiment was of a randomised blocks design with variance analysis. The Tukey test was used for comparisons of acceptance after experimental treatments.

3. Results and discussion

3.1. Myofibrillar fragmentation index

Fig. 1 shows the MFI of intact and excised broiler breast meat. As can be seen, the MFI increased from its initiation up to a 24 h duration of refrigeration. It became virtually constant after 72 h of storage for both samples. These results indicated that the endogenous muscle proteases system reached its maximum activity under conditions of 2 ± 2 °C after approximately 24 h of refrigeration. However, entire breast presented MFIs higher ($p \leq 0.05$) than deboned samples throughout treatment. This difference was 13.2% from 24 h onwards.




Fig. 1. Myofibrillar fragmentation index (MFI) of intact and excised samples of broiler breast stored at $2 \pm 2^\circ\text{C}$ for 0, 8, 12, 24, 48 and 72 h.

3.2. Sample temperature behaviour

Fig. 2 shows the temperature decline rate for both intact and excised samples. As can be seen, the initial experimental time temperature values were 11.3 ± 0.8 for intact and 9.4 ± 0.1 for excised samples and these gradually fell to 0.9 ± 0.1 and 0.8 ± 0.0 , respectively ($p \leq 0.05$), after ca. 5 h of storage, then remaining constant throughout the experiment.

3.3. Breast meat tenderness measurement

Fig. 3 shows the decrease of shear force value for both samples, this being more noticeable in samples kept for 24 h, then becoming virtually constant throughout the experiment. Again, the tenderness was higher in intact than in excised samples ($p \leq 0.05$). A significant negative correlation ($R = -0.93$, $p \leq 0.05$) was observed between MFI and shear force values, suggesting that the higher the proteases activity the more tender became the breast meat.




Fig. 2. Temperature decline rate of intact and excised samples of broiler breast stored at $2 \pm 2^\circ\text{C}$ measured at 0, 1, 2, 3, 4, 5, 6, 7, 8 and 24 h. Note that the temperature values of intact samples are relatively higher than excised samples.




Fig. 3. Shear force values of intact and excised samples of broiler breast stored at $2 \pm 2^\circ\text{C}$ for 0, 8, 12, 24, 48 and 72 h.

One of these enzyme entities, the calpain enzyme system, digested the sarcomere structure, particularly the Z-lines region (Guarnieri et al., 2004; Taylor et al., 1995), improving the meat tenderness. Synergistically, samples attached to bones were protected against muscle cold-shortening, combined with proteolysis, and this may have provided enhanced tenderization. However, as yet, our experiments could not discern potential factors present in bones to possibly explain the biochemical mechanism involved with tenderness in relation to the relative increase of enzymatic proteolytic activities. Fig. 2 offers another possible explanation because of the higher temperature values of the intact samples compared to deboned fillet. Although the difference started at 1.9°C at the beginning of storage time (Fig. 2), these values seem to be critical for the protease activities since the pH values measured (not shown) in both samples were not different ($p \leq 0.05$), causing the meat to become tender in boned samples.

3.4. Sensorial analysis

Table 1 shows the results of the acceptance test. Intact samples refrigerated at 24 and 72 h were those most appreciated by the subjects ($p \leq 0.05$), with the hedonic values of 6.75 and 6.93, respectively. On the other hand, fillets without refrigeration and excised broiler breast meat refrigerated

Table 1

Acceptance of intact and excised samples of broiler breast meat refrigerated at $2 \pm 2^\circ\text{C}$ for 0, 24 and 72 h

Refrigerated samples	Hedonic value ¹
Breast fillet 0 h	4.18 ^b \pm 1.85
Intact breast meat 24 h	6.75 ^a \pm 1.97
Excised breast meat 24 h	5.36 ^b \pm 1.93
Intact breast meat 72 h	6.93 ^a \pm 1.41
Excised breast meat 72 h	5.32 ^b \pm 2.25

Different letters indicate significant differences ($p \leq 0.05$) by Tukey test.
¹ Structured hedonic scale of 9 points, 1 = dislike very much to 9 = like very much.

ated for periods of 24 and 72 h showed lower acceptances ($p \leq 0.05$) with values of 5.36 and 5.32, respectively. These results indicated that subjects preferred samples with higher MFI and lower values of shear.

4. Conclusions

The results of myofibrillar fragmentation index, shear force measurement and sensory analysis demonstrated that storing intact broiler breast fillet, refrigerated at 2 ± 2 °C for a minimum of 24 h, was a feasible procedure for obtaining a very tender meat.

Acknowledgements

PRK is a graduate student of the Food Science Program, Department of Food and Drugs Technology, Londrina State University, ALS is a member of the Post Doctoral CNPq Program, Proc. CNPq 150179/2004-9. MS and EII are CNPq Research Fellows.

References

- Avery, N. C., Sims, T. J., Warkup, C., & Bailey, A. J. (1996). Collagen cross-linking in porcine *M. longissimus lumborum*: Absence of a relationship with variation in texture at pork weight. *Meat Science*, *42*, 355–369.
- Coro, F. A. G., Youssef, E. Y., & Shimokomaki, M. (2002). Age related changes in poultry breast meat collagen pyridinoline and texture. *Journal of Food Biochemistry*, *26*, 533–541.
- Culler, R. D., Parrish, F. C., Jr., Smith, G. C., & Cross, H. R. (1978). Relationship of myofibril fragmentation index to certain chemical, physical, and sensory characteristics of bovine *longissimus* muscle. *Journal of Food Science*, *43*, 1177–1180.
- Goll, D. E., Boehm, M. L., Geesink, G. H., & Thompson, V. F. (1997). What causes postmortem tenderization? In *Proceedings of 50th reciprocal meat conference* (pp. 60). Ames: American Meat Science Association.
- Guarnieri, P. D., Soares, A. L., Olivo, R., Schneider, J. P., Macedo, R. M., Ida, E. I., et al. (2004). Preslaughter handling with water shower spray inhibits PSE (*Pale, Soft, Exudative*) broiler breast meat in a commercial plant. Biochemical and ultrastructural observations. *Journal of Food Biochemistry*, *28*, 269–277.
- Koohmaraie, M. (1996). Biochemical factors regulating the toughening and tenderization processes of meat. *Meat Science*, *43*, 193–201.
- McKee, S. R., Hirschler, E. M., & Sams, A. R. (1997). Physical and biochemical effects associated with tenderization of broiler breast fillets during aging after pre-rigor deboning. *Journal of Food Science*, *62*, 959–962.
- Meilgaard, M., Civille, G. V., & Carr, B. T. (1999). *Sensory evaluation techniques* (3rd ed.). Boca Raton: CRC Press, p. 386.
- Morgan, J. B., Savell, J. W., Hale, D. S., Miller, R. K., Gaiffin, D. B., Cross, H. R., et al. (1991). National beef tenderness survey. *Journal Animal Science*, *69*, 3274–3283.
- Northcutt, J. K. (2001). Preslaughter factors affecting poultry meat quality. In A. R. Sams (Ed.), *Poultry meat processing* (pp. 5–18). Boca Raton: CRC Press.
- Northcutt, J. K., Buhr, R. J., Young, L. L., Lyon, C. E., & Ware, G. O. (2001). Influence of age and postchill carcass aging duration on chicken breast meat. *Poultry Science*, *80*, 808–812.
- Papinaho, P. A., & Fletcher, D. L. (1996). The influence of temperature on broiler breast muscle shortening and extensibility. *Poultry Science*, *75*, 797–802.
- Pearson, A. M. (1987). Muscle function and postmortem changes. In J. F. Price & B. S. Schweigert (Eds.), *The science of meat and meat products* (3rd ed., pp. 307–327). Westport: Food and Nutrition Press, Inc.
- Sams, A. R. (1999). Meat quality during processing. *Poultry Science*, *78*, 798–803.
- Sams, A. R., & Janky, D. M. (1986). The influence of brine chilling on tenderness of hot boned chill-boned and age-boned broiler fillets. *Poultry Science*, *65*, 1316–1321.
- Smith, D. P., Fletcher, D. L., & Papa, C. M. (1992). Duckling and chicken processing yields and breast meat tenderness. *Poultry Science*, *71*, 197–202.
- Taylor, R. G., Gessink, G. H., Thompson, V. F., Koohmaraie, M., & Goll, D. E. (1995). Is Z-disk degradation responsible for post-mortem tenderization? *Journal Animal Science*, *73*, 1351–1367.