

## Effect of $\alpha$ -tocopherol, $\alpha$ -tocotrienol and Rosa mosqueta shell extract on the performance of antioxidant-stripped canola oil (*Brassica* sp.) at high temperature

Nalda Romero<sup>a,\*</sup>, Paz Robert<sup>a</sup>, Lilia Masson<sup>a</sup>, Jaime Ortiz<sup>a</sup>, Karina González<sup>a</sup>, Karla Tapia<sup>a</sup>, Carmen Dobaganes<sup>b</sup>

<sup>a</sup> *Departamento de Ciencia de los Alimentos y Tecnología Química, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Casilla 233, Santiago 1, Chile*

<sup>b</sup> *Instituto de la Grasa (CSIC), Avda Padre Tejero 4, E-41012 Sevilla, Spain*

Received 5 June 2006; received in revised form 20 November 2006; accepted 23 November 2006

### Abstract

The antioxidant effects of tocopherols ( $\alpha$ -tocopherol and  $\alpha$ -tocotrienol) and Rosa mosqueta shell extract added to antioxidant-stripped canola oil (TCO) were evaluated and compared with the non-stripped oil (CO) under the same conditions. Seven oil systems were subjected to thermal treatment at 180 °C for 18 h. Polar compounds formation, degradation of tocopherols and carotenoid pigments were studied. The addition of Rosa mosqueta shell extract gave a great stability to TCO, similar to CO, with a low polar compound formation and a high retention of  $\alpha$ -tocopherol compared with other TCO samples, which suggested the protective action of the minor components present in the extract. Alpha-tocopherol showed a higher effectiveness than  $\alpha$ -tocotrienol at high temperature. However, an increase in the level of  $\alpha$ -tocopherol did not improve its action.

© 2007 Elsevier Ltd. All rights reserved.

**Keywords:** Antioxidant;  $\alpha$ -Tocopherol;  $\alpha$ -Tocotrienol; Canola oil; Carotenoids; Rosa mosqueta; Thermal oxidation

### 1. Introduction

Synthetic antioxidants are being questioned while natural antioxidants such as tocopherol, polyphenols and carotenoid pigments are having a greater relevance in the protection against lipid oxidation. Alpha and gamma tocopherols are the most abundant natural antioxidants in seed oils. Tocopherols act as antioxidant by donating a hydrogen atom to peroxy radicals of unsaturated lipid molecules, forming a hydroperoxide and a tocopheroxy radical, which reacts with other peroxy or tocopheroxy radicals forming more stable adducts (Lampi, Kataja, Kamal-Eldin, & Piironen, 1999).

In processes carried out at high temperatures, such as the process of frying, it is of primary interest to know about the behavior of the natural antioxidants in the oils and to define their significance in connection with oil deterioration (Barrera-Arellano, Ruiz-Méndez, Márquez Ruiz, & Dobaganes, 1999). Some studies have investigated the behavior of tocopherol in different unsaturated oils during frying (Barrera-Arellano, Ruiz-Méndez, Velasco, Márquez-Ruiz, & Dobaganes, 2002; Lampi et al., 1999; Lampi & Kamal-Eldin, 1998; Normand, Eskin, & Przybylski, 2001; Romero et al., 2004).

The use of natural compounds to prevent degradation during frying has been reported earlier, e.g. an ethanol extract from the spice *Satureja hortensis* L. (summer savory) improved the oxidative stability of sunflower oil at frying temperature (Yanishlieva, Marinova, Marekov,

\* Corresponding author. Tel.: +56 02 9781666; fax: +56 02 2227900.  
E-mail address: [nromero@uchile.cl](mailto:nromero@uchile.cl) (N. Romero).

& Gordon, 1997); a rosemary extract showed a protective action against dimer formation and tocopherol degradation in rapeseed oil used in potato frying (Gordon & Kourisma, 1995) and in another study it improved the resistance to oxidative rancidity in a vegetable oil used in deep frying (Lalas & Dourtoglou, 2003).

Carotenoids are present in a wide variety of foods and they are responsible for the red to yellow coloration in fruits and vegetables. The role of pro-vitamin A in some carotenoids has been known for a long time, but carotenoids also have an important antioxidant role in the cellular protection against lipid peroxidation (Basu, Del Vecchio, Flider, & Orthefer, 2001), thus preventing the risk of degenerative illnesses such as cancer, cardiovascular disease and macular degeneration. It also reduces the risk of cataracts and strengthens the immune system (Bermudez, Ribaya-Mercado, Talegawkar, & Tucker, 2005).

Rosa mosqueta (*Rosa rubiginosa*) is widely spread throughout the world. Its fruit has highly interesting carotenoid pigments such as  $\beta$ -carotene, lycopene and rubixanthin; and it can be used as a natural source of carotenoids in food technology (Hornero-Méndez & Minguéz-Mosquera, 2000; Robert, Carlsson, Romero, & Masson, 2003). Robert et al. reported lycopene,  $\beta$ -carotene and rubixanthin as the main carotenoid pigments in commercial shell and fruits (Robert et al., 2003).

So far, there are only a few studies about the antioxidant effect of carotenoid-containing plant extracts at high temperature (Henry, Catignani, & Schwartz, 1998a). Rosa mosqueta shell is an under utilized by-product, that remains after the extraction of the seed from the fruit, for the oil preparation. This by-product rich in carotenoid pigments presents an opportunity for the obtaining of extracts with antioxidant and pigmenting properties and therefore to increase the economic value of the Rose mosqueta and its utilization for any related industries.

The objective of this work was to study the antioxidant effect of  $\alpha$ -tocopherol,  $\alpha$ -tocotrienol and a Rosa mosqueta shell extract on the stability of antioxidant-stripped canola oil at high temperature.

## 2. Materials and methods

### 2.1. Materials

Rapeseed oil (canola variety) or canola oil (*Brassica* sp.) (CO) was supplied by Agromaule S.A. (Santiago, Chile). Commercial dried shells of Rosa mosqueta (*Rosa rubiginosa*) were purchased from Noveltec S.A. (Santiago, Chile). DL- $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ -tocopherols and tocotrienols were obtained by Calbiochem (Merck, Darmstadt, Germany). Solvents used were HPLC-grade or analytical grade.

### 2.2. Samples and treatments

Oil samples. Seven canola oil experimental systems were studied:

- (a) Canola oil (CO).
- (b) Antioxidant-stripped canola oil (canola oil treated with alumina to remove tocopherols) (TCO).
- (c) TCO supplemented with 155 mg/kg of  $\alpha$ -tocopherol (TCO + AT).
- (d) TCO supplemented with 138 mg/kg of  $\alpha$ -tocotrienol (TCO + AT3).
- (e) TCO supplemented with a mixture of 66 mg/kg of  $\alpha$ -tocopherol and 72 mg/kg of  $\alpha$ -tocotrienol (TCO + M).
- (f) TCO supplemented with Rosa mosqueta shell extract (TCO + RME) containing carotenoid pigments (465 mg/kg) and  $\alpha$ -tocopherol (432 mg/kg).
- (g) TCO supplemented with 432 mg/kg of  $\alpha$ -tocopherol (TCO + 432AT).

The levels used of  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol in the samples (c), (d) and (e) were chosen according to previous reports about of their antioxidant effect (Fuster, Lampi, Hopia, & Kamal-Eldin, 1998), considering to study the antioxidant behaviour of each one in independent form and together.

As the RME supplied 432 mg/kg of  $\alpha$ -tocopherol to TCO + RME was necessary to supplement a sample of TCO with 432 mg/kg of AT isolating thus the carotenoid pigments effect.

### 2.3. Preparation of antioxidant-stripped canola oil

Canola oil was removed of its antioxidants via adsorption chromatography using a glass column packed with activated alumina (Merck, Darmstadt, Germany), according to Yoshida, Kondo, and Kajimoto (1992). The absence of tocopherols in the antioxidant-stripped canola oil was confirmed by HPLC, according to AOCS standard method (1993).

### 2.4. Preparation of Rosa mosqueta shell extract

Ground Rosa mosqueta (*Rosa rubiginosa*) shell (3.2 kg) was extracted with hexane (3.2 L) and shaken in multi-wrist shaker at room temperature by 2 h. The hexane containing the carotenoid pigments was filtered by Büchner funnel using vacuum. Three consecutive extractions were made as the cycle described above until the pulp was light red. Extracts were combined; the solvent was removed in a Büchi rotatory evaporator at 40 °C and finally diluted to 500 ml volumetric flask with hexane.

### 2.5. Preparation of TCO supplemented with Rosa mosqueta shell extract

380 g of TCO was mixed with 200 ml of Rosa mosqueta shell extract. The solvent was then evaporated; first in a Büchi rotatory evaporator at 40 °C and then under a stream of nitrogen.

## 2.6. Thermal oxidation assays

Standard Rancimat tubes containing 10 g of oil were heated at  $180 \pm 1$  °C in a Rancimat apparatus (Metrohm Ltd, Herisau, Switzerland). Air was not bubbled during heating and the tubes were left open. Independent tubes containing TCO + RME and TCO + 432AT were heated for 0.5, 1, 1.5, 2, 8 and 18 h, while the rest of the samples were heated for 2, 4, 6, 8, 10, 14 and 18 h. Thermal oxidation was carried out in triplicate.

## 2.7. Fatty acid composition

Fatty acids were analyzed as methyl esters derivatives (AENOR, 1991) by gas chromatography (GLC) using a HP 5890 chromatograph (Hewlett-Packard, Palo Alto, CA, USA). A fused silica capillary column BPX70 (50 m, 0.25  $\mu$ m film; SGE, Incorporated, Austin, TX, USA) was used. Temperature increase was programmed between 160 and 230 °C at 2 °C/min, and samples were run with hydrogen as carrier gas. Standard fatty acid methyl esters (FAME) from Merck (Merck, Darmstadt, Germany) were used for identification purposes.

## 2.8. Analysis of tocopherols and tocotrienols

Tocols were determined by high-performance liquid chromatography (HPLC) with fluorescence detection, according to AOCS standard method (1993) for non-esterified tocopherols and tocotrienols. The HPLC system consisted of a Merck-Hitachi L-6200 A pump (Merck, Darmstadt, Germany) equipped with a Rheodyne 7725i injector with 20  $\mu$ l sample loop, a Merck-Hitachi F-1050 fluorescence detector and a Merck-Hitachi D-2500 chromatographic integrator. Peaks were detected at 290 nm (excitation wavelength) and 330 nm (emission wavelength). A LiChro-CART Superspher Si 60 column (25 cm  $\times$  4 mm id, 5  $\mu$ m particle size; Merck, Darmstadt, Germany) was used. The mobile phase was propan-2-ol in hexane (0.5:99.5 v/v) at a flow rate of 1 ml/min. Tocols were identified and quantified using Calbiochem tocopherols and tocotrienols (Merck, Darmstadt, Germany) as external standards.

## 2.9. Analysis of polar compounds

Quantification and distribution of polar compounds (PC) were determined by a combination of adsorption column chromatography and high-performance size exclusion chromatography (HPSEC) according to Dobarganes, Perez-Camino, and Marquez-Ruiz (1988). The HPLC system consisted of a Merck-Hitachi L-6200 A pump, a 20  $\mu$ l injection loop, a Merck RI-71 refractive index detector and a Merck-Hitachi D-2500 chromatographic integrator. The separation was performed on two serial 500 and 100 Å columns (PLGEL, 30 cm  $\times$  0.8 cm id, 5  $\mu$ m particle size; Hewlett Packard, Amherst, MA, USA). The mobile phase was tetrahydrofuran at a flow rate of 1 ml/min. Identification was

performed in agreement with Dobarganes et al. (1988) and the quantification of each compound was based on peak areas assuming equal detector response.

## 2.10. Oxidative stability

Induction period (IP) was determined using a Rancimat Oxidative Stability Instrument (Metrohm Ltd., Herisau, Switzerland) at 100 °C and an air flow of 20 l/h, according to AOCS standard method (1993).

## 2.11. Analysis of carotenoids

Pigment extraction from oils was carried out according to Henry, Catignani, and Schwartz (1998b). Carotenoid analysis was determined by HPLC. The HPLC system consisted of a Merck-Hitachi L-6200 pump, a Waters 996 Photodiode Array detector coupled to a computer with Millennium 32 software and a Waters symmetry column (C18, 5  $\mu$ m particle size, 4.6 mm id  $\times$  25 cm; Waters, Milford, MA). The isocratic mobile phase was methanol:acetonitrile:ethyl acetate (20:65:15 v/v) at a flow rate of 1 ml/min. Carotenoids were detected at 450 nm. Identification and quantification were carried out using external standards. Standards for *all-trans*- $\beta$ -carotene, *all-trans*-lycopene and *all-trans*-rubixanthin were obtained from carrots, tomatoes and Rosa mosqueta shells, respectively. The pigments were purified by open column chromatography, as described by Rodriguez-Amaya (1999). Concentrations of standards were determined by spectrophotometry using their respective  $A_{1\text{cm}}^{1\%}$ . Calibration curves were determined for each compound. *Cis*-isomers were identified according to elution order as reported by Henry et al. (1998b).

## 2.12. Statistical analysis

The linear regression (95% confidence limit) was used to determine the reaction order and degradation rate constant of tocols. To determine statistical differences in the formation of PC and their distribution a multivariate ANOVA was performed considering treatment and time as independent variables and PC as dependent variable. Induction period and PC for a same time were analyzed by one factor ANOVA. These statistical tests were carried out by using Statgraphics Plus, version 7.0 (Manugistics Inc., Statistical Graphics Corporation, Rockville, MD).

## 3. Results and discussion

Table 1 shows the concentrations of major carotenoid pigments and tocopherols in Rosa mosqueta shell extract (RME) alone, and in antioxidant-stripped canola oil with added RME extract, (TCO + RME). Lycopene,  $\beta$ -carotene and rubixanthin were found in RME, being *trans*- $\beta$ -carotene and *trans*-rubixanthin the main carotenoid pigments. RME had  $\alpha$ -tocopherol too; therefore the addition of RME to

Table 1

Composition of major carotenoid pigments and tocopherols in Rosa mosqueta (*Rosa rubiginosa*) shell extract (RME) and in antioxidant-stripped canola oil with added RME extract (TCO + RME)

	RME ( $\mu\text{g/ml}$ )	TCO + RME (mg/kg)
<i>trans</i> -Rubixanthin	207	125
<i>cis</i> -Rubixanthin	146	87
<i>trans</i> -Lycopene	81	42
<i>cis</i> -Lycopene	55	33
<i>trans</i> - $\beta$ -Carotene	251	144
<i>cis</i> - $\beta$ -Carotene	85	34
$\alpha$ -Tocopherol	825	432

TCO supplied 432 mg/kg of  $\alpha$ -tocopherol (TCO + RME), then it was necessary to add AT (432 mg/kg) to antioxidant-stripped canola oil (TCO + 432AT) for to isolate its effect.

Table 2 shows the fatty acid composition, the tocopherols, the total polar compounds and the polar compound distribution in the initial CO and TCO. CO consisted of totally 61% monounsaturated fatty acids and the major fatty acid was oleic acid (total content 56%). In addition, CO had a low content of saturated fatty acids (8.1%) and a medium content of polyunsaturated fatty acids (30.7%), mainly represented by linoleic and linolenic acids (21.5% and 8.0%, respectively).

CO showed a higher concentration in total tocopherols than the content previously reported by Normand et al. (2001) and Barrera-Arellano et al. (2002). In CO,  $\alpha$  and  $\gamma$ -tocopherols were the main natural antioxidants present, with contents of 247 and 348 mg/kg, respectively.

TCO did not show important changes in its fatty acid composition compared to CO. However, it can be observed that the treatment with alumina not only removed the tocopherols, but also the polar compounds (PC) present in CO.

Oxidative stability results are shown in Table 3. A high induction period (IP) of 19.6 h at 100 °C was found for CO, which agreed with its high levels of monounsaturated fatty acids and tocopherols. Barrera-Arellano et al. (2002) found an IP value of 10.4 h for rapeseed oil under similar conditions.

The IP value in TCO was 1.4 h and reflected the importance of antioxidative compounds for oil stability. The addition of tocopherols and Rosa mosqueta shell extract to TCO improved the oxidative stability of the oil ( $P < 0.05$ ). At similar concentrations of tocopherols not significant difference in stability was observed. The Rosa mosqueta shell extract gave the highest stability to treated oil compared with the other treated samples ( $P < 0.05$ ) in the accelerated test, but did not reach the oxidative stability of CO.

### 3.1. Canola oil thermal oxidation assays

Table 4 shows the tocopherol degradation and the polar compound formation in all the oil systems studied. It can be observed that tocopherols remained in CO during the

Table 2

Fatty acid composition, tocopherols, total polar compounds and polar compound distribution of canola oil (CO) and antioxidant-stripped canola oil (TCO)

	CO	TCO
<i>Fatty acid (%)</i>		
Myristic (14:0)	0.1	0.1
Palmitic (16:0)	4.8	4.8
Palmitoleic (16:1)	tr	tr
Palmitoleic isomer (16:1)	0.2	0.2
Heptadecanoic (17:0)	0.1	0.1
Heptadecenoic (17:1)	0.1	tr
Stearic (18:0)	2.1	1.9
Octadecenoic isomer (18:1)	0.2	tr
Oleic (18:1w9)	56.5	55.8
Octadecenoic isomer (18:1w7)	2.6	3.2
Octadecadienoic isomer (18:2)	0.3	0.2
Linoleic (18:2w6)	21.5	21.6
Octadecatrienoic isomer (18:3)	0.9	0.7
Linolenic (18:3w3)	8.0	8.6
Arachidic (20:0)	0.6	0.6
Eicosenoic (20:1)	1.3	1.3
Eicosenoic isomer (20:1)	0.1	ND
Behenic (22:0)	0.4	0.4
Docosenoic (22:1)	tr	0.3
Tetracosanoic (24:0)	0.1	0.1
Tetracosenoic (24:1)	0.1	0.1
Total saturated	8.2	8.0
Total monounsaturated	61.1	60.9
Total polyunsaturated	30.7	31.1
<i>Tocopherols (mg/kg)</i>		
Alpha tocopherol	247	ND
Gamma tocopherol	348	ND
Delta tocopherol	16	ND
<i>Polar compound (wt%) and polar compound distribution (wt%)</i>		
PC	2.6	0.1
TGP	ND	ND
TGD	0.2	ND
oxTGM	1.1	0.1
DG	0.8	ND
FA	0.5	ND

Abbreviations: tr, trace; ND, not detected; PC, total polar compounds; TGP, triacylglycerol polymers; TGD, triacylglycerol dimers; oxTGM, oxidized triacylglycerol monomers; DG, diacylglycerols; FA, free fatty acids.

Table 3

Oxidative stability of canola oil (CO), antioxidant-stripped canola oil (TCO), antioxidant-stripped canola oil with addition of 155 mg/kg of  $\alpha$ -tocopherol (TCO + AT), 138 mg/kg of  $\alpha$ -tocotrienol (TCO + AT3),  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol mixture (TCO + M), Rosa mosqueta shell extract (TCO + RME) and 432 mg/kg of  $\alpha$ -tocopherol (TCO + 432AT)

Oils	Induction period (IP) (h)
CO	19.6 $\pm$ 0.6 <sup>c</sup>
TCO	1.4 $\pm$ 0.0 <sup>a</sup>
TCO + AT	6.9 $\pm$ 0.4 <sup>b</sup>
TCO + AT3	7.6 $\pm$ 0.3 <sup>b</sup>
TCO + M	7.0 $\pm$ 0.4 <sup>b</sup>
TCO + RME	9.5 $\pm$ 0.1 <sup>d</sup>
TCO + 432AT	8.7 $\pm$ 0.4 <sup>c</sup>

Values are reported as means  $\pm$  SD ( $n = 3$ ).

Different letters for IP mean significant differences between samples ( $P < 0.05$ ).

Table 4

Loss of tocopherols (mg/kg) and formation of polar compounds (wt%) in canola oil (CO), antioxidant-stripped canola oil (TCO), antioxidant-stripped canola oil with addition of  $\alpha$ -tocopherol (TCO + AT),  $\alpha$ -tocotrienol (TCO + AT3),  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol mixture (TCO + M), Rosa mosqueta shell extract (TCO + RME) and 432 mg/kg of  $\alpha$ -tocopherol (TCO + 432AT)

Samples/heating period (h)		0	2	4	6	8	10	14	18
CO	AT	247 ± 3	234 ± 3	–	–	196 ± 4	–	172 ± 7	168 ± 8
	GT	348 ± 7	322 ± 10	–	–	288 ± 3	–	270 ± 6	257 ± 8
	DT	16 ± 0.4	15 ± 0.3	–	–	14 ± 0.6	–	14 ± 0.1	14 ± 0.2
	PC <sup>a</sup>	2.6 ± 0.2	4.5 ± 0.3	–	–	6.8 ± 0.2	–	8.7 ± 0.3	9.3 ± 0.4 <sup>d</sup>
TCO	PC <sup>c</sup>	0.1 ± 0.1	4.5 ± 0.2	7.3 ± 0.8	10.9 ± 0.6	14.1 ± 0.7	17.1 ± 0.7	23.7 ± 0.2	30.7 ± 0.5 <sup>j</sup>
	AT	155 ± 1	129 ± 3	87 ± 3	48 ± 2	28 ± 3	15 ± 1	ND	ND
TCO + AT	PC <sup>ab</sup>	0.1 ± 0.1	3.7 ± 0.4	5.3 ± 0.4	6.4 ± 0.1	8.1 ± 0.1	10.9 ± 0.4	13.3 ± 0.4	16.2 ± 0.4 <sup>f</sup>
	AT3	138 ± 1	43 ± 2	26 ± 2	tr	tr	ND	ND	ND
TCO + AT3	PC <sup>c</sup>	0.1 ± 0.1	4.2 ± 0.4	6.5 ± 0.3	9.0 ± 0.8	13.3 ± 0.4	16.4 ± 0.5	23.1 ± 0.6	29.3 ± 0.3 <sup>i</sup>
	AT	66 ± 2	31 ± 1	12 ± 1	tr	tr	ND	ND	ND
TCO + M	AT3	72 ± 1	28 ± 1	10 ± 1	tr	tr	ND	ND	ND
	PC <sup>c</sup>	0.1 ± 0.1	3.2 ± 0.4	6.7 ± 0.5	7.9 ± 0.3	12.7 ± 0.5	16.8 ± 0.7	19.8 ± 0.5	27.9 ± 0.1 <sup>h</sup>
		0	0.5	1	1.5	2	8	18	
TCO + RME	AT	432 ± 4	329 ± 5	325 ± 2	314 ± 3	303 ± 9	213 ± 7	62 ± 8	
	PC <sup>ab</sup>	0.1 ± 0.3	2.4 ± 0.3	3.0 ± 0.1	3.6 ± 0.2	4.6 ± 0.2	5.6 ± 0.1	12.4 ± 0.2 <sup>e</sup>	
TCO + 432AT	AT	432 ± 3	294 ± 6	255 ± 4	237 ± 1	174 ± 3	14 ± 1	ND	
	PC <sup>bc</sup>	0.1 ± 0.1	0.7 ± 0.0	1.6 ± 0.1	2.9 ± 0.1	3.9 ± 0.2	13.9 ± 0.3	23.9 ± 0.5 <sup>e</sup>	

Abbreviations: AT, alpha tocopherol; AT3, alpha tocotrienol; GT, gamma tocopherol; DT, delta tocopherol; PC, polar compounds; ND, not detected; tr,  $\leq 10$  mg/kg. Values are reported as means  $\pm$  SD ( $n = 3$ ). Different letters for PC mean significant differences between samples ( $P < 0.05$ ).

whole thermal oxidation assay. After 18 h 70% of total tocopherols were remaining which was concordant with the high value for IP, as commented above. The result for tocols in CO was in agreement with the data published by Normand et al. (2001) on the frying stability of regular canola oil which presented a high level of tocopherols at 20 h of frying.

Fig. 1 shows the logarithm of the percent retention vs. time (h) for  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol in six samples.




Fig. 1. Pseudo first-order degradation plots for  $\alpha$ -tocopherol or  $\alpha$ -tocotrienol at 180 °C in canola oil (CO), antioxidant-stripped canola oil with addition of  $\alpha$ -tocopherol (TCO + AT),  $\alpha$ -tocotrienol (TCO + AT3),  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol mixture (TCO + M), Rosa mosqueta shell extract (TCO + RME) and 432 mg/kg of  $\alpha$ -tocopherol (TCO + 432AT). Different letters mean significant differences between samples for degradation rate constants ( $k$ ) ( $P < 0.05$ ).

The degradation of  $\alpha$ -tocopherol and  $\alpha$ -tocotrienol followed pseudo first-order kinetics in all the samples. The correlation coefficient was used as a parameter to determine the reaction order. The degradation rate constants ( $k$ ) were obtained from the slopes of the plots of Fig. 1 (Robert, Romero, Ortiz, Masson, & Barrera-Arellano, 2006). The degradation rates for  $\alpha$ -tocopherol varied greatly between the different samples containing the oil treated with alumina ( $P < 0.05$ ). Alpha-tocopherol showed a lower degradation rate ( $P < 0.05$ ) when it was added as the only antioxidant in TCO + AT ( $0.27 \pm 0.004$  h<sup>-1</sup>,  $r^2 = 0.995$ ) than when it was added together with  $\alpha$ -tocotrienol in TCO + M ( $0.44 \pm 0.01$  h<sup>-1</sup>,  $r^2 = 0.999$ ). This same behaviour was observed and explained in a previous research in Chilean hazelnut oil (Romero et al., 2004). Degradation rate of AT was lower than AT3 ( $P < 0.05$ ) as can be observed in samples TCO + AT3 ( $0.54 \pm 0.03$  h<sup>-1</sup>,




Fig. 2. Degradation of carotenoid pigments in antioxidant-stripped canola oil with added Rosa mosqueta shell extract (TCO + RME) during heating at 180 °C. (a) *trans*-isomers; (b) *cis*-isomers.

$r^2 = 0.993$ ) and TCO + M ( $0.53 \pm 0.02 \text{ h}^{-1}$ ,  $r^2 = 0.994$ ), Fig. 1, showing a higher stability of AT than AT3 at high temperature. The increase of the  $\alpha$ -tocopherol level accelerated its loss in TCO + 432AT ( $0.42 \pm 0.01 \text{ h}^{-1}$ ,  $r^2 = 0.996$ ), being a degradation rate similar to that obtained in TCO + M ( $P > 0.05$ ). This is in agreement with another study, in which high concentration of  $\alpha$ -tocopherol added to purified sunflower triacylglycerols showed lower effectiveness (Fuster et al., 1998). Alpha-tocopherol showed the highest stability in TCO + RME with 14% of residual tocopherols after 18 h of heating at 180 °C, whereas only a 3% of this antioxidant was found in TCO + 432AT after 8 h of heating. These two oil systems (TCO + RME and TCO + 432AT) had the same initial content of AT; however TCO + RME showed higher retention of this antioxidant ( $k$ :  $0.10 \pm 0.01 \text{ h}^{-1}$ ,  $r^2 = 0.999$ ). These results could be explained by the presence of carotenoid pigments or other antioxidant compounds in the Rosa mosqueta shell extract.

Fig. 2 shows the degradation of carotenoid pigments in TCO + RME at a high temperature (180 °C). As it can be observed, the carotenoid pigments abruptly decayed in the first half hour of heating, and after two hours only a 13% of  $\beta$ -carotene remained in the oil. An increase of the isomer 13-*cis*  $\beta$ -carotene was observed in the first half hour of heating accompanied by a decrease in the isomer *trans*- $\beta$ -carotene. Henry et al. (1998b) described a similar behaviour for  $\beta$ -carotene. This behaviour was not observed for lycopene and rubixanthin, suggesting the absence of an equilibrium *trans*–*cis* reaction, or that this reaction was very fast due to the high temperature. Alpha-tocopherol

and  $\gamma$ -tocopherol have been reported as protective compounds in the degradation of  $\beta$ -carotene (Terao, Yamachi, Murakami, & Matsushita, 1980), lutein and lycopene (Haila, Lievonon, & Heinonen, 1996) in food lipids. The results obtained in our study did not show this protective effect; however as it was mentioned above, a higher retention of  $\alpha$ -tocopherol in TCO + RME was observed, suggesting an interaction of  $\alpha$ -tocopherol with other compounds present in the Rosa mosqueta shell extract.

Polar compounds (PC) gave a good measurement of the lipid deterioration. The formation of PC in CO (Table 4) reached lower values than the other treatments ( $P < 0.05$ ) at the end of heating period, owing to the protective action of their tocopherols. Furthermore, it was one of the most thermally stable oil and its behaviour was comparable to TCO + AT and TCO + RME ( $P > 0.05$ ). The PC in TCO + AT3, TCO + M and TCO + 432AT increased more rapidly due to the early loss of tocopherols in the systems (6–8 h), not showing significant differences between the treatments ( $P > 0.05$ ) and with a behaviour similar to TCO. When one was compared different treatments at 18 h of heating significant differences were observed for all the samples  $P < 0.05$ . These results could be explained because the greater differences in PC values between the samples began around of 8 h of heating, when the loss of tocopherols was almost complete.

TCO + RME did not show significant differences during all period of heating respect TCO + AT and TCO + 432AT ( $P > 0.05$ ), although significant differences in PC values at 18 h of heating in these three treatments were observed ( $P < 0.05$ ), showing a lower value for TCO + RME which

Table 5  
Changes in polar compound distribution (wt%) in the same samples as named in Table 4

Samples/heating period (h)		0	2	4	6	8	10	14	18
CO	oxTGM <sup>a</sup>	1.1 ± 0.1	1.5 ± 0.2			2.7 ± 0.1		3.2 ± 0.0	3.5 ± 0.1 <sup>j</sup>
	TGD <sup>d</sup>	0.2 ± 0.1	1.3 ± 0.2			1.9 ± 0.1		2.9 ± 0.1	3.0 ± 0.2 <sup>o</sup>
	TGP <sup>g</sup>	ND	ND			0.6 ± 0.1		1.0 ± 0.1	1.3 ± 0.1 <sup>u</sup>
TCO	oxTGM <sup>c</sup>	0.1 ± 0.1	1.9 ± 0.1	3.1 ± 0.4	4.2 ± 0.5	5.7 ± 0.3	6.7 ± 0.3	9.3 ± 0.8	11.0 ± 0.3 <sup>n</sup>
	TGD <sup>f</sup>	ND	1.8 ± 0.1	3.1 ± 0.3	4.7 ± 0.2	6.1 ± 0.2	6.9 ± 0.3	9.3 ± 0.3	11.3 ± 0.3 <sup>i</sup>
	TGP <sup>i</sup>	ND	0.8 ± 0.2	1.1 ± 0.1	2.0 ± 0.1	2.3 ± 0.3	3.5 ± 0.4	5.1 ± 0.8	8.4 ± 0.7 <sup>x</sup>
TCO + AT	oxTGM <sup>ab</sup>	0.1 ± 0.1	2.0 ± 0.2	2.9 ± 0.0	2.9 ± 0.1	3.5 ± 0.3	4.7 ± 0.1	5.5 ± 0.1	6.6 ± 0.2 <sup>l</sup>
	TGD <sup>e</sup>	ND	1.2 ± 0.1	1.8 ± 0.2	2.7 ± 0.1	3.5 ± 0.1	4.6 ± 0.3	5.9 ± 0.3	6.8 ± 0.3 <sup>q</sup>
	TGP <sup>gh</sup>	ND	0.5 ± 0.1	0.6 ± 0.1	0.8 ± 0.2	1.1 ± 0.3	1.6 ± 0.1	1.9 ± 0.1	2.8 ± 0.2 <sup>v</sup>
TCO + AT3	oxTGM <sup>c</sup>	0.1 ± 0.1	2.3 ± 0.2	3.0 ± 0.1	3.6 ± 0.3	6.4 ± 0.2	7.1 ± 0.1	9.6 ± 0.3	11.3 ± 0.1 <sup>n</sup>
	TGD <sup>f</sup>	ND	1.5 ± 0.1	2.3 ± 0.2	3.8 ± 0.3	5.1 ± 0.2	6.4 ± 0.2	8.5 ± 0.3	9.9 ± 0.3 <sup>s</sup>
	TGP <sup>i</sup>	ND	0.4 ± 0.1	1.2 ± 0.1	1.6 ± 0.3	1.8 ± 0.2	2.9 ± 0.4	5.0 ± 0.2	8.1 ± 0.2 <sup>x</sup>
TCO + M	oxTGM <sup>bc</sup>	0.1 ± 0.1	1.8 ± 0.1	2.9 ± 0.1	3.8 ± 0.4	5.1 ± 0.1	6.7 ± 0.2	7.5 ± 0.4	8.8 ± 0.2 <sup>n</sup>
	TGD <sup>f</sup>	ND	0.9 ± 0.1	2.9 ± 0.1	3.1 ± 0.5	5.4 ± 0.3	6.9 ± 0.4	8.1 ± 0.4	9.5 ± 0.1 <sup>rs</sup>
	TGP <sup>i</sup>	ND	0.5 ± 0.2	0.9 ± 0.2	1.0 ± 0.2	2.2 ± 0.2	3.2 ± 0.2	4.2 ± 0.2	9.6 ± 0.4 <sup>y</sup>
		0	0.5	1.0	1.5	2.0	8	18	
TCO + RME	oxTGM <sup>ab</sup>	0.1 ± 0.1	2.1 ± 0.2	2.5 ± 0.1	2.9 ± 0.2	2.7 ± 0.1	2.7 ± 0.2	5.1 ± 0.2 <sup>k</sup>	
	TGD <sup>de</sup>	ND	0.3 ± 0.2	0.5 ± 0.1	0.7 ± 0.0	1.9 ± 0.2	2.4 ± 0.1	5.5 ± 0.2 <sup>p</sup>	
	TGP <sup>gh</sup>	ND	ND	ND	ND	ND	0.5 ± 0.1	1.8 ± 0.1 <sup>uv</sup>	
TCO + 432AT	oxTGM <sup>bc</sup>	0.1 ± 0.1	0.4 ± 0.0	0.8 ± 0.2	1.3 ± 0.2	1.9 ± 0.1	5.6 ± 0.2	8.3 ± 0.1 <sup>m</sup>	
	TGD <sup>ef</sup>	ND	0.3 ± 0.0	0.7 ± 0.1	1.3 ± 0.2	1.5 ± 0.1	5.9 ± 0.1	8.9 ± 0.1 <sup>t</sup>	
	TGP <sup>hi</sup>	ND	ND	0.1 ± 0.2	0.3 ± 0.3	0.5 ± 0.1	2.4 ± 0.2	6.7 ± 0.4 <sup>w</sup>	

Abbreviations: oxTGM, oxidized triacylglycerol monomers; TGD, triacylglycerol dimers; TGP, triacylglycerol polymers. Values are reported as means ± SD ( $n = 3$ ). Different letters for the same parameter mean significant differences between samples ( $P < 0.05$ ).

was in accordance with a higher permanence of AT in its system.

Table 3 showed an IP for TCO + AT lower than TCO + 432AT ( $P < 0.05$ ), suggesting differences in behaviour at high temperatures and in the conditions of the Rancimat test.

The results obtained for the distribution of polar compounds are shown in Table 5. All distribution species including oxidized triacylglycerol monomers (oxTGM), triacylglycerol dimers (TGD) and triacylglycerol polymers (TGP) increased continuously during heating in most of the oil systems. In general, low levels of polymers were found in oils with greater retention of tocopherols ( $P < 0.05$ ). TCO + AT3 and TCO + M showed a similar behaviour to TCO ( $P > 0.05$ ). Rosa mosqueta shell extract gave the highest stability to antioxidant-stripped canola oil with the lowest values of polar compounds and distribution species at 18 h ( $P < 0.05$ ).

In conclusion, the addition of Rosa mosqueta shell extract gave a great stability to TCO similar to CO which suggested the protective action of the minor components present in the extract. Alpha-tocopherol showed a high effectiveness as antioxidant at high temperature, however, an increase in the level of  $\alpha$ -tocopherol did not improve its action.

## Acknowledgement

This study was supported by FONDECYT (project 1011070).

## References

- AENOR. (1991). *Asociación Española de Normalización, Norma UNE 55037-73*, Catálogo de Normas UNE, Madrid.
- American Oil Chemists Society (1993). *Official methods and recommended practices of the American oil chemists society* (4th ed.). Champaign, IL: AOCS Press.
- Barrera-Arellano, D., Ruiz-Méndez, V., Márquez Ruiz, G., & Dobarganes, C. (1999). Loss of tocopherols and formation of degradation compounds in triacylglycerol model systems heated at high temperature. *Journal of the Science of Food and Agriculture*, 79, 1–6.
- Barrera-Arellano, D., Ruiz-Méndez, V., Velasco, J., Márquez-Ruiz, G., & Dobarganes, C. (2002). Loss of tocopherols and formation of degradation compounds at frying temperature in oils differing in degree of unsaturation and natural antioxidant content. *Journal of the Science of Food and Agriculture*, 82, 1696–1702.
- Basu, H., Del Vecchio, A., Flider, F., & Orthefer, F. (2001). Nutritional and potential disease prevention properties of carotenoids. *Journal of the American Oil Chemists Society*, 78, 665–675.
- Bermudez, O., Ribaya-Mercado, J., Talegawkar, S., & Tucker, K. (2005). Hispanic and non-hispanic white elders from Massachusetts have different patterns of carotenoid intake and plasma concentrations. *Journal of Nutrition*, 135, 1496–1502.
- Dobarganes, M. C., Perez-Camino, M. C., & Marquez-Ruiz, G. (1988). High performance size exclusion chromatography of polar compounds in heated and non-heated fats. *Fat Science Technology*, 90, 308–311.
- Fuster, M. A., Lampi, A., Hopia, A., & Kamal-Eldin, A. (1998). Effects of  $\alpha$ - and  $\gamma$ -tocopherols on the autoxidation of purified sunflower triacylglycerols. *Lipids*, 33, 715–722.
- Gordon, M. H., & Kourisma, L. (1995). The effects of antioxidants on changes in oils during heating and deep frying. *Journal of the Science of Food and Agriculture*, 68, 347–353.
- Haila, K., Lievonen, S., & Heinonen, M. (1996). Effects of lutein, lycopene, annatto and  $\gamma$ -tocopherol on autooxidation of triglycerides. *Journal of Agricultural and Food Chemistry*, 44, 2096–2100.
- Henry, L. K., Catignani, G. L., & Schwartz, S. J. (1998a). The influence of carotenoids and tocopherols on the stability of safflower seed oil during heat-catalyzed oxidation. *Journal of the American Oil Chemists Society*, 75, 1399–1402.
- Henry, L. K., Catignani, G. L., & Schwartz, S. J. (1998b). Oxidative degradation kinetics of lycopene, lutein and 9-*cis* and all-*trans*- $\beta$ -carotene. *Journal of the American Oil Chemists Society*, 75, 823–828.
- Hornero-Méndez, D., & Minguez-Mosquera, I. (2000). Carotenoid pigments in Rosa mosqueta Hips, an alternative carotenoid source for foods. *Journal of Agricultural and Food Chemistry*, 48, 825–828.
- Lalas, S., & Dourtoglou, V. (2003). Use of rosemary extract in preventing oxidation during deep-fat frying of potato chips. *Journal of the American Oil Chemists Society*, 80, 579–583.
- Lampi, A., & Kamal-Eldin, A. (1998). Effect of  $\alpha$ - and  $\gamma$ -tocopherols on thermal polymerization of purified high-oleic sunflower triacylglycerols. *Journal of the American Oil Chemists Society*, 75, 1699–1703.
- Lampi, A., Kataja, L., Kamal-Eldin, A., & Piironen, V. (1999). Antioxidant activities of  $\alpha$ - and  $\gamma$ -tocopherols in the oxidation of rapeseed oil triacylglycerols. *Journal of the American Oil Chemists Society*, 76, 749–755.
- Normand, L., Eskin, N., & Przybylski, R. (2001). Effect of tocopherols on the frying stability of regular and modified canola oils. *Journal of the American Oil Chemists Society*, 78, 369–373.
- Robert, P., Carlsson, R. M., Romero, N., & Masson, L. (2003). Stability of spray-dried encapsulated carotenoid pigments from rosa mosqueta (*Rosa rubiginosa*) oleoresin. *Journal of the American Oil Chemists Society*, 80, 1115–1120.
- Robert, P., Romero, N., Ortiz, J., Masson, L., & Barrera-Arellano, D. (2006). Effect of Rosa mosqueta (*Rosa rubiginosa*) extract on the performance of Chilean hazelnut oil (*Genuina avellana Mol*) at high temperature. *Journal of the American Oil Chemists Society*, 83, 691–695.
- Rodriguez-Amaya, D. A. (1999). *Guide to carotenoid analysis in food*. Washington, DC: ILSI Press.
- Romero, N., Robert, P., Masson, L., Ortiz, J., Pavez, J., Garrido, C., et al. (2004). Effect of  $\alpha$ -tocopherol and  $\gamma$ -tocotrienol on the performance of Chilean hazelnut oil (*Genuina avellana Mol*) at high temperature. *Journal of the Science of Food and Agriculture*, 84, 943–948.
- Terao, J., Yamauchi, R., Murakami, H., & Matsushita, S. (1980). Inhibitory effect of tocopherols and  $\beta$ -carotene on singlet oxygen-initiated photooxidation of methyl linoleate and soybean oil. *Journal of Food Processing and Preservation*, 4, 79–93.
- Yanishlieva, N. V., Marinova, E. M., Marekov, I. N., & Gordon, M. H. (1997). Effect of an ethanol extract from summer savory (*Satureja hortensis* L.) on the stability of sunflower oil at frying temperature. *Journal of the Science of Food and Agriculture*, 74, 524–530.
- Yoshida, H., Kondo, I., & Kajimoto, G. (1992). Participation of free fatty acid in the oxidation of purified soybean oil during microwave heating. *Journal of the American Oil Chemists Society*, 69, 1136–1140.