

## Lipid composition of the liver oil of the ray, *Himantura bleekeri*

Sandrine Le Néchet, Nolwenn Dubois, Jean-Paul Gouyguo, Jean-Pascal Bergé\*

Département Sciences et Techniques Alimentaires Marines, IFREMER Centre de Nantes, BP 21105, 44311 Nantes Cedex 03, France

Received 12 September 2006; received in revised form 22 November 2006; accepted 12 December 2006

### Abstract

If rays are traditionally fished for their caudal fins, the rest of the body is wasted, except part of the skin that can be transformed into leather. Liver oil of the ray, *Himantura bleekeri*, was characterized in terms of lipid class composition and fatty acid profile. Liver oil content was high and represented 54% of the liver weight (w/w). Neutral lipids were predominant (92%), major components being triacylglycerols (63%). Other neutral lipids identified were hydrocarbons, sterol esters, mono- and di-acylglycerols, free fatty acids, sterols and glyceryl ethers. Polyunsaturated fatty acids of the *n*-3 series, namely eicosapentaenoic and docosahexaenoic acids, were high (4% and 16%, respectively). Cholesterol was the major component in the sterol fractions (free or esterified). Hydrocarbons were up to 30 carbons and squalene was present at the level of 22%. Thus, this liver oil proved to be an adequate source of *n*-3 fatty acids and other valuable lipidic compounds.

© 2007 Elsevier Ltd. All rights reserved.

**Keywords:** Ray liver oil; Fatty acids; Lipids; PUFA; EPA; DHA; Squalene; Sterol

### 1. Introduction

The traditional Inuit diet is known to be rich in long-chain *n*-3 polyunsaturated fatty acids (PUFAs) and believed to account for the low incidence of cardiovascular disease in those populations (Mnari et al., 2007). This observation is a starting point for the increasing interest in these fatty acids (Bergé & Barnathan, 2005). Indeed, PUFAs, and particularly eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), are nowadays recognized to prevent inflammatory conditions, arrhythmias, hypertension and triacylglycerolemia, atherosclerosis, autoimmune disorders (Mnari et al., 2007) and to display a variety of beneficial effects in areas ranging from foetal development to cancer prevention (Connor, 2000).

Fish oils are the main commercial sources of EPA and DHA. However, oils from elasmobranchs (particularly the sharks) also contain large quantities of squalene, a 30-carbon isoprenoid. As this compound occurs naturally

in the surface of the human skin (Navarro-Garcia, Pacheco-Aguilar, Vallejo-Cordova, Ramirez-Suarez, & Bolanos, 2000), it is currently used in cosmetics, pharmaceuticals and sunscreens (Wetherbee & Nichols, 2000).

Among elasmobranchs, ray fishes are traditionally caught around the world. However, only a few parts are eaten, the caudal fins, and most of the rest is considered as waste (body, viscera, skin). Few authors have studied the lipid composition of those by-products, notably liver and gonads, but they contain a high proportion of lipids (Navarro-Garcia, Bringas-Alvarado, Pacheco-Aguilar, & Ortega-Garcia, 2004a; Navarro-garcia, Pacheco-Aguilar, Bringas-Alvarado, & Ortega-Garcia, 2004b; Ould El Kebir, Barnathan, Siau, Miralles, & Gaydou, 2003; Pal, Banerjee, Patra, Patra, & Ghosh, 1998).

As indicated by Navarro-garcia et al. (2004b), little information has been reported for ray liver oil. In the present investigation, the lipids and fatty acids of the liver of the commercial ray, *Himantura bleekeri*, were studied for the first time. This ray belongs to the family *Dasyatidae* of the class *Elasmobranchii*. This cartilaginous fish lives in the Indo-pacific ocean and is traditionally fished for flesh and skin (to produce leather) (Blyth, 1860).

\* Corresponding author. Tel.: +33 2 40 37 40 79; fax: +33 2 40 37 40 71.  
E-mail address: [jpberge@ifremer.fr](mailto:jpberge@ifremer.fr) (J.-P. Bergé).

Results are intended to be used to explore the commercial potential of such biomass as a source of marine oils and particularly PUFAs.

## 2. Materials and methods

### 2.1. Sample collection

*H. bleekeri* was collected on the east coast of India (Chennai), during February 2004. Liver was isolated and kept frozen ( $-18\text{ }^{\circ}\text{C}$ ) prior to analysis.

### 2.2. Lipid extraction

Lipids were extracted by the Folch method (Folch, Lees, & Sloan Stanley, 1957). To 5 g of raw material, 33 ml of methanol were added and the whole mixed for at least 30 min at room temperature. Then, 66 ml of chloroform were added. The solution was stirred for 30 min. After eliminating the non-soluble part by filtration, 0.2 volumes of calcium chloride (0.9% in distilled water) were added. After decantation, in a separatory funnel (12 h at  $4\text{ }^{\circ}\text{C}$ ), the organic phase was recovered, and the solvent evaporated. Lipid fractions were stored in chloroform at  $-20\text{ }^{\circ}\text{C}$  for future use (Fig. 1).

### 2.3. Lipid class separation by column chromatography

A quantity of 85 mg of lipid were subjected to column chromatography, using a Chromabond SiOH column (Macherey-Nagel, Düren, Germany), as described by Dreyfus, Guerols, Freysz, and Hicks (1997). The column

was conditioned with 20 ml of chloroform. The neutral lipids, glyco-, and phospholipids were eluted by chloroform (20 ml), acetone/methanol (9:1, 30 ml) and methanol (20 ml), respectively. Solvents were evaporated and each class of lipids was estimated by weighing with a microbalance. All fractions were kept in chloroform at  $-20\text{ }^{\circ}\text{C}$  before use.

### 2.4. Saponification of lipid extract

The total lipid sample (10 mg) was refluxed with a 1 M solution of potassium hydroxide in 95% ethanol (2 ml) for 1 h (Christie, 1989c). The solution was cooled to room temperature, then 5 ml of water were added and the mixture was extracted thoroughly with hexane-diethyl ether (1:1, v/v;  $3 \times 5\text{ ml}$ ). The organic phase was washed with water and dried over anhydrous sodium sulphate, and the non-saponifiable materials were recovered by removing solvent in a rotary evaporator. The water washings were added to the aqueous layer, acidified by 6 M hydrochloric acid and further extracted with diethyl ether-hexane (1:1, v/v;  $3 \times 5\text{ ml}$ ). Free fatty acids (FFA) were recovered by washing the extract with water, drying it over anhydrous sodium sulphate and removing solvents by evaporation. FFA were kept in chloroform at  $-20\text{ }^{\circ}\text{C}$  before use.

### 2.5. Neutral lipid components identification by thin-layer chromatography

Analytical TLC was performed for the neutral lipids fraction on  $20 \times 20\text{ cm}$  chromatoplates coated with silica




Fig. 1. Flowsheet illustrating the different sample preparation steps.

gel G (0.25 mm thickness; Macherey-Nagel, Düren, Germany). Elutions were obtained by using hexane/ether/acetone (70:30:1, v/v/v) according to Mangold (Mangold, 1961, 1964).

Lipids were visualized by spraying successively with two specific reagents: first, with a primuline solution (5 mg in 100 ml of acetone/water, 80:20, v/v), which allows lipid spot identification under a UV lamp (Bergé, Gouyguo, Dubacq, & Durand, 1995); second, with a solution of sulphuric acid-vanillin, leading to colourized spots after heating slowly to 180 °C during 20 min.

All the lipids were identified by comparing their retention factors (Rf) with those of authentic standards (SIGMA, St Quentin Fallavier, France).

#### 2.6. Quantitative analysis of neutral lipid classes by thin-layer chromatography

Preparative TLC was performed on 20 × 20 cm chromatoplates (0.5 mm thickness; Macherey-Nagel, Düren, Germany) with 50 mg of neutral lipid. Compounds were separated by using the same eluting system as analytical TLC. Spots were scraped into glass tubes and eluted from the silica by two washes (2 × 10 ml) with two solvent mixtures: hexane/diethyl ether (50:50; v/v), and hexane/chloroform/methanol (10:10:10; v/v/v). Proportions of compounds were directly estimated in weight after solvent removal.

#### 2.7. Derivatisation of neutral lipid classes

##### 2.7.1. Free fatty acids and those linked to glycerol

A quantity of 500 mg of lipid were evaporated under nitrogen and transmethylated by contact with methanol/sulphuric acid (98:2) in excess for one night at 50 °C. After cooling, 1 ml of distilled water and 2 ml of hexane were added and vortexed. The upper organic phases, containing fatty acid methyl esters (FAMES), were collected and assayed by GC-FID and GC-MS. (Christie, 1993).

Picolinyl esters were prepared by treatment of FAMES with potassium *tert*-butoxide and 3-hydroxymethylpyridine during 45 min at 45 °C. Derivatives were extracted with 2 ml of hexane and 1 ml of water. The upper organic phase was dried with a sodium sulphate column. Derivatives were dried under a flush of nitrogen and dissolved in 1 ml of hexane for injection (Dubois, Barthelemy, & Bergé, 2006).

##### 2.7.2. Acetylation of sterols

A quantity of 2 ml of pyridine-acetic anhydride (2:1; v/v) were added to 500 µg of isolated sterols, at 45 °C for 45 min. After cooling to ambient temperature, 2 ml of distilled water were added. Steryl acetates were extracted with 4 ml of hexane. The upper organic phase was evaporated under nitrogen and 1 ml of hexane was added. The derivatives were further analysed by gas chromatography-mass spectrometry (GC-MS).

#### 2.8. Lipid identification by gas chromatography-mass spectrometry

Analyses of the lipid derivatives were performed by GC-MS on a Hewlett-Packard model 6890 series II gas chromatograph attached to an Agilent model 5973N selective quadrupole mass detector. GC-MS was connected to a computer with a Hewlett-Packard chemstation and the ionisation voltage used was 70 eV at 250 °C. The temperature of injector and interface were maintained at 250 °C and He was used as carrier gas under constant flow (1 ml/min). Separations were realised on a CP-Sil 5 CB low bleed MS (60 m × 0.25 mm i.d., 0.25 µm film thickness; Chrompack, Middleburg, The Netherlands).

FAMES, picolinyl esters, acetates and hydrocarbons were analysed by various methods:

- For FAMES, oven temperature was programmed from 80 to 170 °C at a rate of 30 °C/min, then from 170 to 295 °C at a rate of 3 °C/min.
- Analyses of picolinyl esters were operated with an oven temperature linearly increasing from 220 to 270 °C at 0.4 °C/min.
- Sterol acetates were analysed with linear programming. Oven temperature was programmed from 150 to 350 °C at a rate of 3 °C/min.
- For hydrocarbons (isolated by preparative TLC), oven temperature was programmed from 50 to 350 °C at a rate of 2 °C/min, then kept constant at 350 °C for 10 min.

#### 2.9. Quantification of fatty acids from neutral lipids by gas chromatography-flame ionisation detector

FAMES' quantification was done by GC-FID, using a gas chromatograph (Auto system Gas Chromatography, Perkin-Elmer) equipped with an auto-sampler and fitted with a split/splitless injector and flame ionisation detector. The separation was carried out with a BPX-70 capillary column (60 m long, 0.25 mm internal diameter, 0.25 µm film thickness; SGE, Austin, USA), containing a polar stationary phase (cyanopropyl-siloxane) (Bergé, Debiton, Dumay, Durand, & Barthelemy, 2002; Dumay, Barthelemy, & Bergé, 2004).

### 3. Results and discussion

It is well known that rays, like other elasmobranchs, store fat in liver (Kinsella, 1988). As expected, oil extraction from the liver of *H. bleekeri* confirmed that this organ is relatively rich in lipids (54% of the fresh matter). This is in accordance with previous studies on other elasmobranchs' livers, where, lipid contents ranged from 23 to 67% (Navarro-Garcia et al., 2000; Navarro-Garcia et al., 2004a; Navarro-garcia et al., 2004b; Ould El Kebir et al., 2003; Pal et al., 1998).

Table 1  
Distribution of neutral lipids from the liver of *Himantura bleekeri*

	Rf	% (w/w)
Hydrocarbons/steryl esters	0.94	6.3
Triacylglycerols	0.64	62.5
Free fatty acids/fatty alcohols	0.37	8.8
1,3-Diacylglycerols/sterols	0.21	8.3
1,2-Diacylglycerols	0.17	6.9
Glyceryl ethers/monoacylglycerols	0.02	7.3

Neutral lipid was the major lipid class (92%), while glycolipids and phospholipids were present in relatively low amounts (5% and 3%, respectively). Such distribution confirmed previous results obtained with liver oil of the ray *Dasyatis bleekeri* (Pal et al., 1998).

Composition of NL is presented in Table 1. Triacylglycerols (TAG) were found to represent about two-thirds of the NL. This predominance has been previously reported, irrespective of the species of rays: 67% in *Dasyatis brevis* and 67% in *Gymnura marmorata* (Navarro-garcia et al., 2004b), 75% in *Rhinoptera steindechneri* (Navarro-Garcia et al., 2004a), 93% in *D. bleekeri* (Pal et al., 1998) or sharks: *G. curvier* (37.5%), *C. falciformis* (56.9%) (Navarro-Garcia et al., 2000). However, high levels of NL particularly rich in TAG, are not specific to elasmobranchs but can be extended to all liver oils of marine organisms (Pigott, 1996), as TAG are known to be the main energy source (Navarro-garcia et al., 2004b).

As expected, intermediate compounds of the biosynthesis of TAG are also found in minor proportions, and they are free fatty acids (FFA), monoacylglycerol (MAG), 1,2 diacylglycerol (1,2-DAG) and 1,3 diacylglycerol (1,3-DAG). Sterols, hydrocarbons, steryl esters and glyceryl ethers were also found but, under our conditions, they were eluted with other components (see Table 1). A saponifica-

tion step is thus needed to quantify and identify them (see below).

Total fatty acids and those linked to glycerol and sterols are presented in Table 2. Up to 13 different FA (3 saturated, 4 mono-unsaturated and 6 PUFAs) have been identified by GC-MS using FAME and picolinyl derivatives, as previously described (Dubois et al., 2006). Their quantification was done by using GC/FID and FAME as previously mentioned (Bergé et al., 2002). Palmitic acid (16:0) was found to be predominant (28%), while the level of saturated fatty acids was found to be 45%. Thus, unsaturated fatty acids represented more than half of the total fatty acids (55%), as previously reported (Navarro-Garcia et al., 2000; Navarro-Garcia et al., 2004a; Navarro-garcia et al., 2004b; Pal et al., 1998). Among those unsaturated fatty acids, the *n*-3 PUFAs content was 27% with 16% of DHA and 4% of EPA, which is in accordance with the marine liver oil compositions already found (Bergé & Barnathan, 2005). Indeed, in cod liver oil, traditionally used as an omega-3 PUFAs source, the amount of EPA and DHA was around 16% (Méndez, Gonzalez, Inocente, Giudice, & Grompone, 1996). Our study confirms previous results (Bergé & Barnathan, 2005; Navarro-garcia et al., 2004b), indicating that ray oil can also be an alternative source for omega-3 (18% of EPA and DHA in *G. marmorata*, and 16% for *D. brevis*). In addition to these omega-3 compounds, omega-6 fatty acids were also found: arachidonic acid (20:4 *n*-6) represented 4% and the level of linoleic acid (18:2 *n*-6) was 1%. The resulting ratio of *n*-3 to *n*-6 fatty acids was 5.7 which is what seems to be needed for the prevention of chronic diseases (Bergé et al., 2005).

After a saponification step, total sterols represented 46% of the unsaponifiable fraction (Table 3). Cholesterol was the only one found free, while 12 were esterified with FA. Whatever the form (free or esterified), cholesterol was predominant and the other compounds were present in very

Table 2  
Fatty acid composition of liver oil of *Himantura bleekeri*

Fatty acid	Total FA (%)	MAG (%)	1,2-DAG (%)	1,3-DAG (%)	FFA (%)	TAG (%)	FFA linked to sterols (%)
C14:0	3.12	4.90	4.12	2.63	2.16	3.50	0.00
C16:0	28.3	36.9	30.6	27.5	24.6	32.3	29.3
C18:0	13.9	13.3	10.3	12.7	11.9	15.5	10.8
C16:1 w7	6.82	10.5	12.1	9.09	9.87	6.71	0.00
C18:1 w9	7.76	8.26	7.51	9.01	11.8	7.82	11.0
C18:1 w7	6.22	7.98	7.95	8.20	9.65	6.19	6.71
C20:1 w9	2.03	0.00	0.00	2.67	2.07	1.92	0.00
C18:2 w6	1.03	0.00	0.00	0.00	1.70	0.00	0.00
C20:4 w3	3.36	0.00	2.52	3.00	4.31	2.97	0.00
C20:4 w6	3.73	2.11	2.86	3.48	4.16	3.01	16.6
C20:5 w3	4.10	0.00	3.17	2.40	3.71	3.63	7.09
C22:5 w3	4.01	3.92	4.48	3.75	3.68	3.39	0.00
C22:6 w3	15.6	12.1	14.4	15.6	10.3	13.0	18.5
SFA	45.3	55.1	45.0	42.8	38.7	51.3	40.1
MUFA	22.8	26.8	27.6	29.0	33.4	22.6	17.7
PUFA	31.9	18.2	27.5	28.2	27.9	26.0	42.2
<i>n</i> -3 PUFA	27.1	16.1	24.6	24.7	22.0	23.0	25.6
<i>n</i> -6 PUFA	4.76	2.11	2.86	3.48	5.86	3.01	16.6

Table 3  
Composition of free and esterified sterols from the liver of *Himantura bleekeri*, analysed as acetates by gas chromatography

	Percentage area (%)
<i>Sterols (free)</i>	
Cholesterol	100
<i>Sterols (esterified)</i>	
22-Dehydrocholesterol	0.89
Cholesterol	97.8
Unidentified	Traces
Desmosterol	0.09
Latosterol	Traces
24-Methylene-cholesterol	0.45
Campesterol (R)	0.52
22-Dehydro-brassicasterol (S)	0.07
24-Methyl-desmosterol	Traces
Unidentified	0.02
Sitosterol	0.02
Fucosterol	0.08

low concentrations. Such results confirm a previous study on the liver oil of *D. bleekeri*, where cholesterol was found at a level of 68% while 22-dehydrocholesterol, campesterol and sitosterol were detected in low amounts (Pal et al., 1998). These observations suggest that cholesterol best performs the structural function of sterols in membranes (Nes, 1974).

Hydrocarbon composition of the liver oil of *H. bleekeri* is presented in Table 4. *n*-Alkanes were the major components (74%). Similar results have been previously published by Pal et al. (1998) for *D. bleekeri*. Up to 11 different hydrocarbons were identified. Squalene (22%) was at least twice more abundant than others; 9 alkanes were identified, ranging from 17 to 28 carbons. Squalene is recognized as an important intermediate in biosynthesis of cholesterol. It is used as a food additive and as a precursor of squalane; the latter compound is utilized largely as an excipient in cosmetics and pharmaceuticals (Bordier, Sellier, Foucault, & Le Goffic, 1996). Its potential clinical uses are various as it may have several roles as an antioxidant for human skin and eyes but also in detoxification of some xenobiotics (Kelly, 1999). This compound has been found in liver oil of many deep-sea shark species (Bordier et al., 1996;

Table 4  
Hydrocarbon compositions in ray liver oil

Hydrocarbon	Carbon number	Percentage (%)
Heptadecane	17	3.96
Octadecene	18	3.63
Heneicosane	21	4.42
Docosane	22	12.7
Tricosane	23	7.56
Tetracosane	24	13.8
Pentacosane	25	10.4
Hexacosane	26	9.92
Heptacosane	27	5.76
Octacosane	28	5.75
Squalene	30	22.1

Wetherbee & Nichols, 2000) but also in the ray, *D. bleekeri* (Pal et al., 1998).

At least, as expected, glyceryl ethers were also detected. Such compounds have been previously found in ray liver and shark liver (Bordier et al., 1996; Hayashi & Takagi, 1981; Navarro-Garcia et al., 2000; Pal et al., 1998; Wetherbee & Nichols, 2000). They are of interest since many benefits for human health, e.g., bacteriostatic and fungistatic properties, anti-inflammatory activities and hematopoietic effects, were attributed to those compounds (Bordier et al., 1996).

#### 4. Conclusions

In conclusion, this study demonstrates that the liver of the ray, *H. bleekeri*, which is actually wasted, could be used as new raw material for oil production. Indeed, such oil, which could be obtained in relatively high amounts, is an excellent source of omega-3 PUFAs, particularly DHA. Also, the presence of squalene and glyceryl ethers can justify such exploitation for nutritional or cosmetic purposes.

#### Acknowledgements

The authors thank SIBIO Biosoft technologies (Romainville, France) for their help and advice.

#### References

- Bergé, J. P., & Barnathan, G. (2005). Fatty acids from lipids of marine organisms: molecular biodiversity, roles as biomarkers, biologically active compounds, and economical aspects. *Advances in Biochemical and Engineering/Biotechnology*, 96, 49–125.
- Bergé, J. P., Gouygou, J. P., Dubacq, J. P., & Durand, P. (1995). Reassessment of lipid composition of the diatom *Skeletonema costatum*. *Phytochemistry*, 39, 1017–1021.
- Bergé, J. P., Debiton, E., Dumay, J., Durand, P., & Barthomeuf, C. (2002). In vitro anti-inflammatory and anti-proliferative activity of sulfolipids from the red alga *Porphyridium cruentum*. *Journal of Agricultural and Food Chemistry*, 50, 6227–6232.
- Blyth, E. (1860). The cartilaginous fishes of lower Bengal. *Journal of the Asiatic Society of Bengal*, 35–45.
- Bordier, C. G., Sellier, N., Foucault, A. P., & Le Goffic, F. (1996). Purification and characterization of deep sea shark *Centropristis squamosus* liver oil 1-*O*-alkylglycerol ether lipids. *Lipids*, 31, 521–528.
- Christie, W. W. (1989c). Gas chromatography and lipids. *Gas Chromatography–Mass Spectrometry and Fatty Acids*, 161–183.
- Christie, W. W. (1993). Preparation of ester derivatives of fatty acids for chromatographic analysis. *Advances in lipid methodology* (2nd ed.). (pp. 69–112).
- Connor, W. E. (2000). Importance of *n*-3 fatty acids in health and disease. *American Journal of Clinical Nutrition*, 71, 171–175.
- Dreyfus, H., Guerols, B., Freysz, L., & Hicks, D. (1997). Successive isolation and separation of the major lipid fractions including gangliosides from single biological samples. *Analytical Biochemistry*, 249, 67–78.
- Dubois, N., Barthomeuf, C., & Bergé, J. P. (2006). Convenient preparation of picolinyl derivatives from fatty acid esters. *European Journal of Lipid Science and Technology*, 108, 28–32.
- Dumay, J., Barthomeuf, C., & Bergé, J. P. (2004). How enzymes may be helpful for upgrading fish by-products: enhancement of fat extraction. *Journal of Aquatic Food Product Technology*, 13, 69–84.

- Folch, J., Lees, M., & Sloan Stanley, G. H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*, 226, 497–509.
- Hayashi, K., & Takagi, T. (1981). Distribution of squalene and diacyl glyceryl ethers in the different tissues of deep-sea shark, *Dalatias licha*. *Bulletin of the Japanese Society of Scientific Fisheries*, 47, 281–288.
- Kelly, N. D. G. S. (1999). Squalene and its potential clinical uses. *Alternative Medicine Review*, 4, 29–36.
- Kinsella, J. E. (1988). Foods lipids and fatty acids: importance in food quality, nutrition and health. *Food Technology*, 124–145.
- Mangold, H. K. (1961). Thin layer chromatography of lipids. *Journal of the American Oil Chemists Society*, 38, 708–727.
- Mangold, H. K. (1964). Thin layer chromatography of lipids. *Journal of American Oil Chemists Society*, 47, 762–773.
- Méndez, E., Gonzalez, R. M., Inocente, G., Giudice, H., & Grompone, M. A. (1996). Lipid content and fatty acid composition of fillets of six fishes from the Rio de la Plata. *Journal of Food Composition and Analysis*, 9, 163–170.
- Mnari, A., Bouhlef, I., Chraief, I., Hammami, M., Romdhane, M. S., El Cafsi, M., et al. (2007). Fatty acids in muscles and liver of tunisian wild and farmed gilthead sea bream, *Sparus aurata*. *Food Chemistry*, 100, 1393–1397.
- Navarro-Garcia, G., Pacheco-Aguilar, R., Vallejo-Cordova, B., Ramirez-Suarez, J. C., & Bolanos, A. (2000). Lipid composition of the liver oil of shark species from the caribbean and gulf of california waters. *Journal of Food Composition and Analysis*, 13, 791–798.
- Navarro-Garcia, G., Bringas-Alvarado, L., Pacheco-Aguilar, R., & Ortega-Garcia, J. (2004a). Oxidative resistance, carotenes, tocopherols and lipid profile of liver oil of the ray *Rhinoptera steinchneri*. *Journal of Food Composition and Analysis*, 17, 699–706.
- Navarro-garcia, G., Pacheco-Aguilar, R., Bringas-Alvarado, L., & Ortega-Garcia, J. (2004b). Characterization of the lipid composition and natural antioxidants in the liver oil of *Dasyatis brevis* and *Gymnura marmorata* rays. *Food Chemistry*, 87, 89–96.
- Nes, W. R. (1974). Role of sterols in membranes. *Lipids*, 9, 596–612.
- Ould El Kebir, M. V., Barnathan, G., Siau, Y., Miralles, J., & Gaydou, E. (2003). Fatty acids distribution in muscle, liver, and gonads of rays (*Dasyatis marmorata*, *Rhinobatos cemiculus*, and *Rhinoptera marginata*) from the east tropical atlantic ocean. *Journal of Agricultural and Food Chemistry*, 51, 1942–1947.
- Pal, D., Banerjee, D., Patra, T. K., Patra, A., & Ghosh, A. (1998). Liver lipids and fatty acids of the sting ray *Dasyatis bleekeri* (Blyth). *Journal of the American Oil Chemists Society*, 75, 1373–1378.
- Pigott, M. G. (1996). Marine oils (5th ed.). In Y. H. Hui (Ed.). *Bailey's Industrial Oil and Fat Products* (Vol. 3, pp. 225–254). New York: Wiley J. & Sons, Inc.
- Wetherbee, B. M., & Nichols, P. D. (2000). Lipid composition of the liver oil of deep-sea sharks from the Chatham Rise, New Zealand. *Comparative Biochemistry and Physiology Part B*, 125, 511–521.