

Fumonisin level in corn-based food and feed from Linxian County, a high-risk area for esophageal cancer in china

Jiansheng Wang^a, Ying Zhou^a, Wanbo Liu^b, Xiangcheng Zhu^b,
Liangcheng Du^b, Qiaomei Wang^{a,*}

^a Department of Horticulture, Zhejiang University, The State Agriculture Ministry Laboratory of Horticultural, Plant Growth Development and Biotechnology, Hangzhou 310029, China

^b Department of Chemistry, University of Nebraska-Lincoln, Lincoln, NE 68588-0304, USA

Received 21 January 2007; received in revised form 28 March 2007; accepted 30 May 2007

Abstract

The level of mycotoxin fumonisins in corn-based food and feed collected from Linxian County, a high-risk area for esophageal cancer in China, has been analyzed using high-performance liquid chromatographic coupled with evaporative laser scattering detector (HPLC-ELSD). A total of 104 corn kernel samples were obtained from local households, granaries, wholesale markets (central markets), and retail markets (stores and supermarkets). Fumonisin B₁ (FB₁) was detected in the samples from households, granaries, central markets, and stores, with a positive rate of 61.5%, 50%, 33.3%, and 17%, respectively. No fumonisin was detected in samples from the supermarket. The highest FB₁ levels (0.30–3.20 µg/g; mean, 1.42 µg/g) were found in samples from the granary, followed by household (0.25–1.80 µg/g; mean, 0.73 µg/g), central market (0.25–1.10 µg/g; mean, 0.51 µg/g), and store (0.22–0.34 µg/g; mean, 0.28 µg/g). Among the 80 corn kernel samples collected from local households, 18 of 24 (75.0%) moldy samples contained high levels of FB₁ (0.28–3.30 µg/g; mean, 1.58 µg/g), and 20 of 56 (35.7%) apparently healthy samples contained low levels of FB₁ (0.21–0.82 µg/g; mean, 0.46 µg/g). As the central market plays an important role in trade of corn-based food and feed in China, a total of 115 corn-based food and feed samples were collected from the local central market. The highest FB₁ levels (0.30–3.13 µg/g; mean, 1.50 µg/g) were found in feed, followed by unprocessed food (0.31–0.63 µg/g; mean, 0.47 µg/g) and processed food (0.21–0.28 µg/g; mean, 0.25 µg/g). The positive incidence of FB₁ in feed, unprocessed, and processed food were 53.6%, 33.3% and 17.9%, respectively. In conclusion, the results showed that corn-based food and feed from Linxian County contained low level of FB₁ (<2 µg/g) in general, but efforts should be made to control the fumonisin contamination in corn kernels stored in granaries and households.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Fumonisin; HPLC-ELSD; Corn products

1. Introduction

Fumonisin are a class of mycotoxins produced mainly by *Fusarium moniliforme* (synonym *F. verticillioides*), a primary fungal contaminant of corn and corn-derived products throughout the world. So far, twenty eight fumonisin analogs have been characterized (Rheeder, Marasas, & Vismer, 2002). They are classified into four main

groups, the A, B, C, and P-series fumonisins. The B-series fumonisins are the most abundant analogs produced by *F. moniliforme* strains, with fumonisin B₁ (FB₁) being the most abundant and toxic constituent. FB₁ accounts for approximately 70% of total fumonisins found in nature (Marasas, 2001; Nelson, Desjardins, & Plattner, 1993).

Elevated fumonisin level in corn has become an area of concern since ingestion of fumonisin-contaminated corn has been associated with livestock loss and human health risks. Fumonisin were reported to cause leukoencephalomalacia (ELEM) in horses (Marasas, 2001), pulmonary oedema syndrome (PES) in pigs (Harrison, Colvin, Greene,

* Corresponding author. Tel.: +86 571 85909333; fax: +86 571 87420554.

E-mail address: qmwang@zju.edu.cn (Q. Wang).

Newman, & Cole, 1990), and hepatocarcinoma in rats (Gelderblom, Abel, & Smuts, 2001). In addition to the diseases in livestock, consumption of corn-based food contaminated with fumonisins is known to epidemiologically relate to human esophageal cancer (HEC) in people from the Transkei region of South Africa and the Linxian region of China (Chu & Li, 1994; Rheeder et al., 1992; Sydenham et al., 1990). It was also reported that gestational exposure to fumonisins may result in birth defects, such as neural tube defects (Hendricks, 1999). Several modes of action have been postulated to explain fumonisin-induced toxicity, but the primary hypothesis involves disruption of de novo sphingolipid biosynthesis (Merrill, Sullards, Voss, & Riley, 2001). Fumonisins are specific inhibitors of sphinganine *N*-acyltransferase (ceramide synthase), a key enzyme in the biosynthesis of sphingolipids.

The major high-risk areas for HEC in China are located in Henan and Hebei Provinces, with Linxian County in the Henan Province being the highest-risk area. Several surveys have been conducted concerning the natural occurrence of fumonisin in corn samples collected from households in Linxian County. Chu and Li (1994) detected high level of FB₁ (18–155 µg/g; mean, 74 µg/g) in all moldy samples from households in Linxian County, and relatively lower level of FB₁ (20–60 µg/g; mean, 35.3 µg/g) in all normal samples from the same households. However, lower incidence and level of FB₁ were also reported in other surveys conducted in Linxian County by Yoshizawa and his colleagues (1994) and Wang and Zhu (2002). The content of FB₁ varied greatly among different surveys conducted in Linxian County, which may be due to the different methods used for fumonisin analysis. As reported recently (Shepherd, Van der Westhuizen, Gatyeni, Katerere, & Marasas, 2005), fumonisins detected in food using indirect analytical methods could sometimes produce false positive results, and it is important to have more definitive techniques to validate the results, because Linxian County is a main corn-producing and HEC-high-risk area in China.

Moreover, corns produced by farmers in Linxian County are not only stored and consumed by themselves in household, but also stored in granary or traded in local wholesale and retail markets. The trade of corn-based foods and feeds has been increasing in the local markets. However, limited information is available on fumonisin contamination in corn-based foods and feeds traded in Linxian County. It is important to investigate fumonisin contamination in various sources. The objective of this work was to use HPLC-ELSD to directly identify and measure fumonisins in corn and corn-based food and feed from different sources in Linxian County.

2. Materials and methods

2.1. Chemicals

High-performance liquid chromatography (HPLC)-grade acetonitrile was from EM Science (Darmstadt, Germany)

or from Fisher Scientific (Pittsburgh, PA). Standard FB₁, FB₂, FB₃, and FB₄ were gifts from Ronald D. Plattner (US Department of Agriculture [USDA], Peoria, Ill.).

2.2. Sample collection

Corn kernel samples were collected from local farmers (household), granaries, central markets, stores, and supermarkets, while unprocessed corn kernels, processed corn foods, and corn-based feed were taken from local central market in Linxian County, Henan Province, in the period from July 2005 to May 2006. The processed corn foods included corn grit, corn flour, corn meal, corn tortilla chips and vacuum-packaging cooked corn. The samples were sent to the laboratory as soon as they were collected, and tested upon arrival or stored at –20 °C to arrest any fumonisin formation up to the time of analysis.

2.3. Determination of fumonisin

Analysis of fumonisins was performed according to our former method (Bojja, Cerny, Proctor, & Du, 2004), with minor modifications.

2.3.1. Extraction and clean up

A 10 g aliquot of the sample was placed in a flask containing 25 ml acetonitrile/water (1:1, v/v). Samples were placed in an orbital shaker overnight, filtered and a 10-ml aliquot was transferred to a 50 ml centrifugal tube and added 40 ml deionized water. Clean up of the extract was performed in 300 mg Amberlite XAD-4, previously conditioned with 2 ml of methanol and 20 ml of deionized water. After the initial washing with 200 ml deionized water, the extract was eluted with 3 ml 100% methanol. The eluent was dried under vacuum with freezing at –65 °C and stored at –20 °C, up to the moment of quantification.

2.3.2. Chromatographic quantification of fumonisins by HPLC-ELSD analysis

Samples were dissolved in 200 µl water and filtered through a 0.2 µm syringe-filter. A 20 µl aliquot was injected directly into the HPLC-ELSD system and used for quantification of the fumonisins. The HPLC system was a Pro-Star, Model 210 (Varian Walnut Creek, CA) with a column of Alltima C18LL, 5 µm, 250 × 4.6 mm inner diameter (Alltech, Deerfield, Ill.). The HPLC-ELSD method conditions were performed according to our previous studies with some modifications (Bojja et al., 2004). The mobile phases were (A) water-TFA (100:0.025, v/v) and (B) acetonitrile-TFA 100:0.025, (v/v), with a gradient of 0–20% B in A in the first 5 min, 20–40% B from 5 to 10 min, 40–80% B from 10 to 15 min, 80% B from 15 to 20 min, and 80–0% B from 20 to 25 min. The flow rate was 1.0 ml/min. The conditions set for ELSD (ELSD 2000, Alltech, USA) were 45 °C of drift tube temperature, 2.0-l/min nitrogen gas flow, and gain value of 1 in the impactor-on mode.

In quantitative analyses, a volume of 20 μl standard FB_2 , FB_3 and FB_4 samples (1 $\mu\text{g}/\mu\text{l}$) was injected for HPLC-ELSD analysis, and standard FB_1 (10 $\mu\text{g}/\mu\text{l}$) was diluted in four different concentrations of 0.15, 0.3, 0.6, 1.2 $\mu\text{g}/\mu\text{l}$ with the same method above. The peak area from the responding peak was integrated using on-system tools provided by Varian. At least 2 injections were made for each concentration, and the peak areas were then plotted against the absolute amounts of FB_1 used to obtain a standard curve. Under the established conditions, standard FB_1 gave a peak at a retention time of 16.26 min. Retention times for standard FB_3 , FB_2 and FB_4 were 16.9, 17.2 and 17.9 min, respectively.

The method gave a linear response in the range of 3–24 μg FB_1 with $R^2 = 0.9828$. In five independent spiking experiments, where a known amount of standard FB_1 was added to the food sample, the recovery rate was also determined. The rates varied from 88.6% to 112.7%, with an average of 102.6% and a standard deviation of 8.9% ($n = 5$). The limit of determination was 3 $\text{ng}/\mu\text{l}$ for FBs.

3. Results and discussion

Linxian County is one of the main maize-producing areas in Henan Province. Corn produced in Linxian County is either stored by local farmers and in granaries, or traded in local markets. The newly harvested corn cobs without husk, are usually dried under the sun on the ground floor in order to reduce the moisture content; then the sufficiently dried cobs are shelled, and the corn kernels are either stored or traded. Local farmers usually store the corn kernels in gunnysacks or put them in cement tanks inside the house without any ventilated equipment, for household storage. As for granary storage, no ventilated equipment is used either. The corn kernels are stored in cement tanks after wrapped with plastic film, and the top of cement tanks are then covered with straw mat to avoid insect and rodent attack. The newly harvested corn kernels with good visual quality and low moisture content, are usually traded in the markets. In all the corn samples analyzed, FB_1 was the only fumonisin detected, and no detectable levels of FB_2 , FB_3 and FB_4 were found. In previous studies, the co-occurrence of FB_1 and FB_2 in corn (Yoshizawa et al., 1994), corn-based food (Bittencourt, Oliveira, Dilkin, & Corrêa, 2005), and feed (Sanchis et al., 1995), with FB_1 being the predominant fumonisin have been reported. In our test, only FB_1 was detectable in corn samples, which agrees with some other surveys in Linxian County (Chu & Li, 1994; Wang, Wei, Ma, & Luo, 2000; Wang & Zhu, 2002).

We collected a total of 104 corn kernel samples from local households, granaries, central markets, stores and supermarkets for fumonisin analysis. The highest incidence of FB_1 contamination (61.5%) was found in samples from households, followed by from granaries (50%), central markets (33.3%), and stores (17.0%). No FB_1 was detected

in samples from supermarkets. The highest FB_1 level (0.30–3.20 $\mu\text{g}/\text{g}$; mean, 1.42 $\mu\text{g}/\text{g}$) was found in samples from granaries, followed by households (0.25–1.80 $\mu\text{g}/\text{g}$; mean, 0.73 $\mu\text{g}/\text{g}$), central markets (0.25–1.10 $\mu\text{g}/\text{g}$; mean, 0.51 $\mu\text{g}/\text{g}$), and stores (0.22–0.34 $\mu\text{g}/\text{g}$; mean, 0.28 $\mu\text{g}/\text{g}$) (Table 1).

No moldy corn kernel samples were found in the central and retail markets during our sampling, and all the samples collected were apparently healthy corn without any visible mold contamination, which may explain the low incidence and levels of FB_1 in corn samples traded in the markets in the present survey. As most of the moldy samples were from households, we conducted another survey, randomly collecting 80 samples from household and dividing them into two groups, moldy and normal healthy, according to their surface character. FB_1 analysis of the samples showed that high levels of FB_1 (from 0.28 to 3.30 $\mu\text{g}/\text{g}$; mean, 1.58 $\mu\text{g}/\text{g}$) were found in 18 of 24 (75%) moldy samples, while low levels (from 0.21 to 0.82 $\mu\text{g}/\text{g}$; mean, 0.46 $\mu\text{g}/\text{g}$) were found in 20 of 56 (35.7%) apparently healthy samples (Table 2), which suggested that even corn samples with normal surface character might contain fumonisin.

Our results indicated that fumonisin levels of all samples were significantly lower compared to the maximum level of FDA listed (2–4 ppm for human food) (FDA, 2002). Specifically, the samples from supermarket were free of FB_1 contamination, although FB_1 contamination was slightly more serious in corn kernel samples from local households and granaries than those from local markets. Several previous surveys have been conducted concerning the natural occurrence of fumonisin in corn

Table 1
Fumonisin contamination of corn from different sources

Source	Samples		Level of FB_1 in positive samples	
	Number	Positive (incidence)	Mean ($\mu\text{g}/\text{g}$)	Range ($\mu\text{g}/\text{g}$)
Household	26	16 (61.5%)	0.73 \pm 0.18 ¹	0.25–1.80
Granary	24	12 (50.0%)	1.42 \pm 1.51	0.30–3.20
Central market	36	12 (33.3%)	0.51 \pm 0.28	0.25–1.10
Store	12	2 (17.0%)	0.28 \pm 0.08	0.22–0.34
Supermarket	6	0	– ²	–

¹ Mean \pm standard derivation.

² Below detection limit.

Table 2
Fumonisin contamination of corn samples from household

Surface character	Samples		Level of FB_1 in positive samples	
	Number	Positive (incidence)	Mean ($\mu\text{g}/\text{g}$)	Range ($\mu\text{g}/\text{g}$)
Moldy samples	24	18 (75.0%)	1.58 \pm 1.21	0.28–3.30
Normal samples	56	20 (35.7%)	0.46 \pm 0.20	0.21–0.82

samples collected from households in Linxian County. Chu and Li (1994) detected high level of FB₁ in concentrations ranging from 18 to 155 µg/g (mean 74 µg/g) in all moldy samples from households in Linxian County, and relatively lower level of FB₁ in concentrations ranging from 20 to 60 µg/g (mean 35.3 µg/g) was detected in all normal samples from the same households. However, lower incidence and level of FB₁ were also reported in another survey conducted in Linxian County by Yoshizawa and his colleagues, who described FB₁ contamination frequencies of 48% with average FB₁ level of 0.872 µg/g (Yoshizawa et al., 1994). A recent report by Wang and Zhu (2002) showed that FB₁ at the concentrations ranging from 1.07 to 2.56 µg/g was detected in 50% of moldy corn samples collected from households of Jingtou village in Linxian County, while low level (0.21–0.737 µg/g) of FB₁ was detected in 10.5% of 19 normal samples from the same households. Both the incidence and level of FB₁ in either moldy or healthy corn samples from household in Linxian County in the current survey were much lower than those reported by Chu and Li (1994), but similar to those reported by Yoshizawa and his colleagues (1994), and Wang and Zhu (2002).

The variation of FB₁ content among different surveys in corn from households in Linxian County may be related to the climate conditions during and after harvest. For example, harvest occurring during a rainy period favours for fungal growth and fumonisin production (Jackson & Jalonski, 2004). Our results indicate, that although some helpful means have been adopted to control the fumonisin contamination of corn from household in Lianxian County in recent years, there are still many areas requiring improvements. For example, moldy corn is still harvested, stored, and consumed by the rural population in Linxian County, which represents a health risk for the people.

A direct detection method was used to determine the level of fumonisin contaminations in corn from granaries and markets in Linxian County. The results suggested that both the incidence and level of FB₁ in corn samples from granaries were higher than those of markets, with FB₁ level as high as 3.20 µg/g being determined. Although fumonisin formation is believed to occur predominately in corn before harvest, it was also reported that the toxins could be formed during post-harvest storage, especially when corn was inadequately stored at high temperature and high relative moisture (Jackson & Jalonski, 2004). The relatively heavy FB₁ contamination in corn from granary might be related to the storage environment and method. It is easy to create partial high moisture in corn stored in such a non-ventilated storage system as granary, which is suitable for fungal (*Fusarium*) growth and mycotoxin production. A simple and effective method to reduce fumonisin contamination is to keep corn kernels sufficiently dried (moisture content less than 13%) before storage to ensure unfavourable conditions for fungal growth. To do this, the corn cob without husk can either be dried on the floor during sunny days, or put them on an indoor platform during

rainy period before storing them in households or granaries.

Corn kernels are processed to varieties of corn-based foods and feed besides being directly consumed as food. We conducted another survey of fumonisins contamination in corn-based food and feed samples from central markets of Lianxian County. The results showed that FB₁ was detected in the ranges from 0.21 to 0.28 µg/g (mean, 0.25 ± 0.04 µg/g) in 7 of 39 (17.9%) processed corn food samples. No fumonisin was detected in all corn flour, corn meal and corn tortilla chip samples, only one of six vacuum-packaging cooked corn cob samples detected positive FB₁ at the level of 0.28 µg/g. However, FB₁ was detected at the concentration ranging from 0.21 to 0.27 µg/g in 6 of 12 (50%) corn grit samples (Table 3). Our results showed that less or no contamination of fumonisin was found in the processed corn products, which is coincident with previous reports (Weidenbörner, 2001; Humpf & Voss, 2004). Several previous studies were conducted to investigate the effect of processing method on fumonisin content. It was reported that the use of alkaline solution, water and temperature, used in the prepared corn-based foods might contribute to variations in the occurrence of fumonisin (Soriano & Dragacci, 2004). Cleaning corn to remove damaged or moldy kernels reduces fumonisins in foods while milling increases their concentration in some and reduces their concentration in other products (Jackson & Jalonski, 2004). Humpf and Voss (2004) found that baking, frying, and extrusion cooking of corn at high temperatures (≥ 190 °C) also reduced fumonisin concentrations in foods, with the amount of reduction achieved depending on cooking time, temperature, recipe, and other factors. It was also suggested that the fumonisin level in corn-based products was reduced during processing because fumonisins become 'hidden' and not recoverable under currently used extraction/purification procedures (Kim, Scott, & Lau, 2003). There are only a few reports on fumonisin contamination in corn-based products in China, while relatively more surveys are conducted in the occurrence of fumonisins in corn-based products in some other countries (Soriano &

Table 3
Fumonisin contamination of corn-based foods and feeds from central market

Use purpose	Samples		Level of FB ₁ in positive samples	
	Number	Positive (incidence)	Mean (µg/g)	Range (µg/g)
Processed food	39	7 (17.9%)	0.25 ± 0.04	0.21–0.28
Corn grit	12	6 (50%)	0.23 ± 0.03	0.21–0.27
Corn flour	6	0	–	–
Corn meal	9	0	–	–
Corn tortilla chips	6	0	–	–
Vacuum-packaging cooked corn cob	6	1	0.28	0.28
Unprocessed food	48	16 (33.3%)	0.47 ± 0.12	0.31–0.63
Feed	28	15 (53.6%)	1.50 ± 1.24	0.30–3.13

Dragacci, 2004; Weidenbörner, 2001). Moreover, the consumption of corn is different from other countries in China as some special processing methods are used. It will be interesting to further investigate the effect and mechanism of different processing methods on fumonisin level of corn-based foods.

High levels of FB₁ ($1.50 \pm 1.24 \mu\text{g/g}$) were found in 15 of 28 (53.6%) feed samples with the maximal level of $3.13 \mu\text{g/g}$, which suggests that the feed produced in Linxian County need some more suitable practice for fumonisin control. In Linxian County, the corn kernels with good visual quality and low moisture content are directly traded in the market or used to made corn-based foods, while those with relatively poor quality are stored and consumed by the farmers or used to produce feed. Our results indicated that the fumonisin contamination is more serious in corn feed than in corn-based foods obtained from central markets. The Food and Drug Administration of USA has announced guidance levels for total fumonisin levels in corn products, 2–4 ppm for human foods and 5–100 ppm for animal feeds depending on the species and the proportion of the contaminated material in the total diets (FDA, 2002). Therefore, the feed traded in central markets of Linxian County is still safe according to this guidance standard.

4. Conclusion

FB₁ was the only fumonisin detected by HPLC-ELSD in all corn-based food and feed samples collected in Linxian County, a high-risk area for esophageal cancer in China. No fumonisin was detected in corn samples from local supermarket, and processed corn foods (corn flour, corn meal, and corn tortilla chips) from local central markets. Low average levels of FB₁ ($<2 \mu\text{g/g}$) were found in all corn samples collected locally, and all corn-based food and feed samples from local central markets, although relatively high levels of FB₁ were detected in corn samples from local granaries, and moldy corn samples from local households with maximal FB₁ level of 3.20 and $3.30 \mu\text{g/g}$, respectively. The results indicate that new practices should be adopted to control the fumonisin contamination of corn kernels stored in granaries and households.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (30428023, 30320974), Fok Ying Tong Education Foundation (104034), NCET-05-0516, and Natural Science Foundation of Zhejiang Province (R304103).

References

Bittencourt, A. B. F., Oliveira, C. A. F., Dilkin, P., & Corrêa, B. (2005). Mycotoxin occurrence in corn meal and flour traded in São Paulo, Brazil. *Food Control*, 16, 117–120.

- Bojja, R. S., Cerny, R. L., Proctor, R. H., & Du, L. C. (2004). Determining the biosynthetic sequence in the early steps of the fumonisin pathway by use of three gene-disruption mutants of *Fusarium verticillioides*. *Journal of Agricultural and Food Chemistry*, 52, 2855–2860.
- Chu, F. S., & Li, G. Y. (1994). Simultaneous occurrence of fumonisin B₁ and other mycotoxins in moldy corn collected from the People's Republic of China in regions with high incidences of esophageal cancer. *Applied and Environmental Microbiology*, 60, 847–852.
- FDA. (2002). Draft guidance for industry: fumonisin levels in human foods and animal feeds; availability. Fed.Regist, 65, 35945. (Available from <<http://www.cfsan.fda.gov/~dms/fumongui.html>>).
- Gelderblom, W. C., Abel, S., & Smuts, C. M. (2001). Fumonisin-induced hepatocarcinogenesis: mechanisms related to cancer initiation and promotion. *Environmental Health Perspectives*, 109(Suppl 2), 291–300.
- Harrison, L. R., Colvin, B. M., Greene, J. T., Newman, L. E., & Cole, J. R. (1990). Pulmonary oedema and hydrothorax in swine produced by fumonisin B₁, a toxic metabolite of *Fusarium moniliforme*. *Journal of Veterinary Diagnostic Investigation*, 2, 217–221.
- Hendricks, K. (1999). Fumonisin and neural tube defects in South Texas. *Epidemiology*, 10, 198–200.
- Humpf, H. U., & Voss, K. A. (2004). Effects of thermal food processing on the chemical structure and toxicity of fumonisin mycotoxins. *Molecular Nutrition and Food Research*, 48, 225–269.
- Jackson, L., & Jalonski, J. (2004). Fumonisin. In O. M. Magan (Ed.), *Mycotoxins in food* (pp. 380–385). Boca Raton, Boston, New York, Washington, DC, USA: CRC Press.
- Kim, E. K., Scott, P. M., & Lau, B. P. Y. (2003). Hidden fumonisin in corn flakes. *Food Additives and Contaminants*, 20(2), 161–169.
- Marasas, W. F. O. (2001). Discovery and occurrence of the fumonisins: A historical perspective. *Environmental Health Perspectives*, 109, 239–243.
- Merrill, A. H., Jr., Sullards, M. C., Voss, K. A., & Riley, R. T. (2001). Sphingolipid metabolism: roles in signal transduction and disruption by fumonisins. *Environmental Health Perspectives*, 109(Suppl 2), 283–289.
- Nelson, P. E., Desjardins, A. E., & Plattner, R. D. (1993). Fumonisin, mycotoxins produced by *Fusarium* species: Biology, chemistry, and significance. *Annual Review of Phytopathology*, 31, 233–252.
- Rheeder, J. P., Marasas, W. F. O., Thiel, P. G., Sydenham, E. W., Shephard, G. S., & Van Schalkwyk, D. J. (1992). *Fusarium moniliforme* and fumonisins in corn in relation to human oesophageal cancer in Transkei. *Phytopathology*, 82, 353–357.
- Rheeder, J. P., Marasas, W. F., & Vismer, H. F. (2002). Production of fumonisin analogs by *Fusarium* species. *Applied and Environmental Microbiology*, 68, 2101–2105.
- Sanchis, V., Abadias, M., Oncins, L., Sala, N., Vinas, I., & Canela, R. (1995). Fumonisin B₁ and B₂ and toxigenic *Fusarium* strains in feeds from the Spanish market. *International Journal of Food Microbiology*, 27, 37–44.
- Shephard, G. S., Van der Westhuizen, L., Gatyeni, P. M., Katerere, D. R., & Marasas, W. F. O. (2005). Do fumonisin mycotoxins occur in wheat? *Journal of Agricultural and Food Chemistry*, 53, 9293–9296.
- Soriano, J. M., & Dragacci, S. (2004). Intake, decontamination and legislation of fumonisins in foods. *Food Research International*, 37, 367–374.
- Sydenham, E. W., Thiel, P. G., Marasas, W. F. O., Shephard, G. S., Van Schalkwyk, D. J., & Koch, K. R. (1990). Natural occurrence of some *Fusarium* mycotoxins in corn from low and high esophageal cancer prevalence areas of the Transkei, Southern Africa. *Journal of Agricultural and Food Chemistry*, 38, 1900–1903.
- Wang, H., Wei, H., Ma, J., & Luo, X. (2000). The fumonisin B₁ content in corn from North China, a high-risk area of oesophageal cancer.

- Journal of Environmental Pathology Toxicology and Oncology*, 19, 139–141.
- Wang, Y., & Zhu, T. X. (2002). Determination of FB₁ in corn with high incidence of esophageal cancer in Linxian Jingtoucun. *Journal of China Agricultural University*, 7(1), 9–13 (in Chinese).
- Weidenbörner, M. (2001). Foods and fumonisins. *European Food Research and Technology*, 212, 262–273.
- Yoshizawa, T., Yamashita, A., & Luo, Y. (1994). Fumonisin occurrence in corn from high- and low-risk areas for human esophageal cancer in China. *Applied and Environmental Microbiology*, 60, 1626–1629.