

Cytotoxic chalcones and antioxidants from the fruits of *Syzygium samarangense* (Wax Jambu)

Mario J. Simirgiotis^a, Seiji Adachi^b, Satoshi To^b, Hui Yang^a, Kurt A. Reynertson^c,
Margaret J. Basile^d, Roberto R. Gil^e, I. Bernard Weinstein^b, Edward J. Kennelly^{a,*}

^a Department of Biological Sciences, Lehman College and The Graduate Center, City University of New York,
250 Bedford Park Boulevard West, Bronx, NY 10468, United States

^b Herbert Irving Comprehensive Cancer Center, College of Physicians and Surgeons of Columbia University,
701 West 168th Street, New York, NY 10032, United States

^c Weill Medical College of Cornell University, Department of Pharmacology and the Center for Complementary and Integrative Medicine,
1300 York Avenue, New York, NY 10021, United States

^d Department of Neurology, University of Miami Miller School of Medicine, 1501 NW 9th Avenue, Miami, FL 33136, United States

^e Department of Chemistry, Carnegie Mellon University, 4400 Fifth Avenue, Pittsburgh, PA 15213, United States

Received 22 June 2007; received in revised form 8 August 2007; accepted 31 August 2007

Abstract

Bioassay-guided fractionation of the methanolic extracts of the pulp and seeds of the fruits of *Syzygium samarangense* (Blume) Merr. and L.M. Perry led to the identification of four cytotoxic compounds and eight antioxidants on the basis of HPLC-PDA analysis, MS, and various NMR spectroscopic techniques. Three *C*-methylated chalcones, 2',4'-dihydroxy-3',5'-dimethyl-6'-methoxychalcone (**1**), 2',4'-dihydroxy-3'-methyl-6'-methoxychalcone (stercurensin, **2**), and 2',4'-dihydroxy-6'-methoxychalcone (cardamonin, **3**), were isolated and displayed cytotoxic activity ($IC_{50} = 10, 35, \text{ and } 35 \mu\text{M}$, respectively) against the SW-480 human colon cancer cell line. Also a number of known antioxidants were obtained including six quercetin glycosides: reynoutrin (**4**), hyperin (**5**), myricitrin (**6**), quercitrin (**7**), quercetin (**9**), and guaijaverin (**10**), one flavanone: (*S*)-pinocembrin (**8**), and two phenolic acids: gallic acid (**11**) and ellagic acid (**12**).
© 2007 Elsevier Ltd. All rights reserved.

Keywords: *Syzygium samarangense*; Chalcones; Flavonoids; Cytotoxic activity; Antioxidants

1. Introduction

Syzygium samarangense (Bloom) Merr. and L.M. Perry (Myrtaceae), known commonly as wax jambu, is an evergreen tree with origins in Asia. It produces a pink fleshy fruit which is eaten fresh. Many cultivars have been developed and they are grown throughout the tropical and subtropical parts of the world. The fruit is oblong, pear-shaped, and 5–12 cm in length, with four fleshy calyx lobes and 1–4 seeds (1–2 cm in diameter). The tree can be grown as an ornamental, and attains a height of seven meters.

Wax jambu belongs to the same genus as *Syzygium aromaticum*, the source of cloves, a common spice.

In Malaysia, the green fruits of wax jambu are eaten raw with salt or cooked as a sauce. The flowers, which contain tannins, desmethoxymatteucinol, 5-*O*-methyl-4'-desmethoxymatteucinol, oleanic acid, and β -sitosterol, are used in Taiwan to treat fever and halt diarrhea (Morton, 1987).

Previous phytochemical studies of the leaves of *S. samarangense* have shown the presence of ellagitannins (Nonaka, Aiko, Aritake, & Nishioka, 1992), flavanones (Kuo, Yang, & Lin, 2004), flavonol glycosides (Kuo et al., 2004; Nair, Krishnan, Ravikrishna, & Madhusudan, 1999), proanthocyanidins (Nonaka et al., 1992), anthocyanidins (Kuo et al., 2004; Nonaka et al., 1992),

* Corresponding author. Tel.: +1 718 960 1105; fax: +1 718 960 8236.
E-mail address: edward.kennelly@lehman.cuny.edu (E.J. Kennelly).

triterpenoids (Srivastava, Shaw, & Kulshreshtha, 1995), chalcones (Resurreccion-Magno, Villasenor, Harada, & Monde, 2005; Srivastava et al., 1995), and volatile terpenoids (Wong & Lai, 1996).

Ethanol leaf extract exhibited immunostimulant activity (Srivastava et al., 1995), the hexane extract was found to relax the hypermotility of the gut (Ghayur et al., 2006), while the alcoholic extract of the stem bark showed antibacterial activity (Chattopadhyay, Sinha, & Vaid, 1998).

The immunomodulatory (Kuo et al., 2004), antihyperglycaemic (Resurreccion-Magno et al., 2005), spasmolytic (Amor, Villasenor, Ghayur, Gilani, & Choudhary, 2005), and prolyl endopeptidase inhibitor effects of chalcones **1** and **2** and the flavanone 5-*O*-methyl-4-desmethoxymatteucinol isolated from the leaves have also been reported (Amor, Villasenor, Yasin, & Choudhary, 2004).

Chalcones are a group of plant-derived polyphenolic compounds that are intermediates in the biosynthesis of flavonoids and are associated with several biological activities, including antiviral (Kiat et al., 2006), antifungal (Svetaz et al., 2004) anti-inflammatory (Lespagnol et al., 1972) and antioxidant (Han, Kang, Windono, Lee, & Seo, 2006). They also displayed anticancer and cytotoxic activity (Go, Wu, & Liu, 2005).

As part of a programme to find novel antioxidant and cytotoxic compounds from plants, (Yang et al., 2005), we have investigated the anticancer activity and antioxidant properties of wax jambu, and we report here the bioassay-guided identification of polyphenolic constituents and cytotoxic chalcones from the pulp and seeds of the fruits of *S. samarangense*.

2. Materials and methods

2.1. General methods

Optical rotation was measured on an Autopol III Automatic Polarimeter (Rudolph Research Analytical, Newburgh Hacktstown, NJ, USA); ^1H and ^{13}C NMR spectra were recorded using a Bruker Avance 300, operating at 300 and 75 MHz, respectively. Spectra were obtained in CD_3OD or CDCl_3 , with chemical shifts expressed in δ and coupling constant (J) in Hertz (Hz). Electrospray ionization mass spectrometry (ESI-MS) was performed with a ThermoFinnigan LCQ instrument (San Jose, CA, USA), equipped with Xcalibur software. Samples were dissolved in HPLC grade MeOH and introduced by direct injection. Nitrogen was used as both an auxiliary and sheath gas, the capillary voltage was 10 V, the spray voltage was 4.5 kV, the capillary temperature was 230 °C and the tube lens offset was 0 V. HPLC analyses were carried out on a Waters 2695 Separations Module (Milford, MA, USA) equipped with a model 996 photodiode array detector and Empower software, using a 250 × 4.6 mm i.d., 5 μm , Aqua C-18 column (Phenomenex, Torrance, CA, USA). Preparative HPLC was carried out using a Waters 600 controller with a Waters 486 tunable absorbance detector and Waters

Empower software with a 250 × 21.20 mm i.d., 10 μm Luna C-18 column (Phenomenex Torrance, CA, USA) and the mobile phase consisted of 10% aqueous formic acid (A) and HPLC grade acetonitrile (B). The flow rate was 20 ml/min, with a linear gradient consisting of 40% A–30% A in 20 min run time and the column at room temperature.

A Molecular Devices (Sunnyvale, CA, USA) Versamax tunable absorbance detector was used for the 1,1-diphenyl-2-picrylhydrazyl (DPPH) antiradical assay, total flavonoid content (TFC), and total phenolic content measurements (TPC).

TLC analyses were performed on Si gel 60 F₂₅₄ (1 mm layer thickness, EM Science, Darmstadt, Germany) and RP-18 F₂₅₄ plates (1 mm layer thickness, EM Science, Darmstadt, Germany), with compounds visualized by spraying with a vanillin solution (1.0 g of vanillin in 10 ml of concentrated H_2SO_4 and 90 ml of EtOH) and heating at 50 °C. Sephadex LH-20 (25–100 μm) (Pharmacia Fine Chemicals, Piscataway, NJ, USA), silica gel (230–400 mesh) (EM Science, Darmstadt, Germany), Diaion HP-20 (HP-20) (Supelco, Bellefonte, PA, USA), and C-18 reversed-phase Si gel (40 μm) (J. T. Baker, Phillipsburg, NJ, USA) were used for column chromatography. Solvents for chromatography, HPLC-grade acetonitrile, MeOH, formic acid, and HPLC-grade water, were obtained from J.T. Baker (Phillipsburg, NJ, USA), and GR-grade MeOH, EtOAc and acetone from VWR Inc (Bridgeport, PA, USA). AlCl_3 , DMSO, FeCl_3 , anhydrous Na_2CO_3 , NaNO_2 , NaOAc, NaOH, trolox, 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical, 2,4,6-tri-pyridyl-*s*-triazine, and the Folin–Ciocalteu phenol reagent were purchased from Sigma–Aldrich (St. Louis, MO, USA). Ellagic acid, gallic acid, myricitrin, quercetin, and quercitrin were also obtained from Sigma–Aldrich (St. Louis, MO, USA).

2.2. Plant material

Fruits of *S. samarangense* were collected from the Fruit and Spice Park (Homestead, FL, USA) in June 2001. Fresh fruits were shipped to New York City by overnight courier and stored at –20 °C until extracted. A voucher specimen (Reynertson 17) was prepared, identified, and deposited at the Steere Herbarium of The New York Botanical Garden (Bronx, NY).

2.3. Extraction

The fresh frozen pulp (3.2 kg) of the fruits of *S. samarangense* were extracted twice with MeOH (5 l) at room temperature for 1 h per extraction. After the MeOH was removed *in vacuo*, the resulting dark extract (27.0 g) was suspended in water and sequentially partitioned with hexane (1 l, ×3), EtOAc (1 l, ×3), and *n*-BuOH (1 l, ×3), respectively. The EtOAc and *n*-BuOH partitions were concentrated *in vacuo* to give 3.6 g and 18.0 g of dark brown extracts, respectively.

The seeds (400.0 g) of *S. samarangense* were processed in the same way as the pulp, and 15.0 g of a dark brown extract were obtained, which were suspended in water and sequentially partitioned with hexane (300 ml, $\times 3$), EtOAc (300 ml, $\times 3$), and *n*-BuOH (300 ml, $\times 3$). The EtOAc and *n*-BuOH partitions were concentrated *in vacuo* to give 1.7 g and 10.4 g of two dark brown residues, respectively.

Each selected fraction obtained was screened for free radical-scavenging capacity (Wu et al., 2005). Briefly, sub-fractions were loaded individually on a baseline of the RP-18 TLC. The TLC plate was developed using a 1:1 MeOH/H₂O solvent system and the layer was dried and stained with 0.2% DPPH[•] (w/v) solution in EtOH. The appearance of yellow colour in the spots indicates free radical-scavenging capacity of the test samples. The antioxidant activity of each active fraction was then assessed by the standard DPPH[•] assay.

2.4. Isolation and purification

The EtOAc partition (15.0 g) of the seeds was subjected to repeated column chromatography over Sephadex LH-20 using MeOH as eluent and eight subfractions (SEA-1–8) were collected. Fraction SEA-5 (28.0 mg, IC₅₀ = 18.3 μ g/ml in the DPPH[•] assay and IC₅₀ = 10.0 μ g/ml in MTT assay) was subjected to preparative C-18 HPLC to obtain 2',4'-dihydroxy-3',5'-dimethyl-6'-methoxychalcone (**1**, 14.0 mg), 2',4'-dihydroxy-3'-methyl-6'-methoxychalcone (**2**, 6.0 mg), and 2',4'-dihydroxy-6'-methoxychalcone (**3**, 0.6 mg).

The *n*-BuOH partition (10.4 g, IC₅₀ = 16.8 μ g/ml in the DPPH[•] assay and IC₅₀ = 55.0 μ g/ml in MTT assay) was processed using a similar procedure of EtOAc partition and 15.7 mg of gallic acid (**11**) and 12.9 mg of ellagic acid (**12**) were isolated.

The EtOAc fraction (1.8 g, IC₅₀ = 54.7 μ g/ml in the DPPH[•] assay and IC₅₀ = 20.0 μ g/ml in MTT assay) of the pulp was subjected to column chromatography over Sephadex LH-20 and eluted with an isocratic system using MeOH and 13 fractions (100 ml each) were collected. These fractions were combined into five fractions (EA-A to E) on the basis of RP-18 TLC (7:3 MeOH/H₂O) analysis. All fractions were tested in the MTT and DPPH[•] assays; fraction EA-C (1.1 g, IC₅₀ = 45.0 μ g/ml in the MTT assay and 25.0 μ g/ml in the DPPH[•] assay) and fraction EA-E (80 mg, 20.4 μ g/ml in the DPPH[•] assay) showed cytotoxic and antioxidant activities.

Fraction EA-C was chromatographed by Sephadex LH-20 (100% MeOH) and repeated RP-18 CC, eluting with a gradient of 3:7 MeOH/H₂O to 9:1 MeOH/H₂O (10% increments) to yield (*S*)-pinocembrin (**8**) (7.2 mg) and quercetin (**9**) (2.0 mg).

Fraction EA-E was chromatographed using the same procedure as described above to afford reynoutrin (**4**) (6.3 mg), hyperin (1.6 mg) (**5**), myricitrin (**6**) (0.5 mg), quercitrin (1.6 mg) (**7**), and guajaverin (**10**) (4.5 mg). The *n*-BuOH partition (18.0 g, IC₅₀ = 87.8 μ g/ml in the DPPH[•] assay) was processed using a similar procedure and 12.0 mg of gallic acid (**11**) and 11.6 mg of ellagic acid (**12**) were isolated.

Those nine compounds from the pulp and five compounds from the seeds of *S. samarangense* were identified by the spectroscopic methods and HPLC–PDA analysis with authentic standards (Fig. 1):

2',4'-Dihydroxy-3',5'-dimethyl-6'-methoxychalcone (**1**): yellow-orange crystals (14.0 mg) (Table 1), the yield was 35.0 mg/kg fresh weight from the seeds. Negative ESIMS: *m/z* 297 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Resurreccion-Magno et al., 2005).




Fig. 1. Structure of compounds (1–12) isolated from *S. samarangense*.

Table 1

Results of cytotoxic activity on the SW-480 human colon cancer cell line, antiradical DPPH assay, ferric reducing antioxidant power, total flavonoid content, and total phenolic content of methanol extracts and compounds **1–3** and **8** from the pulp and seeds of the fruits of *S. samarangense*

Compounds/methanol extract	Cytotoxic activity ^a , IC ₅₀	DPPH ^a , IC ₅₀	FRAP ^c	TFC ^d	TPC ^e
1	10	205 ± 1.2 ^b	196 ± 0.0	NT	NT
2	35	141 ± 2.3	191 ± 0.1	NT	NT
3	35	124 ± 3.4	173 ± 0.0	NT	NT
8	60	199 ± 0.8	196 ± 0.5	NT	NT
Pulp methanol extract	60	72.9 ± 0.0	14.8 ± 0.2	294 ± 0.0	460 ± 0.0
Seeds methanol extract	NS	78.4 ± 0.0	31.3 ± 0.2	19.9 ± 0.0	1278 ± 0.1
EGCG	50				
Gallic acid		25.0 ± 0.1			

NS: not completely soluble. NT: not tested.

^a Cytotoxic and antiradical DPPH activity are expressed as IC₅₀ in μM for pure compound and μg/ml for extracts.

^b Mean ± S.D. (*n* = 3).

^c Expressed as μM trolox equivalents/500 μmol for pure compound and μmol trolox/g fresh weight for extracts.

^d Total flavonoid content (TFC) expressed as mg quercetin/100 g fresh weight.

^e Total phenolic content (TPC) expressed as mg gallic acid/100 g fresh weight.

2',4'-Dihydroxy-3'-methyl-6'-methoxychalcone (stercurensin) (**2**): yellow-orange crystals (3.9 mg) (Table 1); the yield was 9.7 mg/kg fresh weight from the seeds. Negative ESIMS: *m/z* 283 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Resurreccion-Magno et al., 2005).

2',4'-Dihydroxy-6'-methoxychalcone (cardamonin) (**3**): yellow-orange crystals (0.8 mg) (Table 1); the yield was 2.0 mg/kg fresh weight from the seeds. Negative ESIMS: *m/z* 269 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Jaipetch et al., 1982).

Reynoutrin (**4**): yellow powder (6.3 mg); the yield was 2.37 mg/kg fresh weight from the fruits. Negative ESIMS: *m/z* 433 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Lu & Foo, 1997).

Hyperin (**5**): yellow powder (1.6 mg); the yield was 0.5 mg/kg fresh weight from the fruits. Negative ESIMS: *m/z* 463 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Lu & Foo, 1997).

Myricitrin (**6**): yellow powder (0.5 mg); the yield was 0.2 mg/kg fresh weight from the fruits. The compound was identified by HPLC–PDA analysis (retention time and UV spectrum) with authentic standard.

Quercitrin (**7**): yellow powder (1.6 mg); the yield was 1.4 mg/kg fresh weight from the fruits. The compound was identified by HPLC–PDA analysis (retention time and UV spectrum) with authentic standard.

(*S*)-Pinocembrin (**8**): white powder (7.2 mg); the yield was 2.2 mg/kg fresh weight from the fruits. Positive ESIMS: *m/z* 255 [M + H]⁺; ¹H and ¹³C NMR data are consistent with previously published data (Kuroyanagi et al., 1983). The specific rotation of **8** was measured as [α]_D^{29.0} = –25.9 (c, 0.004 in MeOH); therefore, **8** was identified as (*S*)-pinocembrin (Kuroyanagi et al., 1983).

Quercetin (**9**): yellow powder (2.0 mg); the yield was 0.6 mg/kg fresh weight from the fruits. The compound was identified by HPLC–PDA analysis (retention time and UV spectrum) with authentic standard.

Guaijaverin (**10**): yellow powder (4.5 mg); the yield was 1.5 mg/kg fresh weight from the fruits. Negative ESIMS:

m/z 433 [M–H][–]; ¹H and ¹³C NMR data are consistent with previously published data (Stark, Bareuther, & Hofmann, 2005).

Gallic acid (**11**): white powder (29.7 mg); the yield was 3.8 mg/kg fresh weight from the fruits and 39.8 mg/kg fresh weight from the seeds. Negative ESIMS: *m/z* 169 [M–1][–]; ¹H and ¹³C NMR data are consistent with previously published data (Khac, Tran-Van, Campos, Lallemand, & Fetizon, 1990). The compound was also identified by HPLC–PDA analysis (retention time and UV spectrum) with authentic standard.

Ellagic acid (**12**): brown powder (27.5 mg); the yield was 3.6 mg/kg fresh weight from the fruits and 44.3 mg/kg fresh weight from the seeds. Negative ESIMS: *m/z* 301 [M–1][–]; ¹H and ¹³C NMR data are consistent with previously published data (Khac et al., 1990). The compound was also identified by HPLC–PDA analysis (retention time and UV spectrum) with authentic standard.

The above mentioned 12 compounds and MeOH extracts of the pulp and seeds were tested for antioxidant activity in the DPPH[•] and FRAP assays and cytotoxic activity against SW-480 cell lines in the MTT assay (Table 1).

2.5. Total phenolic and total flavonoid contents (TPC and TFC)

The TPCs were determined by the Folin–Ciocalteu method (Yildirim, Mavi, & Kara, 2001). Briefly, 2.0 g of freeze-dried samples of the fruits–seeds were extracted twice with MeOH for 1 h at room temperature. Filtered MeOH extract (100 μl) and 10% Folin–Ciocalteu reagent (1.0 ml) in ethanol were placed in a test tube and incubated for 5 min at room temperature (22 °C). The mixture was left to stand for 5 min. and 1.0 ml of 10% sodium carbonate was added. After 90 min of incubation at room temperature, the resulting absorbance was measured at 765 nm. The calibration curve was performed with gallic acid, (concentrations ranging from 31.3 μg/ml to 500.0 μg/ml) and the results were expressed as μg of gallic acid equivalents per 100 g of fresh material. (Liu et al., 2002). The total fla-

vonoid contents of the extracts were determined. Briefly, 250 μ l of the sample (0.6 mg of the freeze dry extract per ml) was diluted with 1.25 ml of water. Then 75 μ l of 5% NaNO₂ solution were added to the mixture. After 5 min, 150 μ l of 10% AlCl₃ · 6H₂O were added and the mixture was allowed to stand for 5 min. Then 500 μ l of 1 M NaOH solution and 275 μ l of distilled water were added to make a total of 2.5 ml. The absorbance was measured immediately against the prepared blank at 510 nm. The results were expressed as milligrammes of quercetin equivalents per 100 g of fresh material. All data are reported as means \pm SD for at least three replications.

2.6. Cell culture

The SW-480 human colon cell cancer line was purchased from The American Type Culture Collection. The cell line was maintained in Dulbecco's modified Eagle's medium (Gibco-BRL, Grand Island, NY, USA) with 10% fetal bovine serum (Gibco) at 5% CO₂ and 37 °C. The culture was passaged weekly, and the medium was changed three times per week. No antibiotics were added at any time during the experiments. In all experiments, chalcones and plant extracts were dissolved in DMSO and added to the medium at the start of the incubation. The incubation time was 72 h for the MTT experiment.

2.7. Microtetrazolium (MTT) assay

The MTT assay (Boehringer-Mannheim, Indianapolis, IN, USA) was carried out according to the manufacturer's instructions. In brief, about 3000 cells were plated in 96-well flat-bottom plates in 100 μ l of medium. When cells reached 40% confluence, the medium was changed and cells were exposed to the plant extracts or isolates. After 72 h, cells were washed once with PBS followed by the addition of 100 μ l of Dulbecco's modified Eagle medium, and 10 μ l of 5 mg/ml MTT solution in PBS were added to each well for 6 h. Finally, 100 μ l of MTT solubilization solution were added to each well to dissolve the formazan crystals. The absorbance at 575 nm was determined using a Biokinetics plate reader (Bio-Tek Instruments Inc., Winooski, VT, USA). Octuplicate wells were assayed for each condition, and mean as well as standard deviations were determined. The IC₅₀ values were determined by linear regression analysis. Epigallocatechin gallate (EGCG) was used as a positive control (IC₅₀ = 50 μ M).

2.8. Antioxidant assays

Antioxidant properties of MeOH extracts and pure compounds isolated from the fruits and seeds of *S. samarangense* were determined by the DPPH free-radical-scavenging and the ferric reducing antioxidant power (FRAP) assays. The DPPH[•] assay was performed on fractions and purified isolates as previously described (Smith, Reeves, Dage, & Schnettler, 1987). Gallic acid was used as the posi-

tive control and percent inhibition by sample treatment was determined by comparison with a DMSO-treated control group. FRAP assay (Benzie & Strain, 1996) was adapted for use in a 96-well microplate spectrophotometer (Versamax Molecular Devices, Sunnyvale, CA, USA). FRAP values were expressed in μ mol trolox equivalents (TE) per g of fresh material for the extracts and μ mol TE per 500 μ mol of pure compounds.

3. Results and discussion

The methanol extract of the pulp and seeds of the fruits of *S. samarangense* (Table 1) were each partitioned with hexane, EtOAc and *n*-BuOH. The EtOAc partitions were each subjected to bioassay-guided fractionation with an initial separation by Sephadex LH-20 column chromatography. Subsequent purification of active fractions led to the isolation of four cytotoxic compounds against the SW-480 human colon cancer cell line: 2',4'-dihydroxy-3',5'-dimethyl-6'-methoxychalcone (**1**), stercurensin (**2**), cardamonin (**3**), and (*S*)-pinocembrin (**8**) (Table 1, Fig. 1), and eight well known antioxidants: reynoutrin (**4**), hyperin (**5**), myricitrin (**6**), quercitrin (**7**), quercetin (**9**), guaijaverin (**10**), gallic acid (**11**) and ellagic acid (**12**) (Fig. 1). Compounds **1–3**, **11**, and **12** were isolated from the seeds, and compounds **4–12** were obtained from the pulp. Compounds were identified using a combination of data from MS, NMR and HPLC–PDA comparison (retention time and UV spectra) with authentic standards. The yields of the 12 polyphenols isolated ranged from 0.2 to 3.8 mg per kg fresh weight from the pulp and 1.5–39.7 mg per kg fresh weight from the seeds, respectively. The total phenolic content, total flavonoid content, DPPH[•]-scavenging activity and FRAP results for the extracts are shown in Table 1.

Chalcones are flavonoids that lack the C ring. The presence of a 4-OH group, and an α , β -unsaturated double bond are essential features for the cytotoxic activity of this class of compound (Cai, Sun, Jie, Luo, & Corke, 2006). In a report on structure activity relationships of chalcones, a series of *O*-methylated chalcones showed activity against human tumor cell replication (KB, KB-VCR and A 549 cell lines) (Tatsuzaki et al., 2006). Another investigation showed that several ring-A methoxylated chalcones exhibited cytotoxic activity by inhibiting tubulin polymerization against a variety of tumor cell lines in the low micromolar range (IC₅₀ < 50 μ M) (Go et al., 2005).

Compounds **1**, **2**, and **3** are chalcones that have a 2'-hydroxy, a 4'-hydroxy and 6'-methoxy group substituting the A-ring. An uncommon feature is the additional C-methylation in **1** and **2**; these compounds have been previously isolated from the leaves of this plant and reported to have antidiabetic properties (Resurreccion-Magno et al., 2005), antibacterial (Gafner, Wolfender, Mavi, & Hostettmann, 1996), spasmolytic (Amor et al., 2005; Ghayur et al., 2006) properties, and inhibited prolyl endopeptidase (Amor et al., 2004).

This is the first report of the isolation of chalcone **3** from *S. samarangense*. This chalcone, cardamonin (2',4'-dihydroxy-6'-methoxychalcone), is one of the main constituents from the seeds of *Alpinia katsumadai*, and was first isolated from the seeds of *Amomum subulatum* (Bheemasankara, Namosiva Rao, & Suryaprakasam, 1976). It has anti-inflammatory activity (Lee et al., 2006) and anti-mutagenic effects upon activation of heterocyclic amines (Trakoontivakorn et al., 2001), and inhibits lipopolysaccharide-induced expression of inducible nitric oxide synthase and tumor necrosis factor- α in RAW 264.7 cells (Ban et al., 2004) and proinflammatory mediators (Syahida et al., 2006).

The flavanone (*S*)-pinocembrin (**8**), and the chalcones **1**, **2**, and **3**, displayed cytotoxic and weak antioxidant activity (Table 1). Compounds **4–7** and **9–10** had values higher than 100 μ M in the MTT assay, and compounds **11** and **12** were not tested in this assay. The lower cytotoxic activity of the flavanone (*S*)-pinocembrin (**8**, IC₅₀ = 60 μ M in the MTT assay) in relation to the chalcones isolated could be related to the rigidity of the flavanone structure (Fig. 1). The 5'-methyl group appears to be an important structural requirement for cytotoxic activity among the chalcones **1**, **2**, and **3**, as seen from the fall in activity when the structure loses this methyl group (10 μ M for compound **1** and 35 μ M for compounds **2** and **3**). The weak antioxidant activity displayed by the isolated chalcones could be attributed to the lack of hydroxyl groups in ring-B (Cai et al., 2006). There is no apparent correlation between cytotoxic and antioxidant activity.

The edible fruits of *S. samarangense* represent potential benefits for human health because they are a rich dietary source of polyphenolic antioxidants. In addition, the seeds are a rich source of the cytotoxic chalcones **1–3**, especially true for compound **1**, since it is present in high concentration (35.0 mg per kg fresh weight).

Acknowledgements

We would like to thank Chris Rollins, Director of the Fruit and Spice Park in Homestead, FL, and Larry Schokman, Director of the Kampong in Coconut Grove, FL for providing the fruits in this study. This research was supported by the funds from the National Institutes of Health-National Institute of General Medical Sciences SCORE award S06GM08225, NIH-NCCAM award F31000801, and from the United States Department of Agriculture.

References

Amor, E. C., Villasenor, I. M., Ghayur, M. N., Gilani, A. H., & Choudhary, M. I. (2005). Spasmolytic flavonoids from *Syzygium samarangense* (Blume) Merr. and I.M. Perry. *Zeitschrift fuer Naturforschung, C: Journal of Biosciences*, 60(1/2), 67–71.

Amor, E. C., Villasenor, I. M., Yasin, A., & Choudhary, M. I. (2004). Prolyl endopeptidase inhibitors from *Syzygium samarangense* (Blume) Merr. and I.M. Perry. *Zeitschrift fuer Naturforschung, C: Journal of Biosciences*, 59(1/2), 86–92.

Ban, H. S., Suzuki, K., Lim, S. S., Jung, S. H., Lee, S., Ji, J., et al. (2004). Inhibition of lipopolysaccharide-induced expression of inducible nitric oxide synthase and tumor necrosis factor- α by 2'-hydroxychalcone derivatives in raw 264.7 cells. *Biochemical Pharmacology*, 67(8), 1549–1557.

Benzie, I. F. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (frap) as a measure of antioxidant power: The frap assay. *Analytical Biochemistry*, 239(1), 70–76.

Bheemasankara, R. C., Namosiva Rao, T., & Suryaprakasam, S. (1976). Cardamonin and alpinetin from the seeds of *Amomum subulatum*. *Planta Medica*, 29(4), 391–392.

Cai, Y.-Z., Sun, M., Jie, X., Luo, Q., & Corke, H. (2006). Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Sciences*, 78(25), 2872–2888.

Chattopadhyay, E. D., Sinha, B. K., & Vaid, L. K. (1998). Antibacterial activity of *syzygium species*. *Fitoterapia*, 119(4), 365–367.

Gafner, S., Wolfender, J.-L., Mavi, S., & Hostettmann, K. (1996). Antifungal and antibacterial chalcones from *Myrica serrata*. *Planta Medica*, 62(1), 67–69.

Ghayur, M. N., Gilani, A. H., Khan, A., Amor, E. C., Villasenor, I. M., & Choudhary, M. I. (2006). Presence of calcium antagonist activity explains the use of *Syzygium samarangense* in diarrhoea. *Phytotherapy Research*, 20(1), 49–52.

Go, M. L., Wu, X., & Liu, X. L. (2005). Chalcones: An update on cytotoxic and chemoprotective properties. *Current Medicinal Chemistry*, 12(4), 483–499.

Han, A. R., Kang, Y. J., Windono, T., Lee, S. K., & Seo, E. K. (2006). Prenylated flavonoids from the heartwood of *Artocarpus communis* with inhibitory activity on lipopolysaccharide-induced nitric oxide production. *Journal of Natural Products*, 69(4), 719–721.

Jaipetch, T., Kangahe, S., Pancharoen, O., Patrick, V. A., Reutrakul, V., Tuntiwachwuttikul, P., et al. (1982). Constituents of *Boesenbergia pandurata* (syn. *Kaempferia pandurata*) isolation, crystal structure and synthesis of (+)-boesenbergin a. *Australian Journal of Chemistry*, 35, 351–361.

Khac, D., Tran-Van, S., Campos, M. A., Lallemand, Y., & Fetizon, M. (1990). Ellagic compounds from *Diplopanax stachyanthus*. *Phytochemistry*, 29, 251–256.

Kiat, T. S., Phippen, R., Yusof, R., Ibrahim, H., Khalid, N., & Rahman, N. A. (2006). Inhibitory activity of cyclohexenyl chalcone derivatives and flavonoids of Fingerroot, *Boesenbergia rotunda* (L.), towards dengue-2 virus ns3 protease. *Bioorganic & Medicinal Chemistry Letters*, 16(12), 3337–3340.

Kuo, Y.-C., Yang, L.-M., & Lin, L.-C. (2004). Isolation and immunomodulatory effects of flavonoids from *Syzygium samarangense*. *Planta Medica*, 70(1), 1237–1239.

Kuroyanagi, M., Noro, T., Fukushima, S., Aiyama, R., Ikuta, A., Itokawa, H., et al. (1983). Studies on the constituents of the seeds of *Alpinia katsumadai* Hayata. *Chemical & Pharmaceutical Bulletin*, 31(5), 1544–1550.

Lee, J.-H., Jung, H. S., Giang, P. M., Jin, X., Lee, S., Son, P. T., et al. (2006). Blockade of nuclear factor- κ B signaling pathway and anti-inflammatory activity of cardamonin, a chalcone analog from *Alpinia conchigera*. *Journal of Pharmacology and Experimental Therapeutics*, 316(1), 271–278.

Lespagnol, A., Lespagnol, C., Lesieur, D., Cazin, J. C., Cazin, M., Beerens, H., et al. (1972). Analgesic, antiinflammatory and antimicrobial activities of chalcones and dihydrochalcones derived from salicylic acid. *Chimica Therapeutica*, 7(5), 365–369.

Liu, M., Li, X. Q., Weber, C., Lee, C. Y., Brown, J., & Liu, R. H. (2002). Antioxidant and antiproliferative activities of raspberries. *Journal of Agricultural and Food Chemistry*, 50(10), 2926–2930.

Lu, Y., & Foo, Y. (1997). Identification and quantification of major polyphenols in Apple pomace. *Food Chemistry*, 59, 187–194.

Morton, J. (Ed.). (1987). *Fruits of Warm Climates* (pp. 386–388). Winterville, NC: Julia Morton.

- Nair, A. G. R., Krishnan, S., Ravikrishna, C., & Madhusudanan, K. P. (1999). New and rare flavonol glycosides from leaves of *Syzygium samarangense*. *Fitoterapia*, 70(2), 148–151.
- Nonaka, G., Aiko, Y., Aritake, K., & Nishioka, I. (1992). Tannins and related compounds. CXIX. Samarangenins a and b, novel proanthocyanidins with doubly bonded structures, from *Syzygium samarangense* and *S. aqueum*. *Chemical & Pharmaceutical Bulletin*, 40(10), 2671–2673.
- Resurreccion-Magno, M. H. C., Villasenor, I. M., Harada, N., & Monde, K. (2005). Antihyperglycaemic flavonoids from *Syzygium samarangense* (Blume) Merr. and Perry. *Phytotherapy Research*, 19(3), 246–253.
- Smith, R. C., Reeves, J. C., Dage, R. C., & Schnettler, R. A. (1987). Antioxidant properties of 2-imidazolones and 2-imidazolthiones. *Biochemical Pharmacology*, 36(9), 1457–1460.
- Srivastava, R., Shaw, A. K., & Kulshreshtha, D. K. (1995). Triterpenoids and chalcone from *Syzygium samarangense*. *Phytochemistry*, 38(3), 687–689.
- Stark, T., Bareuther, S., & Hofmann, T. (2005). Sensory guided decomposition of roasted cocoa nibs (*Theobroma cacao*) and structure determination of taste active polyphenols. *Journal of Agricultural and Food Chemistry*, 53(13), 5407–5418.
- Svetaz, L., Tapia, A., Lopez, S. N., Furlan, R. L. E., Petenatti, E., Pioli, R., et al. (2004). Antifungal chalcones and new caffeic acid esters from *Zuccagnia punctata* acting against soybean infecting fungi. *Journal of Agricultural and Food Chemistry*, 52(11), 3297–3300.
- Syahida, A., Israf, D. A., Lajisa, N. H., Shaaria, K., Mohamede, H., Wahaba, A. A., et al. (2006). Cardamonin, inhibits pro-inflammatory mediators in activated raw 264.7 cells and whole blood. *European Journal of Pharmacology*, 538(1–3), 188–194.
- Tatsuzaki, J., Bastow, K. F., Nakagawa-Goto, K., Nakamura, S., Itokawa, H., & Lee, K.-H. (2006). Dehydrozingerone, chalcone, and isoeugenol analogues as in vitro anticancer agents. *Journal of Natural Products*, 69(10), 1445–1449.
- Trakoontivakorn, K., Nakahara, H., Shinmoto, M., Takenaka, M., Onishi-Kameyama, H., Ono, M., et al. (2001). Structural analysis of a novel antimutagenic compound, 4-hydroxy-panduratin A, and the antimutagenic activity of flavonoids in a Thai spice, fingerroot (*Boesenbergia pandurata* Schult.) against mutagenic heterocyclic amines. *Journal of Agricultural and Food Chemistry*, 49(6), 3046–3050.
- Wong, K. C., & Lai, F. Y. (1996). Volatile constituents from the fruits of four *Syzygium* species grown in Malaysia. *Flavour and Fragrance Journal*, 11(1), 61–66.
- Wu, J.-H., Tung, Y.-T., Wang, S.-Y., Shyur, L.-F., Kuo, Y.-H., & Chang, S.-T. (2005). Phenolic antioxidants from the heartwood of *Acacia confusa*. *Journal of Agricultural and Food Chemistry*, 53(15), 5917–5921.
- Yang, H., Protiva, P., Gil, R. R., Jiang, B., Baggett, S., Basile, M. J., et al. (2005). Antioxidant and cytotoxic isoprenylated coumarins from *Mammea americana*. *Planta Medica*, 71(9), 852–860.
- Yildirim, A., Mavi, A., & Kara, A. (2001). Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. Extracts. *Journal of Agricultural and Food Chemistry*, 49(1), 4083–4089.