

The use of water and ice with bactericide to prevent onboard and onshore spoilage of refrigerated megrim (*Lepidorhombus whiffiagonis*)

Laura Pastoriza^{*}, Marta Bernárdez, Gabriel Sampedro, Marta L. Cabo, Juan J.R. Herrera

Instituto de Investigaciones Marinas (CSIC), Eduardo Cabello, 6-36208 Vigo, Spain

Received 30 August 2007; received in revised form 17 November 2007; accepted 24 January 2008

Abstract

This study investigates the effectiveness of ozonated water and flake ice (combined Petfrost system) to increase the quality and stability of fresh megrim on fishing boats. The captured fish were washed, placed in plastic boxes, covered with flake ice and refrigerated at 2 °C for up to 2-weeks onboard and, thereafter, for 11 days onshore. The experiments employed sterile, filtered and ozonated water at a concentration of 2 ppm for washing the fish and making the flake ice. The results are compared with samples from a traditional treatment consisting of water and flake ice of marine origin. Fish were caught in four different hauls, which took 14, 12, 8 and 3 days in being landed. Subsequently, fish were stored for 1, 5, 7 and 11 days at 3 °C. The different treatments were evaluated using sensory, microbiological and chemical techniques. Fish treated with ozone always showed the best quality. Megrim treated with ozone was still suitable for consumption after 14 days on board, and megrim stored for 12, 8 and 3 days on board could be stored for a further five days in the ice state once landed with an acceptable quality. In contrast, control fish were not suitable for consumption if stored for longer than three days on board. The results indicate that treatment with water and ice flakes made from sterile and ozonated water maintains the quality of fresh megrim onboard fishing boats and upon arrival onshore.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: Megrim; Ozonised water; Flake ice; Refrigeration; On board

1. Introduction

The Codex Alimentarius defines ozone as an antimicrobial agent and disinfectant for use in foodstuffs, both in the water destined for direct consumption and in ice or substances for indirect consumption, such as the water used to preserve fish, agricultural products and other perishable foods (Pérez, Palacios, & Amigo, 2006). In 2001, the US Food and Drug Administration (FDA) listed ozone as a safe (GRAS: Generally Recognized as Safe) antimicrobial agent for use in direct contact with food products, including fish, meat and poultry (Mielcke & Ried, 2006). The Electric Power Research Institute and the Agriculture and Food Technology Alliance submitted a petition to the FDA demanding the use of ozone in food processing

without limitations (Graham, 2000). In response, the FDA amended the food additive regulations to include gaseous and aqueous ozone as an antimicrobial agent (Molloy et al., 2001). This approval should encourage an increased use of ozone in food processing (Kim, Yousef, & Khadre, 2003).

Ozone at low concentration (0.01 ppm) is toxic for Gram-positive and Gram-negative bacteria (Mielcke & Ried, 2006) and has an oxidation potential 1.5 times greater than that of chlorine (Lin & Yeh, 1993). The bactericidal effect of ozone depends on the ozone concentration, the contact time, temperature and the production system used. Low ozone concentrations in solution decrease the half life of ozone in some cases and the bactericidal effect is unobserved. However, when the concentrations of ozone in solution are sufficiently high (~2 ppm), the effect of ozonated water and ice can improve the quality of refrigerated fish (Pastoriza et al., 2007). The decomposition of ozone in air occurs at much lower rates than in solution

^{*} Corresponding author. Tel.: +34 986 214465; fax: +34 986 292762.
E-mail address: laura@iim.csic.es (L. Pastoriza).

(Glaze, 1986) and it can react with other components on the surface of fish to produce a bactericide effect. Additionally, Ravesi, Licciardello, and Racicot (1987) reported that ozone may react with some components in sea water to produce a bactericidal ion or compound. The use of a gaseous ozone system provides fisherman with a practical alternative to improve catch quality and marketability (da Silva, Gibbs, & Kirby, 1998). The absence of adverse sensory effects and harmful oxidation by-products makes ozone a desirable antimicrobial agent in the processing of fish products for human consumption (Kim et al., 2003).

In Spain, the majority of fish catches sold at fresh fish markets comes from the Grand Sole fishing bank (North Atlantic). The trawlers typically spend 2 weeks at sea, during which the daily catch is stored in ice and refrigerated onboard. The storage time thus varies between 3 and 15 days, depending on which day the fish is caught, before arrival onshore and sale at auction. Among the marine flat fish, megrim is highly prized, by both domestic consumers and restaurants. The present study focusses on increasing the lifetime of megrim when refrigerated onboard. Two treatments are compared: (i) washing of the fish with ozonated water, followed by refrigeration in ozonated flake ice, and (ii) the traditional treatment employing sea water for the washing and ice-making. The quality and grade of freshness of the megrim once on land and stored for 11 days at 2–3 °C is also investigated.

2. Material and methods

2.1. Water and flake ice preparation

The Petfrost system was used to filter, sterilize and ozonate the water required for fish washing and flake ice production (Taboada, 2004). For both treatments, the flake ice was prepared onboard with a saline solution (0.10–0.15%):

(1) Control flake ice was prepared onboard in an ice machine using the homogenized saline solution (Icematic, Castel MAC, Castelfranco Veneto, Italia). This is termed control flake ice in the text. (2) Flake ice with bactericide was prepared from the saline solution using the Petfrost system with an ozone concentration of 2 ppm. This is termed Petfrost flake ice in the text.

2.2. Sample preparation

A total of 50 kg of megrim (length 22 ± 2 cm) caught by trawlers working in the North Atlantic during June 2006 was used for the experiments (Table 1). After the fish were landed, they were separated from the by-catch and washed with water from the Petfrost system for 3 s. Subsequently, the fish were placed in boxes with Petfrost flake ice in a 2:1 fish:ice ratio and stored onboard at 2 °C. The fish used in this work had been stored onboard and at sea for 14, 12, 8 and 3 days, and are identified in the text as batches 1,

Table 1
Sample preparation and days of storage of the batches

	Onboard	Onshore
Petfrost samples:	(Water + ice) Petfrost →	Ice Petfrost
Control samples:	(Water + ice) Control →	Ice Control
No. batch	Days onboard + Days onshore = Total days	
1	14	1, 5, 7 15, 19, 21
2	12	1, 5, 7 13, 17, 19
3	8	1, 5, 7, 11 9, 13, 15, 19
4	3	1, 5, 7, 11 4, 8, 10, 14

2, 3 and 4, respectively. The sealed and refrigerated fish boxes were sent to the Marine Research Institute (IIM, Vigo, Spain) where additional ice was added to the control and Petfrost boxes in the same 2:1 fish:ice ratio, and then refrigerated at 2 °C. These samples were compared with control samples which used saline water only (0.10–0.15%) for washing and flake ice production. Laboratory analyses of the fish were carried out after 1, 5, 7 and 11 days of their arrival at IIM. For both the control and Petfrost samples, only the fish stored onboard for 8 and 3 days were stored for a further 11 days onshore since the catches which had spent 12 and 14 days onboard were rejected on the basis of chemical, microbiological and sensory analyses.

2.3. Proximate analysis

Moisture, crude protein, ash and crude fat were measured in triplicate, following the procedures proposed by AOAC (1995). Moisture was determined by oven-drying at 105 °C. Crude protein was determined by mineralization of the sample with sulfuric acid and a Se/Cu catalyst, followed by distillation, and then analyzed with 0.1 N HCl using a 2300 Kjeltec Analyzer Unit (FossTecator, Höganäs, Sweden). Crude fat was measured by Soxhlet extraction of the fat with ethyl ether, and ash was determined by oven-heating at 550 °C for 24 h.

2.4. Microbiological analysis

Under sterile conditions inside a vertical laminar-flow cabinet (Telstar, AV-100, Tarrasa, Spain), 10 g of megrim muscle were placed in a sterilized plastic filter bag (Seward, Thetford, Norfolk, UK) with 90 ml of peptone water. Appropriate dilutions were prepared, following 1 min sample mixing in a stomacher blender (ITUL Instruments, 2997/400, Barcelona, Spain). The total number of aerobic microorganisms (total viable counts, TVC) was used as an indicator of the limit of product acceptance. The TVC analysis was performed on Plate Count Agar (Panreac, Barcelona, Spain) supplemented with 1% NaCl (Panreac) incubated at 17 °C for 3 days. Microbiological counts were expressed as log colony forming units per gramme of sample (CFU g⁻¹).

2.5. Chemical analysis

Triplicate samples of homogenized fish muscle with distilled water (ratio 1:2) were analyzed using an electrode (Crison, Barcelona, Spain) to determine the pH variation. Total volatile nitrogenous bases (TVB-N) were determined according to Lücke and Geidel (1935) and Antonacopoulos (1960). The analysis was based on titration with 0.1 N HCl, using a solution of a boric acid as indicator (pH 5.5), of a distillate of fish muscle triturate (10 g) in water and MgO (2 g) (Panreac, Barcelona, Spain). The results are expressed in mg TVB-N per 100 g of muscle. Trimethylamine-nitrogen (TMA-N) values were determined by the picrate method (Dyer, 1945), employing a 7.5% trichloroacetic acid (Panreac) extract of fish muscle (20 g/100 ml). The concentration is expressed in mg TMA-N per 100 g of muscle. Dissolved ozone concentration was measured by the indigo method of Bader and Hoigné (1981), in which the ozone reacts selectively with the carbon double bond of the sulfonated indigo molecule. Therefore, the ozone measurement was not affected by the presence of hydrogen peroxide, organic peroxides, manganese ions or oxidized species in the aqueous medium (Kim et al., 2003).

2.6. Sensory analysis

For raw megrim analysis, eight panellists were familiarized with the sensory analysis. The EU grading scheme (Council Regulation (EC) Nr 2406/96 of 26 November, 1996) was used to describe each attribute, choosing the attributes which avoided cutting or dissection. Gill colour was not intense and differences were observed between individuals rather than between the treatment methods, and this attribute was thus excluded from the evaluation. Each single descriptor was associated with demerit points on a scale of 0–3. In total, 19 demerit points were possible, and a quality index (QI) was calculated as the QI ratio = $ss/19$, where “ss” is the sum of the scores of each characteristic assessed. Thus QI ranged from 0 (best) to 1 (worst).

The sensory analysis of cooked fish was performed by 10 trained panellists and was based on colour, smell, taste, texture and palatability after 3–5 min of microwave cooking on a scale from 0 to 5 (Lupin, Giannini, Soulé, Davidovich, & Boeri, 1980). The fish was cooked whole, filleted and presented to the tasting panel. The total score for each characteristic indicates the loss in quality of the product, with a possible maximum of 25. The quality index of cooked fish (QIC) was calculated as the QIC ratio = $cs/25$, where “cs” is the sum of scores of each attribute of the cooked fish assessed. Thus, QIC ranged from 0 (best) to 1 (worst).

2.7. Statistical analysis of results

The results obtained for the control and Petfrost treatments, with regard to storage time, were subjected to an

analysis of variance (Statistica 6.0 software) for each sampling period, using a Student's *t*-test with a significance level of 95%. Statistical differences, shown in the Tables, are identifiable with different letters. Parameter correlations were also calculated using the statistical software.

3. Results and discussion

3.1. Proximate analysis

The approximate composition of megrim in $\text{g kg}^{-1} \pm \text{SD}$ was: water 782.6 ± 7.7 ; crude protein 191 ± 6.5 (total nitrogen $\times 6.25$), crude fat 19.3 ± 0.5 and ash 14.2 ± 0.7 .

3.2. Microbiological results

When compared to the untreated samples, the Petfrost treatment significantly reduced the bacterial population in the fresh megrim, with significantly lower ($p < 0.05$) TVC of undesirable bacteria commonly present in refrigerated fish (Fig. 1). Upon arrival onshore, after having been refrigerated onboard for 14, 12, 8 and 3 days, the bacterial counts were 5.75, 5.01, 4.77 and 4.27 log CFU g^{-1} , respectively, for the Petfrost samples and 6.26, 6.08, 5.49 and 5.04 log CFU g^{-1} for the control samples. In all cases, the differences between treatments were significant. At the IIM (day 1) the fish were stored at 2 °C, and the control samples which had been onboard for 14 and 12 days were unsuitable for consumption since they surpassed the limit of bacterial content established by European legislation (>1 million bacteria per gramme). However, the Petfrost samples had values below this limit and significant differences were observed between the treated and control samples from the 12 and 14 day batches. After 5 days of




Fig. 1. Evolution of total viable counts (TVC) in refrigerated megrim for Petfrost treatment (T: solid lines) versus control treatment (C: discontinuous lines). Samples follow the same convention; batch 1 (14 days onboard storage); batch 2 (12 days onboard storage); batch 3 (8 days onboard storage), and batch 4 (3 days onboard storage).

refrigeration, the control and Petfrost samples surpassed 1 million bacteria per gramme with significantly different bacterial counts.

When the samples which had been stored onboard for 8 days arrived onshore (day 1 at IIM, in total 9 days of storage), the total counts of aerobes between the control and Petfrost treatments were significantly different, yet in both cases lower than 6 log units (5.49 and 4.77 log CFU g⁻¹, respectively). After 5 days of storage at IIM at 2 °C, the samples again had significantly different values and the control samples were discarded (6.55 log units) whereas the treated samples were still within permitted limits (5.63 log units). Therefore, this Petfrost-treated fish was considered suitable for consumption after a total of 13 days of refrigeration.

The control and Petfrost-treated fish which arrived onshore after only 3 days onboard could be maintained refrigerated onshore for a further 5 and 7 days, respectively (a total of 8 and 10 days storage). However, subsequent analysis showed that the Petfrost samples could be refrigerated onshore for 11 days (14 days in total), at which point the total aerobic count was 5.9 log units. The corresponding control surpassed permitted limits (6.9 log units). In addition, bacterial counts were significantly different for the control and Petfrost samples after 1, 5, 7 and 11 days of refrigerated storage at IIM. In summary, megrim treated onboard with ozonated water and flake ice (Petfrost system) can be maintained refrigerated on land for a further 1, 5, 7 or 11 days, depending on the day of catch.

The antimicrobial affect of ozone was also noted for gutted mackerel by Haraguchi, Simidu, and Aiso (1969), who observed a logarithmic reduction in viable bacteria after submerging the fish in an aqueous solution of 3% NaCl containing 0.6 ppm of ozone for 30–60 min. Crapo, Himelbloom, Vitt, and Pedersen (2004) observed that ozonated water (0.6–1.5 ppm) was an effective bactericide for steel and plastic surfaces in fresh fish preparation rooms. Furthermore, da Silva et al. (1998) noted a 1 log CFU cm⁻² reduction in specific microorganisms in skin and muscle when fish was treated daily with ozone. Other authors concluded that ozone reduced contamination on the surface of fresh vegetables, and suggested that washing with ozonated water could be an effective disinfectant for foods (Karaca & Velioglu, 2007; Rico, Martin-Diana, Frias, Henehan, & Barry-Ryan, 2006; Selma, Beltran, Chacon-Vera, & Gil, 2006; Zambuchini, Giosia, & Sturba, 2006). Restaino, Frampton, Hemphill, & Painikar 1995 reported that ozonated water was highly effective in killing both Gram-negative and Gram-positive food-associated bacteria. The microorganisms usually responsible for food spoilage are Gram-negative bacteria, which cause tissue breakdown (Barriga, Trachy, Willomot, & Simard, 1991). In this work, it can be shown that ozonated water and ice were effective in reducing bacterial numbers on fresh fish, thus decreasing microbial spoilage and assuring food safety.

3.3. TMA-N

For the batches corresponding to 14, 12, 8 and 3 days after capture, the Petfrost fish showed significantly lower levels of TMA nitrogen (7.38, 1.91, 0.83 and 0.54 mg TMA-N/100 g) than the control fish (12.0, 3.85, 2.76 and 0.61 mg TMA-N/100 g), respectively (Fig. 2). Considering that the concentration limit established by European legislation is 12 mg/100 g TMA-N (Directive 91/493/EEC), the 14 day control sample was unsuitable for consumption upon arrival onshore (12.0 mg) in contrast to the Petfrost fish (7.38 mg).

The TMA-N values measured on the first day onshore after 12 and 8 days onboard storage (13 and 9 days total storage) were within legal limits for the control and treated samples. However, significant differences were observed between the two treatments. The increase in TMA-N levels after 5 days of storage at the IIM rendered the 13 and 9 day control batches unsuitable (16.5 and 12.9 mg, respectively), yet suitable (8.27 and 1.33 mg) for the Petfrost samples. After 7 and 11 days of refrigeration, only the Petfrost batch stored onboard for 8 days (a total of 15 and 19 days storage) had lower TMA-N values than the permitted limit (3.46 and 10.7 mg, respectively).

The fish from the batch refrigerated onboard for 3 days could be refrigerated for a further 11 days onshore (a total of 14 days storage) for the control and treated samples (5.24 and 2.95 mg TMA-N/100 g, respectively), whilst not exceeding permitted TMA-N limits. In this same batch, the amine values were significantly different for the control and Petfrost samples after 1, 5, 7 and 11 days of onshore refrigeration at the IIM.

3.4. TVB-N

Similar trends to the above were observed for TVB nitrogen (Table 2). When the fish arrived onshore (day 1)




Fig. 2. Evolution of trimethylamine nitrogen (TMA-N) in refrigerated megrim for Petfrost treatment versus control treatment. Sample identification follows the same convention as in Fig. 1.

Table 2
Changes in N-BVT during megrim storage in ice

Inshore days	C1 (14)*	C2 (12)	C3 (8)	C4 (3)
1	29.6 (1.80) ka	18.3 (0.43) ka	18.1 (0.53) ka	8.30 (0.70) ka
5	38.2 (1.25) la	30.3 (1.36) la	24.1 (0.18) la	13.7 (0.91) la
7	51.3 (0.54) ma	41.7 (1.79) ma	36.0 (0.12) ma	17.7 (1.12) la
11	NA	NA	45.0 (2.24) na	24.5 (0.88) ma
	T1 (14)	T2 (12)	T3 (8)	T4 (3)
1	22.4 (0.84) kb	13.4 (0.58) kb	13.4 (0.34) kb	7.96 (0.59) ka
5	35.4 (2.22) la	20.9 (1.00) lb	16.6 (0.40) lb	12.5 (0.75) la
7	49.2 (1.59) ma	26.3 (0.30) mb	19.9 (1.37) lb	15.7 (0.86) la
11	NA	NA	30.7 (0.99) mb	23.6 (0.42) ma

* (days) on board. Each value represents mean (standard deviation). Values followed by letters: a and b are significantly different ($p < 0.05$) between control and "Petfrost" batches (C1:T1, C2:T2, etc); $k - n$ between days of storage of each batch; NA is not analysed.

after 14 days onboard refrigerated storage, the Petfrost samples showed significantly lower levels of TVB (22.4, 13.4, 13.4, 7.96 mg TVB-N/100 g) than the samples treated with sea water and ice (29.6, 18.3, 18.1, 8.30 mg TVB-N/100 g). The presence of ozone in both water and flake ice thus had a favourable effect on TVB nitrogen for all fish, regardless of the day of capture.

The legal limit of TVB-N is 30 mg/100 g (Directive 95/149/EEC). Fish refrigerated for 7 days onshore and which had previously been stored on the trawler for 12 and 8 days (a total of 19 and 15 days of storage) showed permissible TVB levels for the Petfrost sample (26.3 mg and 19.9 mg) but not for the control (41.7 mg and 36.0 mg).

Finally, the TVB levels in fish stored onboard for 3 days and analyzed on day 1 onshore were within legal limits for both the control and treated batches. In all the control and treated samples, no significant differences in TVB levels were observed following refrigeration for 1, 5, 7 and 11 days and they were within legal limits.

Variability in the nitrogen compounds in fish occurs as a result of both endogenous enzymatic and bacterial activity (Matches, 1982). TMA-N and TVB-N are not always adequate for quantifying the freshness of all fish species (Civera, Turi, Bisio, Parisi, & Fazio, 1993; Price, Melvin, & Bell, 1991). With regard to megrim, however, there is a good

relationship between these two parameters and the level of freshness (Aubourg, Losada, Gallardo, & Barros-Velazquez, 2006) established according to European regulations (Civera, Turi, Parisi, & Fazio, 1995). In this study, and in agreement with Civera et al. (1995), both parameters indicate that the Petfrost system (ozonated water and flake ice) is more favourable than is the control (saline water and flake ice) for megrim preservation when refrigerating for 2 weeks onboard, followed by a further week on land or 11 days in some trials.

3.5. pH

The pH values varied according to the day the fish were caught, whereby samples with lower storage time, either onboard or on land, had a lower pH. The Petfrost-treated samples always had lower pH than had the control samples (Table 3). On arrival onshore (day 1), no significant differences were observed in pH between the control and Petfrost samples for any of the batches, and all showed pH < 7. After 5 days of refrigerated on land, the samples from batches corresponding to 12 and 8 days onboard showed significant differences in pH for the Petfrost (7.00 and 6.79) and control samples (7.22 and 7.06), respectively. The pH was always significantly higher in the control

Table 3
Change in pH values during megrim storage in ice

Inshore days	C1 (14)*	C2 (12)	C3 (8)	C4 (3)
1	6.88 (0.11) ka	6.78 (0.06) ka	6.72 (0.08) ka	6.65 (0.06) ka
5	7.39 (0.08) la	7.22 (0.03) la	7.06 (0.04) la	6.69 (0.04) ka
7	7.87 (0.04) ma	7.62 (0.10) ma	7.58 (0.13) ma	6.75 (0.07) ka
11	NA	NA	7.87 (0.10) na	7.06 (0.13) la
	T1 (14)	T2 (12)	T3 (8)	T4 (3)
1	6.85 (0.08) ka	6.74 (0.04) ka	6.70 (0.03) ka	6.59 (0.04) ka
5	7.33 (0.13) la	7.00 (0.06) lb	6.79 (0.07) kb	6.61 (0.06) ka
7	7.44 (0.07) lb	7.19 (0.06) lb	6.88 (0.07) klb	6.70 (0.04) kla
11	NA	NA	7.01 (0.06) lb	6.90 (0.08) la

Each value represents mean (standard deviation). Values followed by different letters: a and b are significantly different ($p < 0.05$) between control (C) and "Petfrost" batches (T); $k - n$ between days of storage of each batch.

* (days) on board; NA is not analysed.

samples. For the batch stored onboard for 3 days, no significant differences were observed in pH for the control and Petfrost samples after 1, 5, 7 and 11 days of refrigeration on land, and in all cases the pH values were <7. Following fish capture, the pH of the muscle tissue for the majority of fish species is usually below 7. This is due to the presence of lactic acid from hydrolysis of the glycogen produced after death. Thereafter, bacterial action produces

undesirable alkaline compounds, such as ammonia and TMA, which increase the pH to 7.0 and above, at which point the fish are usually discarded (Connell, 1980; Hebard, Flick, & Martin, 1982). In the present study, all samples with pH ~7 or less were deemed acceptable for consumption on the basis of the sensory characteristics of the fresh fish.

3.6. Sensory results

The sensorial examination of the fresh fish followed the attributes described in Table 4. The samples with lowest values had a higher level of freshness, and corresponded to the batches which were stored onboard for the least number of days. In general, the Petfrost samples showed a higher sensorial quality throughout storage and had lower values than had the control (Fig. 3). The microbiological data were used to establish the limit of acceptability of the fish. Applying the Pareto rule, only 20% of the cases had a log value >6 in the interval 0–0.65, and no sample with a value of 0.65 or higher had log value below log 6. This threshold coincides with the limit of freshness in Mediterranean hake, applying the ROC curve (Baixas-Nogueras, Bover-Cid, Veciana-Nogués, Nunes, & Vidal-Carou, 2003). According to this criterion, days of difference between the control samples and the treated samples in each of the batches were established, such that the control of batch 1 was unsuitable for consumption upon arrival onshore and the treated sample was not suitable after 4 days of storage on land. For batch 2, the control was unsuitable on day 4 and the treated sample on day 6. For batch 3, the control was unsuitable on day 6 yet the treated was still acceptable on day 11. Finally, both the control and treated fish from batch 4 were still suitable for consumption on day 11.

The condition of the cooked fish filets was better for the Petfrost samples than for the controls, although the differences were not always significant. In this case, half of the

Table 4
Sensory attributes quantified in raw megrim

Parameter	Characteristic	Points	
Dorsal zone	Aspect	Vivid pigment	0
		Vivid pigment, but without sheen	1
		Pigment shows discoloration and lacklustre	2
		Dull, lacklustre pigmentation	3
Mucus		Aqueous, transparent	0
		Lightly opaque	1
		Milky	2
Skin		Yellow-grey, opaque	3
		Smooth surface	0
Rigidity		Wrinkled surface	1
		Firm	0
Eye	Pupils	Lightly flaccid, less elastic	1
		Clear and black, metallic sheen	0
	Forma	Dark grey	1
		Lacklustre, grey	2
Gills	Mucus	Convex	0
		Flat	1
		Sunken, concave	2
Ventral zone	Mucus	Transparent	0
		Lightly turbid	1
	Odour	Milky	2
		Marine algae	0
Colour	Absence of algal smell, neutral	1	
	Fermented, slightly bitter	2	
	Bitter, putrid	3	
Colour	White or light cream without mucus	0	
	Cream or yellow with mucus	1	
	Greyish colour with abundant mucus	2	




Fig. 3. Evolution of the sensory score for the Petfrost treatment versus control treatment for raw (QI in bars) and cooked (QIC in lines) megrim. C1–C5 are control batches and T1–T5 are Petfrost batches (see Fig. 1 legend).

scale is the reference value for the suitable samples, since values >0.5 represent samples with undesirable sensory characteristics. On day 4 of storage there were significant differences between the control and Petfrost samples for batches 1 and 2. In batch 3 the most prominent result corresponds to day 11 of storage, in which the control sample was inedible yet the Petfrost sample was still acceptable. In batch 4, all the values were <0.5 , but the differences between the control and Petfrost were significant throughout the storage.

Sensory analyses of raw fish and cooked fish were considered jointly. Results for raw fish were split into six intervals (0.05–0.2–0.35–0.5–0.65–0.8–0.95) and arithmetic means were plotted against means for sensory analyses of the corresponding cooked fish. In the resulting equation ($y = 0.5077 \ln(x) + 0.9631$; $r^2 = 0.9791$), a value 0.65 for raw fish corresponds to a value of 0.54 for cooked fish. This calculation has been made by excluding three cases for which no explanation between raw fish and cooked fish was found. In cooked fish, panellists noticed the largest differences in odour and texture. Odour was acid and texture had no consistency for values of 0.5. Hyldig and Green-Petersen (2004) pointed out that the sensory quality of a fish/fish product, that the consumer buys, depends on the quality of the raw material used for the final product sold at retail level. The selection of the best descriptors for the spoiling fish allows sensory quality of fish to be objectively determined. This quality corresponds to a typical odour, colour, flavour and/or texture of the cooked product, which is particular for each fish species.

In summary, megrim washed with ozonised water on board and stored in the ice state, by using bactericide ice flakes (both on board and on land), showed a better quality than did control fish. Also the quality of the Petfrost ice flakes was different from that of the control flakes. Most notably, the flakes maintained their initial appearance and form for the first 2 weeks of refrigeration, remained loose and came apart easily, and prevented the formation of air pockets around the fish. Accordingly, the Petfrost flake ice facilitates the ice-fish contact during storage and enhances the quality of the fresh megrim. Thus, megrim treated with ozone was still suitable for consumption after 14 days on board, and megrim stored for 12, 8 and 3 days on board could be stored for five further days in the ice state once landed, with an acceptable quality. In contrast, control fish were not suitable for consumption if stored for longer than three days on board. Similarly, under conditions in which both ozone-treated fish and control fish were acceptable (3 days ice-stored on board and 7 days ice-stored on land), differences were significant too, so the treatment always renders a product with a higher quality.

4. Conclusions

When megrim is washed with ozonised water on board and stored in the ice state by using bactericide ice flakes,

which is called the Petfrost combined system, it can maintain higher quality than that of fish treated with the traditional system (sea water for washing and making flake ice). It allows a better acceptance of megrim at the retail outlets, and increases its shelf-life.

Acknowledgements

The authors wish to thank Alberto Gallego, Bibiana Torres and Carlos Suárez for their technical assistance. This research was supported by Empresa Peter Taboada (Pontevedra-Spain) and by the Xunta de Galicia, project (PGIDIT04TAL029E).

References

- Antonacopoulos, N. (1960). Verbesserte apparatur zur quantitative destillation wasserdampflichtiger stoffe. *Zeitschrift für Lebensmitteluntersuchung und -Forschung A*, 113(2), 113–116.
- AOAC (1995). *Official methods of analysis* (16th ed.). VA: Association of Official Analytical Chemists, Arlington.
- Aubourg, S. P., Losada, V., Gallardo, J. M., & Barros-Velazquez, J. (2006). On-board quality preservation of megrim (*Lepidorhombus whiffiagonis*) by a novel ozonised-slurry ice system. *European Food Research and Technology*, 223, 232–237.
- Bader, H., & Hoigné, J. (1981). Determination of ozone in water by the indigo method. *Water Research*, 1, 449.
- Baixas-Nogueras, S., Bover-Cid, S., Veciana-Nogues, T., Nunes, M. L., & Vidal-Carou, M. C. (2003). Development of quality index method to evaluate freshness in mediterranean hake (*Merluccius merluccius*). *Journal of Food Science*, 68, 1067–1071.
- Barriga, M. I., Trachy, G., Willomot, C., & Simard, R. E. (1991). Microbial changes in shredded iceberg lettuce stored under controlled atmosphere. *Journal of Food Science*, 56, 1586–1588.
- Civera, T., Turi, R. M., Bisio, C., Parisi, E., & Fazio, G. (1993). Sensory and chemical assessment of marine teleosteans. *Scienze Alimentari*, 13, 109–117.
- Civera, T., Turi, R. M., Parisi, E., & Fazio, G. (1995). Further investigations on total volatile basic nitrogen and trimethylamine in some Mediterranean teleosteans during cold storage. *Scienze Alimentari*, 15, 179–186.
- Connell, J. J. (1980). *Control of fish quality* (2nd ed.). Farnham, Surrey, England: Fishing News Books Ltd.
- Council Regulation (EC) No. 2406/96 of 26 November (1996). Laying down common marketing standards for certain fishery products. *Official Journal of the European Communities*. Nr L-334/2.
- Crapo, C., Himelbloom, B., Vitt, S., & Pedersen, L. (2004). Ozone efficacy as a bactericide in seafood processing. *Journal Aquatic Food Production Technology*, 13, 111–123.
- da Silva, M. V., Gibbs, P. A., & Kirby, R. M. (1998). Sensorial and microbial effects of gaseous ozone on fresh scad (*Trachurus trachurus*). *Journal of Applied Microbiology*, 84, 802–810.
- Dyer, W. J. (1945). Amines in fish muscle I Colorimetric determination of trimethylamine as the picrate salt. *Journal of the Fisheries Research Board of Canada*, 6, 351–358.
- Glaze, W. H. (1986). Reaction products of ozone: a review. *Environmental Health Perspectives*, 69, 151–157.
- Graham, D. M. (2000). Direct food additive petition: Ozone as an antimicrobial agent for the treatment, storage, and processing of foods in gas and aqueous phases. *FDA*, submitted for publication.
- Haraguchi, T., Simidu, U., & Aiso, K. (1969). Preserving effect of ozone on fish. *Bulletin of the Japanese Society of Science Fish*, 35, 915–919.

- Hebard, C. E., Flick, G. J., & Martin, R. E. (1982). Occurrence and significance of trimethylamine oxide and its derivatives in fish and shellfish. In R. E. Martin, G. J. Flick, C. E. Hebard, & D. R. Ward (Eds.), *Chemistry and biochemistry of marine food products* (pp. 149–304). Westport (USA): Avi.
- Hyldig, G., & Green-Petersen, D. M. B. (2004). Quality index method – An objective tool for determination of sensory quality. *Journal of Aquatic Food Product Technology*, 13(4), 71–80.
- Karaca, H., & Velioglu, Y. S. (2007). Ozone applications in fruit and vegetable processing. *Food Review International*, 23, 91–106.
- Kim, J. G., Yousef, A. E., & Khadre, M. A. (2003). Ozone and its current and future application in the food industry. *Advances in Food and Nutrition Research*, 45, 167–218.
- Lin, S. H., & Yeh, K. L. (1993). Looking to treat wastewater? Try ozone. *Chemical Engineering*, 100(5), 112–114, 116.
- Lücke, F., & Geidel, W. (1935). Bestimmung des flüchtigen basischen Stickstoffs in Fischen als Maßstab für ihren Frischezustand. *Zeitschrift für Lebensmitteluntersuchung und – Forschung A*, 70(6), 441–458.
- Lupin, H. M., Giannini, D. H., Soulé, C. L., Davidovich, L. A., & Boeri, R. L. (1980). Storage life of chilled patagonian hake (*Merluccius hubbsi*). *Journal of Food Technology*, 159, 285–300.
- Matches, J. R. (1982). Effect of temperature on the decomposition of Pacific coast shrimp (*Pandalus jordani*). *Journal of Food Science*, 47, 1044–1047, 1069.
- Mielcke, J., & Ried, A. (2006). Current state of application of ozone and UV for food processing. In Proceedings of the food protection international conference 2004. Monte da Caparica, Portugal, 20–22 of May.
- Molloy, E., Hassenberg, K., Plöchl, M., Idler, C., Geyer, M., & Barnwes, J. (2001). Secondary direct food additives permitted in food for human consumption. *Federal Register*, 66, 33829–33830.
- Pastoriza, L., Bernárdez, M., Sampedro, G., Cabo, M. L., & Herrera, J. J. R. (2007). Use of sterile and ozonized water and as a strategy to stabilize the quality of stored refrigerated fresh fish. *Food Control*. doi: 10.1016/j.foodcont.2007.08.003.
- Pérez, M., Palacios, A., & Amigo, P. (2006). Estudio de indicadores de la calidad de tomate conservado en atmósfera ozonizada. *Alimentaria*, 115, 124–129.
- Price, R. J., Melvin, E. F., & Bell, J. W. (1991). Postmortem changes in chilled round, bled and dressed Albacore. *Journal of Food Science*, 56, 318–321.
- Ravesi, E. M., Licciardello, J. J., & Racicot, L. D. (1987). Ozone treatment of fresh Atlantic cod, *Gadus morhua*. *Marine Fisheries Review*, 49, 37–42.
- Restaino, L., Frampton, E. W., Hemphill, J. B., & Painikar, P. (1995). Efficacy of ozonated water against various food-related microorganism. *Applied and Environmental Microbiology*, 61, 3471–3475.
- Rico, D., Martín-Diana, A. B., Frias, J. M., Henehan, G. T. M., & Barry-Ryan, C. (2006). Effect of ozone and calcium lactate treatment on browning and texture properties of fresh-cut lettuce. *Journal of the Science of Food and Agriculture*, 86, 2179–2188.
- Selma, M. V., Beltran, D., Chacon-Vera, E., & Gil, M. I. (2006). Effect of ozone on the inactivation of *Yersinia enterocolitica* and the reduction of natural flora on potatoes. *Journal of Food Protection*, 69, 2357–2363.
- Taboada, P. J. M. (2004). Equipo para la obtención de agua salobre ozonizada para ser utilizada como medio refrigerador y germicida de productos pesqueros. España AR038110 A1. Patent A23B 4/24, A23L 3/358, A22C 25/00.
- Zambuchini, B., Giosia, L., & Sturba, M. (2006). Study of the ozone effect on the shelf life of minimally processed salad. *Industria Alimentari*, 45, 636–643.