



Partial dehydration of cherry tomato at different temperature, and nutritional quality of the products

Giuseppe Muratore*, Valeria Rizzo, Fabio Licciardello, Emanuele Maccarone

Sezione Tecnologie Agroalimentari, Dipartimento di Orto-Floro-Arboricoltura e Tecnologie Agroalimentari (DOFATA), University of Catania, Via S. Sofia, 98 – 95123 Catania, Italy

ARTICLE INFO

Article history:

Received 16 July 2007

Received in revised form 5 March 2008

Accepted 1 May 2008

Keywords:

Cherry tomato

Partial drying

L-ascorbic acid

β-Carotene

Lycopene

Hydroxymethylfurfural

ABSTRACT

This study concerns the partial dehydration of cherry tomatoes (*Lycopersicon esculentum*, var. Shiren) to obtain a product with 25% initial water content. Two kinds of dried tomatoes were obtained using a forced air oven at 40, 60 and 80 °C for different lengths of treatment. The first type was dehydrated after immersion of the fresh tomatoes in an aqueous solution of citric acid, sodium and calcium chloride (10:10:24 g/l); the second was obtained with no pre-treatment. The products were characterised by measuring their CIE $L^*a^*b^*$ colour parameters and levels of L-ascorbic acid, lycopene and β-carotene to evaluate thermal damage during processing under the different conditions. Moreover, water activity and the formation of 5-hydroxymethylfurfural were also determined as an index of sugar heat degradation. Treatment with a dipping solution protected both the nutritional and chemical qualities of the partially dried cherry tomatoes. Temperature was directly related to browning, ascorbic acid loss and HMF formation, while no clear influence could be found for carotenoid degradation.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

A significant part of cherry tomato production is concentrated in a brief period during which their value drops to below cost. Therefore, producers aim at alternatives to commercialise excess produce, hopeful of some profit. This can be achieved by creating a new product, which is stable and marketable all year round. Being a popular fruit, tomatoes find numerous uses in both fresh and processed forms. Processed products include ketchup, sauces, pastes and juice. Drying is not a popular way to process tomatoes due to its adverse effect on final product quality (Lewicki, Vu Le, & Pomarańska-Lazuka, 2002) such as tissue browning and remarkable changes in the flavour profile. Many differences in general composition have been highlighted between traditional varieties (big tomatoes) and the new small-sized varieties (cherry and plum tomatoes), the latter characterised by higher dry matter and soluble solid fraction, essentially due to the higher levels of sugars and organic acids (Muratore, Licciardello, & Maccarone, 2005).

Much research has been carried out in order to correlate the assumption of tomatoes and their derivatives, with the ability to prevent some kinds of cancers and cardiovascular diseases (Giovannucci et al., 1995; Giovannucci, 1999; La Vecchia, 1998). These studies have demonstrated that thermal treatment of tomatoes (in every commercial product) positively correlates with low risk for cancers of the digestive tract and prostate. Other works (Nakaga-

wa, Yokozawa, Teresawa, Shu, & Raj Juneja, 2002; Russo et al., 2000) have confirmed that the consumption of natural substances capable of lowering oxidants, can protect against epithelial cancers and other diseases. Abushita, Hebshi, Daood, and Biacs (1997), Crozier, Lean, Mc Donald, and Black (1997), and Sies and Stahl (1998) have highlighted that the biological value of tomatoes is undoubtedly related to their high concentration of antioxidants, such as carotenoids, ascorbic acid and phenolic substances; in particular, lycopene acts as a “scavenger” against free radicals reducing the risk of cancer in humans.

Lycopene, according to the findings of Gartner, Stahl, and Sies (1997) and Stahl and Sies (1992), is stable during heating and industrial treatment, and treatments are able to improve lycopene bioavailability. Nevertheless, research carried out by Shi, Le Maguer, Kakuda, Liptay, and Niekamp (1999) showed significant loss in lycopene content during the dehydration of tomato products. Processes like cooking, cooling and canning do not usually cause large changes in total lycopene content, but during conventional tomato management, most lycopene can be converted from the all-*trans* form into the *cis* isomer which is less bioactive.

The aim of this work was to examine the effects of the partial dehydration of cherry tomatoes at different temperatures, in a forced air oven, with special regard to the effect of a pre-treating solution on the main physical, chemical and organoleptic characteristics of the products, namely, colour, water activity, dry matter, L-ascorbic acid, lycopene, β-carotene and 5-hydroxymethylfurfural. Many studies deal with the drying of tomatoes, but none of these has considered cherry tomatoes as a partially dehydrated product

* Corresponding author. Tel.: +39 95 7580210; fax: +39 95 7141960.
E-mail address: g.muratore@unict.it (G. Muratore).

with 25% initial water content. Today, the market shows increasing interest in products with intermediate humidity, which combine increased stability, due to lower water activity, with good nutritional and organoleptic characteristics. Partially dehydrated cherry tomatoes could be utilised as seasoning or to replace fresh tomatoes as the main ingredient in starters and other recipes.

2. Materials and methods

2.1. Sample preparation

Cherry tomatoes, cultivar “Shiren”, were collected at a local farm when ripe. They were selected by colour and homogenous diameter (29–32 mm), washed with tap water and cut longitudinally in halves.

The samples were divided into two batches, one immersed for 20 h in a solution of sodium chloride (10 g/l), citric acid (10 g/l) and calcium chloride (24 g/l) in tap water. The other was used as the control. Sodium chloride is used in ancient Sicilian tradition to prepare dried tomatoes, citric acid was used to guarantee lower pH for a longer shelf life, and calcium chloride was added because, according to a previous work, it increases the amount of water removed during dehydration (Lewicki et al., 2002).

2.2. Dehydration

Cut tomatoes were placed on four grates. Drying was performed in a forced air oven G[®]-Therm 075 (Galli, Milan, Italy) with the following characteristics: heating power, 1330 W; volume, 75 l; forced air speed, 2000 rpm. The following drying temperatures were chosen for treated and untreated samples: 40, 60 and 80 °C. The tomatoes were dried to 25% of their initial water content.

2.3. Quantitative measurements

For the fresh sample, pH, °Brix, dry matter and colour (CIE parameters) were determined.

The partially dehydrated cherry tomatoes (16–20 halves, 160–200 g), both raw and treated samples, were homogenised immediately at the end of drying and their dry matter and colour were measured. Samples with similar weight and dimensions were packed in PET/PP bags (Plastar Pak s.r.l., Milan, Italy) in a partial vacuum to reduce headspace, and then thermally sealed (Delta model “Delta 30”, Brindisi, Italy) stored at –18 °C, and thawed at +4 °C before analysis.

All the analyses were carried out in duplicate runs: dry matter was determined for 2 g homogenised samples in a thermo ventilated oven at 105 °C until constant weight; pH was determined for five homogenised fresh fruits by digital pH meter (MP 220, Mettler Toledo, Greifensee, Switzerland); total soluble solids were measured at 25 °C by refractometer (2WAJ ABBE bench Refractometer, Optika Microscopes, Bergamo, Italy); the colour coordinates L^* , a^* , b^* , C (colour chromaticity) and h (colour intensity) were measured by calorimeter NR-3000 (Nippon Denshoku Ind. Co. Ltd, Japan).

The colour of fresh cherry tomatoes was measured from the puree of 8–10 halves for both fresh and dehydrated samples. Water activity was determined at 25 °C for fresh and partially dried samples by hygrometer AwVC (Rotronic AG, Bassersdorf, Switzerland).

2.4. Carotenoids

Carotenoids were extracted according to De Sio, Grimaldi, and Louidice (1999); in particular, for partially dehydrated samples, the initial extraction was modified as follows: 2 g were mixed with

50 ml dichloromethane/methanol 2:1 (v/v) solution with 0.1% BHT and the extraction was repeated three times, using a total of 150 ml extraction solution.

Quantification was performed by HPLC (Finnigan SpectraSystem P 2000, Thermo scientific, Waltham, MA, USA) equipped with a Diode Array Detector UV 6000 LP, a Waters-YMC C-30 column for carotenoids (5 µm, 250 mm × 4.6 mm i.d.), and a 20 µl loop. Carotenoids were separated by 95:5 (v/v) methanol/H₂O as mobile phase A and dichloromethane with 0.1% BHT and 0.05% triethylamine (TEA) as mobile phase B. The solvent program, at a flow rate of 0.8 ml/min, was: 95% A for 2 min; 95% A to 30% A linearly in 8 min; followed by 5 min isocratic; 30% A to 10% A linear increase in 5 min; 10 min isocratic and a return to 95% A at 40 min to restore the initial conditions. Triethylamine was used to reduce carotenoid misplaced in the column and to improve peak shape.

The spectra were recorded in the range 320–480 nm, while chromatograms were acquired at wavelength (λ) 480 and 460 nm to determine lycopene and β -carotene. Quantification was made by calibration using appropriate dilutions of external standard lycopene and β -carotene (Sigma-Aldrich, Milan, Italy). All solvents were HPLC grade (J. T. Baker, Deventer, Holland).

2.5. L-ascorbic acid

Ascorbic acid was extracted and quantified by HPLC according to Nisperos-Carriedo, Buslig, and Shaw (1992). HPLC separation was by Finnigan SpectraSystem P 2000 (Thermo scientific, Waltham, MA, USA) equipped with a Diode Array Detector UV 6000 LP, a Supelcosil LC-18 (5 µm, 25 mm × 4.6 mm i.d.) column and a 20 µl loop.

Ascorbic acid was separated by 2% KH₂PO₄ buffer solution at pH 2.3 as mobile phase, at a flow rate of 0.5 ml/min. The spectra were recorded in the range 245–260 nm, where maximum absorbance for ascorbic acid is at 245 nm, but the acquisition of chromatograms was set at 260 nm to avoid interference (Finley & Duang, 1981). The identification and quantification was made by the external standard method, using a pure commercial standard (Carlo Erba Reagenti, Milan, Italy).

2.6. 5-Hydroxymethylfurfural (HMF)

The extraction of HMF was performed by homogenising 10 g of sample with 20 ml HPLC grade water for 15 min; samples were filtered through 0.45 µm nylon filters, and injected into the HPLC. The HPLC was a Varian 9012Q liquid chromatograph equipped with a diode array detector (Varian, Star 330), a Merk Lichrospher 100 RP-18 (5 µm, 125 mm × 4 mm i.d.) column, and a 20 µl loop. HMF was separated using H₂O/acetonitrile 95:5 (v/v) as mobile phase, at a flow rate of 1 ml/min. Chromatograms were acquired from 250 to 300 nm but HMF was monitored at 285 nm. Quantification was performed by external standard calibration using a calibration line obtained with appropriate dilutions of commercial standard (Sigma, Milan, Italy). Water and acetonitrile were HPLC grade (Merck, Darmstadt, Germany).

2.7. Statistical analysis

All analyses were carried out in duplicate. The experimental data was processed by Duncan test using Statgraphics Plus 5.1 software (Manugistic Inc. Rockville, MD, USA). The means and standard deviations were calculated. The data shown is the average of all repetitions. The sets of data obtained for each parameter were statistically treated to obtain a 95% confidence interval.

3. Results and discussion

3.1. Characterization of fresh and treated products

The fresh cherry tomato var. Shiren is characterised by a pH of 4.05, 7.06% soluble solids and 8.15% dry matter. Standard deviation ranged within 6% of mean values. Table 1 reports temperatures and dehydration times to reduce water content to 25% of the initial value and drying kinetics for both samples are shown in Fig. 1. Obviously, the dehydration time was inversely proportional to the treatment temperature. For each temperature the dehydration was quicker for treated cherry tomatoes according to Lewicki et al. (2002). Treated samples showed higher dry matter content, probably due to the presence of salts in the pre-treatment solution (sodium chloride, citric acid and calcium chloride). The water activity changed from 0.97 in the fresh tomatoes to 0.93 in semi-dry ones.

3.2. Colour measurements

Table 2 shows the CIE colour parameters. In particular the a^*/b^* ratio is often used as an indicator of colour development in tomatoes and redder hue, since it produces a good linear regression with

Table 1
Temperatures, partial dehydration times and dry matter content

Sample	Temperature (°C)	Time (h)	Dry matter (g/100 g)
Untreated	40	29	39.16 ± 0.21
	60	9	
	80	4	
Treated	40	24	42.00 ± 0.62
	60	8	
	80	3.5	




Fig. 1. Drying kinetics of untreated and treated cherry tomatoes.

the maturation of the fresh tomatoes (Arias, Lee, Logendra, & Janes, 2000). The results agree with literature data regarding a convection drying system at 95 °C, obtained by Shi et al. (1999).

Yeatman (1969) observed that the b^*L^*/a^* ratio gave a high linear correlation with the visual colour scores of processed tomatoes. The colour of treated semi-dry cherry tomatoes appeared more comparable with the colour characteristics of the fresh test.

Hue values range from 180° (pure green colour) to 0° (pure red colour). In fresh tomatoes, the hue is deeper than in treated tomatoes, according to previous work (Arias et al., 2000).

3.3. Carotenoids, ascorbic acid and 5-hydroxymethylfurfural

Table 3 shows the content of β -carotene, lycopene, ascorbic acid and HMF in the fresh tomato and in the untreated and treated semi-dry products. The fresh cherry tomato contains a high quantity of lycopene (99.8 mg/100 g d.m.) and β -carotene (38.0 mg/100 g d.m.). In untreated semi-dry samples the highest value of lycopene was found in the dehydrated sample at 80 °C (76.4 mg/100 g d.m.), while the lowest value was found at 40 °C (58.5 mg/100 g d.m.), confirming that in these samples lycopene was damaged by the length of the drying process.

β -Carotene in the untreated samples followed the same trend described above. Most protection was achieved at 80 °C (34.8 mg/100 g d.m.), while the highest loss was at 40 °C (28.6 mg/100 g d.m.). β -Carotene was found to be more stable in comparison with lycopene and it decreased by up to 25.9 mg/100 g d.m. in the untreated product dried at 60 °C.

In treated samples, lycopene and β -carotene contents were preserved at low temperature with long process time (40 °C for 24 h) while for other temperatures this effect was less apparent or lacking. The β -carotene was minimum after drying at 60 °C, with losses ranging from 28% to 32% in treated and untreated tomatoes, respectively.

Ascorbic acid loss was clearly temperature-dependent, as it increased with increasing dehydration temperature. The maximum losses were observed in samples dried at 80 °C, which showed a content of 112.7 and 178.5 mg/100 g d.m. in untreated and treated samples, respectively, highlighting the greater stability of ascorbic acid in the former samples, proving treatment was effective.

The formation of HMF in semi-dry cherry tomatoes is lower than that reported in the literature for fully dried tomatoes, as was expected (Zanoni, Peri, Nani, & Lavelli, 1999); the highest concentrations were found in samples dried at the highest temperature (1.8 and 1.7 mg/100 g d.m. for untreated and treated, respectively), while the lowest values were observed at 40 °C (1.1 and 1.2 mg/100 g d.m.).

A high tomato consumption plays a protective role in inhibiting the propagation of radical chain reactions due to the carotenoid constituents, particularly lycopene and β -carotene, and also a variety of natural antioxidants such as ascorbic acid and polyphenols (Lavelli & Giovanelli, 2003; Re, Bramley, & Rice-Evans, 2002). Food processing is generally believed to be responsible for a depletion of naturally occurring antioxidants. Carotenoids are light and temperature-sensitive, so high temperature, long processing time, light and oxygen have been shown to have effects on their degradation. Carotenoids and ascorbic acid decrease has been observed by Shi et al. (1999), Lavelli, Hippeli, Peri, and Elstner (1999) in the conventional air drying of tomatoes.

Pre-treatment effectiveness is strictly linked to dipping solution components, as reported by Baloch, Khan, and Baloch (1997). The lower decrease of nutritional components in the treated samples can be explained by the protective role of sodium chloride and citric acid. In fact the products undergo more rapid water loss in the presence of sodium chloride reducing the dehydration time, while citric acid provides secondary antioxidant activity.

Table 2
CIE colour parameters in fresh, untreated and in treated (dipping solutions of sodium chloride, citric acid and calcium chloride) semi-dry cherry tomatoes

Sample	T (°C)	L*	a*	b*	C	h	a*/b*	b*L*/a*
Fresh	–	22.33 ± 0.56	7.67 ± 0.19	20.53 ± 0.51	21.55 ± 0.54	74.01 ± 1.85	0.37 d	52.27 a
Untreated	40	33.75 ± 1.18	22.48 ± 0.79	20.04 ± 0.70	30.16 ± 1.06	41.85 ± 1.46	1.12 ab	30.08 e
	60	35.11 ± 1.05	22.80 ± 0.68	22.89 ± 0.69	32.44 ± 0.97	46.24 ± 1.39	1.00 abc	35.24 c
	80	36.98 ± 0.92	26.54 ± 0.93	22.12 ± 0.55	35.00 ± 1.58	40.12 ± 1.00	1.20 a	30.82 de
Treated	40	36.65 ± 1.03	18.92 ± 0.53	17.84 ± 0.50	26.35 ± 0.74	42.38 ± 1.19	1.06 abc	35.54 cd
	60	39.10 ± 1.64	19.60 ± 0.82	22.37 ± 0.94	29.98 ± 1.26	48.88 ± 2.05	0.88 c	44.63 b
	80	37.36 ± 1.57	21.99 ± 0.92	25.48 ± 1.07	33.75 ± 1.42	49.05 ± 2.45	0.90 bc	43.27 b

Means in columns followed by different letter are significantly different at $P \leq 0.05$.

Table 3
Mean contents of β -carotene, lycopene, ascorbic acid and 5-hydroxymethylfurfural

Sample	T (°C)	β -Carotene		Lycopene		Ascorbic acid		HMF	
		mg/Kg	mg/100 g d.m.	mg/Kg	mg/100 g d.m.	mg/Kg	mg/100 g d.m.	mg/Kg	mg/100 g d.m.
Fresh	–	31.0 ± 1.18	38.0 a	81.3 ± 2.28	99.8 a	353.3 ± 4.24	433.5 a	–	–
Untreated	40	112.7 ± 0.01	28.6 cd	230.5 ± 2.78	58.5 e	1226.1 ± 13.49	311.5 c	0.8 ± 0.03	1.12 c
	60	101.8 ± 0.22	25.9 e	260.9 ± 4.70	66.5 d	648.9 ± 16.22	165.5 f	1.0 ± 0.03	1.30 bc
	80	135.6 ± 0.14	34.8 b	297.7 ± 0.14	76.4 b	439.0 ± 5.27	112.7 g	1.3 ± 0.05	1.76 a
Treated	40	167.3 ± 0.00	38.5 a	290.8 ± 0.12	74.4 b	1509.8 ± 22.65	386.1 b	0.9 ± 0.02	1.15 c
	60	109.4 ± 0.20	27.4 cd	234.7 ± 0.29	58.7 e	692.0 ± 5.50	230.6 d	1.1 ± 0.02	1.41 abc
	80	118.1 ± 0.20	30.4 c	273.1 ± 0.83	70.4 c	688.0 ± 11.07	178.5 e	1.4 ± 0.03	1.72 ab

Means in columns followed by different letter are significantly different at $P \leq 0.05$.

* Values of treated samples were corrected considering the presence of salts (citric acid, sodium and calcium chloride) on dry matter content, supplied with the dipping solution ($42.00/39.16 = 1.07$).

HMF is commonly associated with sugar thermal damage. It is a thermal damage marker for processed foods and an index of improper storage conditions, its level being close to zero in fresh foods (Babsky, Toribio, & Lozano, 1986). The HMF level is a useful parameter for evaluating the technological history of food products, like honey, fruit juices, etc. The formation of HMF involves hexoses through acid-catalyzed dehydration and cyclisation. The first step of the reaction consists of a structural hexose change into an intermediate 1,2-endiolic form, which rapidly eliminates water (Belitz & Grosch, 1999). The formation of HMF through the dehydration and cyclisation of hexoses, besides low pH values and heat supply, is enhanced by low water activity values (Muratore, Licciardello, Restuccia, Puglisi, & Giudici, 2006); therefore, the reaction is stimulated at the later stages of dehydration. HMF is also a major product of the Maillard reaction; however, considering the small concentrations of free aminic groups compared to sugar concentration and low pH of tomato (Maillard reaction is favoured by high pH values), this pathway to HMF formation in dehydrated cherry tomatoes can be neglected.

The small quantities and little difference of HMF content in treated and untreated samples, indicate no intervention of the Maillard reaction. Therefore, according to the role of HMF as a marker of heat damage, the results qualify the semi-drying treatment of cherry tomatoes as a mild process.

4. Conclusions

The minimum browning effect of drying, evaluated either by the a^*/b^* or the b^*L^*/a^* ratios, was observed for treated samples dehydrated at 60 and 80 °C, while the highest colour degradation was caused by drying untreated samples. Therefore, the colour parameters highlighted the clearly positive effect of pre-treatment by dipping into a citric acid–sodium chloride–calcium chloride solution.

As regards carotenoids, after partial dehydration of the samples the loss of carotenoids did not prove to be related to time of treat-

ment or temperature. No clear dependence was found between the loss of these compounds and temperature and/or whether pre-treatment was performed, the worst results being at 60 °C.

Ascorbic acid was always higher in the treated tomatoes than in the untreated ones; the use of mild temperatures maintained these levels closer to those of fresh tomatoes, while at 80 °C the ascorbic acid loss reached 59% and 74% in treated and untreated samples.

The HMF content confirms the effect of temperature on its formation, but it is so low as to be considered negligible.

Tomatoes and their derivatives are important in the human diet for their carotenoid contribution, and represent the almost exclusive source of lycopene. After partial dehydration of the samples, the loss of carotenoids did not prove to be related to time of treatment or temperature. The above results, supported by statistical analysis, suggest that treatment with a dipping solution protected both the nutritional and chemical qualities of partially dried cherry tomatoes.

Acknowledgment

The authors gratefully acknowledge Filippo Tomaselli for advice on statistical analyses.

References

- Abushita, A. A., Hebshi, E. A., Daood, H. G., & Biacs, P. A. (1997). Determination of antioxidant vitamins in tomatoes. *Food Chemistry*, 60, 207–212.
- Arias, R., Lee, T. C., Logendra, L., & Janes, H. (2000). Correlation of lycopene measured by HPLC with the L^* , a^* , b^* color reading of a hydroponic tomato and the relationship of maturity with color and lycopene content. *Journal of Agricultural and Food Chemistry*, 48, 1697–1702.
- Babsky, N. E., Toribio, J. L., & Lozano, J. E. (1986). Influence of storage on the composition of clarified apple juice concentrate. *Journal of Food Science*, 51, 564–567.
- Baloch, W. A., Khan, S., & Baloch, A. K. (1997). Influence of chemical additives on the stability of dried tomato powder. *International Journal of Food Science and Technology*, 32(2), 117–120.
- Belitz, H.-D., & Grosch, W. (1999). In *Food chemistry* (2nd ed., pp. 252–261). Berlin, Germany: Springer-Verlag.

- Crozier, A., Lean, M. E.J., Mc Donald, M. S., & Black, C. (1997). Quantitative analysis of the flavonoid content of commercial tomatoes, onions, lettuce, and celery. *Journal of Agricultural and Food Chemistry*, *45*, 590–595.
- De Sio, F., Grimaldi, M., & Louidice, R. (1999). *Introduzione all'HPLC e Metodi Analitici per Alimenti Vegetali*. Milan, Italy: Morgan Edizioni Tecniche.
- Finley, J. W., & Duang, E. (1981). Resolution of ascorbic, dehydroascorbic and diketogulonic acids by paired-ion reserved phase chromatography. *Journal of Chromatography*, *207*, 449–453.
- Gartner, C., Stahl, W., & Sies, H. (1997). Lycopene is more bioavailable from tomato paste than from fresh tomatoes. *American Journal of Clinical Nutrition*, *66*, 116–122.
- Giovannucci, E. (1999). Tomatoes, tomato-based products, lycopene and cancer. Review of the epidemiologic literature. *Journal of the National Cancer Institute*, *91*, 317–331.
- Giovannucci, E., Ascherio, A., Rimm, E. B., Meir, J., Stampfer, M. J., Colditz, G. A., et al. (1995). Intake of carotenoids and retinol in relation to risk of prostate cancer. *Journal of the National Cancer Institute*, *87*, 1767–1776.
- La Vecchia, C. (1998). Mediterranean epidemiological evidence on tomatoes and the prevention of digestive tract cancers. *Proceedings of the Society for Experimental Biology and Medicine*, *218*, 125–128.
- Lavelli, V., & Giovanelli, G. (2003). Evaluation of heat and oxidative damage during storage of processed tomato products. II. Study of oxidative damage indices. *Journal of the Science of Food and Agriculture*, *83*, 966–971.
- Lavelli, V., Hippeli, S., Peri, C., & Elstner, E. (1999). Evaluation of radical scavenging activity of fresh and air-dried tomatoes by three model reactions. *Journal of Agricultural and Food Chemistry*, *47*, 3826–3831.
- Lewicki, P. P., Vu Le, H., & Pomarańska-Lazuka, W. (2002). Effect of pre-treatment on convective drying of tomatoes. *Journal of Food Engineering*, *54*, 141–146.
- Muratore, G., Licciardello, F., & Maccarone, E. (2005). Evaluation of the chemical quality of a new type of small-sized tomato cultivar, the plum tomato (*Lycopersicon lycopersicum*). *Italian Journal of Food Science*, *17*(1), 75–81.
- Muratore, G., Licciardello, F., Restuccia, C., Puglisi, M. L., & Giudici, P. (2006). Role of different factors affecting the formation of 5-hydroxymethyl-2-furancarboxaldehyde in heated grape must. *Journal of Agricultural and Food Chemistry*, *54*, 860–863.
- Nakagawa, T., Yokozawa, T., Teresawa, K., Shu, S., & Raj Juneja, L. (2002). Protective activity of green tea against free radical and glucose mediated protein damage. *Journal of Agricultural and Food Chemistry*, *50*, 2418–2422.
- Nisperos-Carriedo, M. O., Buslig, B. S., & Shaw, P. E. (1992). Simultaneous detection of dehydroascorbic, ascorbic, and some organic acids in fruits and vegetables by HPLC. *Journal of Agricultural and Food Chemistry*, *40*, 1127–1130.
- Re, R., Bramley, P. M., & Rice-Evans, C. (2002). Effects of food processing on flavonoids and lycopene status in a mediterranean tomato variety. *Free Radical Research*, *36*(7), 803–810.
- Russo, A., Acquaviva, R., Campisi, A., Sorrenti, V., Di Giacomo, C., Virgata, G., et al. (2000). Bioflavonoids as antiradicals, antioxidants and DNA cleavage protectors. *Cell Biology and Toxicology*, *16*, 91–98.
- Shi, J., Le Maguer, M., Kakuda, Y., Liptay, A., & Niekamp, F. (1999). Lycopene degradation and isomerization in tomato dehydration. *Food Research International*, *32*, 15–21.
- Sies, H., & Stahl, W. (1998). Lycopene: Antioxidant and biological effects and its bioavailability in the human. *Proceedings of the Society for Experimental Biology and Medicine*, *218*, 121–124.
- Stahl, W., & Sies, H. (1992). Uptake of lycopene and its geometrical isomers is greater from heat-processed than from unprocessed tomato juice in humans. *Journal of Nutrition*, *122*, 2161–2166.
- Yeatman, J. N. (1969). Tomato products: Read tomato red. *Food Technology*, *23*, 618–627.
- Zanoni, B., Peri, C., Nani, R., & Lavelli, V. (1999). Oxidative heat damage of tomato halves as affected by drying. *Food Research International*, *31*(5), 395–401.