



## Lipid and fatty acid compositions differentiate between wild and cultured Japanese eel (*Anguilla japonica*)

Takahiro Oku<sup>a</sup>, Akira Sugawara<sup>b</sup>, Malay Choudhury<sup>a</sup>, Masaharu Komatsu<sup>a</sup>, Shoji Yamada<sup>a</sup>, Seiichi Ando<sup>a,\*</sup>

<sup>a</sup> Department of Fisheries Science, Kagoshima University, 4-50-20 Shimoarata, Kagoshima 890-0056, Japan

<sup>b</sup> Hokkaido Central Fisheries Experimental Station, Yoichi, Hokkaido 046-8555, Japan

### ARTICLE INFO

#### Article history:

Received 11 July 2008

Received in revised form 8 December 2008

Accepted 9 December 2008

#### Keywords:

*Anguilla japonica*

Fatty acid composition

Japanese eel

Lipid profile

### ABSTRACT

The lipid profile and fatty acid composition of muscle, liver, and plasma lipoprotein were examined in wild and cultured Japanese eel (*Anguilla japonica*). Although, the muscle lipid levels of wild eels (11.6%) were less than those of cultured eels (13.1%), both eels were classified as typical fatty fish. Compared with the liver lipid composition of cultured eels, triacylglycerol (TG) level of the liver decreased in wild eels, whilst phosphatidylcholine and phosphatidylethanolamine levels of the liver increased in wild eels, reflecting the difference of liver lipid levels in both eels. Wild eels contained more cholesteryl ester (CE) and less TG, phospholipid, and free cholesterol in their plasma than cultured eels. The ratio of TG to CE decreased, whilst that of CE to total cholesterol increased in the plasma of wild eels. Different fatty acid compositions were found between wild and cultured eels. Compared with the fatty acid compositions of cultured eels, wild eels contained high percentages of 18:2, 18:3, and 20:4 and low percentages of 22:6, 20:1, and 20:5 in their muscle, liver, and plasma lipoprotein. The lipid profile and fatty acid composition seemed to be useful criteria for distinguishing between wild and cultured eels.

© 2008 Elsevier Ltd. All rights reserved.

### 1. Introduction

Fish preferentially utilise lipids rather than carbohydrates as an energy source and accumulate considerable amounts of lipids in their muscle, liver, or adipose tissue (Sheridan, 1988; Watanabe, 1982). Fish can be divided into fatty and non-fatty fish based on the lipid level of their muscle. Muscle is the main lipid reservoir in the former, whilst the latter possesses substantial quantities of lipids in the liver, but not muscle (Ando, Mori, Nakamura, & Sugawara, 1993; Sheridan, Allen, & Kerstetter, 1983; Shibata & Kayama, 1989).

Japanese eel *Anguilla japonica* is known to be a typical fatty fish (Ohshima, 1985; Ozaki, Koga, Takahashi, Adachi, & Yamauchi, 2008). Japanese eel is one of the most valuable cultured species in Japan. Some 100,000 and 300 tons of cultured and wild eels, respectively, are supplied to the food market of Japan a year (Tachiki, 1996). A commercial eel diet mainly consists of fish meal and starch and its fatty acid composition is rich in 16:0, 18:1, 22:6, and 20:5 (Ozaki et al., 2008; Suzuki, Okazaki, Hayakawa, Wada, & Tamura, 1986). Cultured Japanese eels contain more than 10% of lipids in their muscle. Most Japanese favour grilled cultured Japanese eels because they accumulate considerable amounts of lipids in their muscle.

In our previous papers, we have reported that cultured Japanese eels were in plasma hyperlipoproteinemia, reflecting their high use of lipids (Ando & Matsuzaki, 1996, 1997). Plasma lipoproteins serve as major carriers of lipids and other hydrophobic compounds. Lipoproteins with high levels of lipids, such as very low density (VLDL) and low density (LDL) lipoproteins, predominated in the plasma of cultured Japanese eels, in contrast to other teleosts with high density lipoprotein (HDL) as the main component. To our knowledge, cultured Japanese eels with high level of VLDL seem to be the only teleosts reported to date, although, plasma VLDL level has been shown to increase in rainbow trout *Oncorhynchus mykiss* treated with oestradiol-17 $\beta$  (Wallert & Babin, 1992). On the other hand, the average plasma lipoprotein levels of wild Japanese eels were lower than those of cultured eels (36.9 vs. 54.1 mg/ml of plasma), and both VLDL and LDL levels of wild Japanese eels were reduced to one-half of cultured ones (Ando, 1999; Ando & Matsuzaki, 1996). The lipoprotein profiles found in the plasma of wild Japanese eels were different from those of cultured eels. Both HDL and VLDL were the predominant lipoproteins in the plasma of wild Japanese eels. VLDL is synthesised from triacylglycerol (TG) and apolipoproteins in the liver and secreted as TG-rich lipoprotein in the blood. TG is hydrolysed by lipoprotein lipase during the circulation of VLDL. The hydrolytic products are absorbed in the peripheral tissues, including muscle and resynthesised into TG, the final storage product. Thus the difference of plasma lipoprotein profiles between wild and cultured Japanese eels seems to influence their

\* Corresponding author. Tel.: +81 99 286 4201; fax: +81 99 286 4015.

E-mail address: [ando@fish.kagoshima-u.ac.jp](mailto:ando@fish.kagoshima-u.ac.jp) (S. Ando).

lipid profiles in muscle and liver. It is of interest whether wild Japanese eels are classified into fatty fish similarly to cultured eels.

In this study, we examined the lipid profiles of muscle, liver, and plasma in wild and cultured Japanese eels. We also analysed fatty acid compositions of muscle, liver, and plasma lipoproteins from both eels. The results indicate that lipid profiles of liver and plasma, and fatty acid compositions of muscle, liver, and plasma lipoproteins enable us to differentiate between wild and cultured Japanese eels.

## 2. Materials and methods

### 2.1. Fish

Wild Japanese eels (average body weight 195 g,  $n = 3$ ) were captured from the Amori River, Kagoshima. Cultured Japanese eels (average body weight 220 g,  $n = 3$ ) starved for two days were purchased from a local fish market.

### 2.2. Proximate and lipid compositions of muscle and liver

The dorsal muscle and liver from wild and cultured Japanese eels were analysed in duplicate for moisture and protein according to the usual methods (AOAC, 1984). Lipid extraction from the dorsal muscle and liver was carried out in duplicate by the method of Bligh and Dyer (1959). The total lipid was analysed quantitatively by thin-layer chromatography (TLC). The TLC plates (Kieselgel 60, a ready-made plate from Merck) were developed using *n*-hexane:diethylether:acetic acid 85:15:1 (by volume) for non-phospholipid, and chloroform:methanol:acetic acid:water 25:15:4:2 (by volume) for phospholipid. The TLC plate was sprayed with 3% copper acetate-8% phosphoric acid, heated on a hot plate, and quantified using a Pharmacia Biotech Image Master. The amount of phospholipid was calculated from lipid-phosphorus assayed using a Phospholipid-Test Wako (Wako Pure Chemical Industries, Osaka, Japan).

### 2.3. Plasma lipoprotein isolation and plasma lipid analysis

Blood was drawn into ethylenediaminetetraacetic acid (EDTA, 5 mg/ml of blood) dissolved in 0.15 M NaCl, by using a syringe inserted in the caudal vasculature of live Japanese eels, and maintained at a temperature of 4 °C throughout the procedure. Plasma was obtained by centrifugation (3000 g for 15 min). VLDL (density ( $d$ ) < 1.006 g/ml), LDL (1.006 <  $d$  < 1.085 g/ml), HDL2 (1.085 <  $d$  < 1.100 g/ml), and HDL3 (1.100 <  $d$  < 1.210 g/ml) in the plasma were obtained by sequential ultracentrifugal flotation (Ando & Matsuzaki, 1996), using a Hitachi CS100 ultracentrifuge equipped with a RP80AT rotor (267,000 g at 15 °C). The plasma lipid composition was estimated using commercially available enzymatic kits from Kyowa Medex (Tokyo, Japan) for TG, phospholipid (PL), free (FC), and total cholesterol (TC). The amount of cholesteryl ester (CE) was calculated from the difference between TC and FC contents. Protein content was determined using Bio-Rad Protein Assay kit (Bio-Rad Laboratories, Hercules, CA) using bovine serum albumin as a standard.

### 2.4. Fatty acid analysis

A 1 ml aliquot of chloroform solution containing 100 mg of total lipid from muscle or liver was poured on a Sep-Pak silica cartridge (2.0 ml; Waters Chromatography Division, Milford, MA) for separating non-phospholipid and phospholipid fractions. Non-phospholipid and phospholipid fractions were eluted with 20 ml chloroform and 30 ml methanol, respectively. The total lipid was

extracted from plasma lipoprotein fractions according to the method of Bligh and Dyer (1959).

Fatty acid compositions of total lipid, non-phospholipid, and phospholipid were determined after methanolysis according to Prevot and Mordret (1976), and subsequent analysis on a gas chromatograph (Hitachi type 263-30). The conditions for gas chromatography were as follows. The separation column consisted of a 40 m wide-bore capillary column G-300, equivalent to polyethylene glycol 20 M (i.d. 1.2 mm) (Chemical Inspection Association, Tokyo). The oven temperature was 190 °C, and the injection port and detector temperature were 245 °C. Helium was used as a carrier gas and peak integration was carried out with a Hitachi D-2500 Chromato-Integrator. Fatty acids were identified by comparing their retention times with those of standards and reported equivalent chain length values.

### 2.5. Statistics

Data were analysed using one-way analysis of variance, followed by Student's *t*-test.

## 3. Results and discussion

The results of this study were obtained from a limited sample size ( $n = 3$  from each wild and cultured Japanese eels) and any statistical analyses of data seemed not to be valid. Nevertheless, some average values significantly differed between wild and cultured eels.

### 3.1. Proximate compositions of muscle and liver from wild and cultured Japanese eels

Table 1 shows the proximate compositions of muscle and liver from wild and cultured Japanese eels. The moisture contents of muscle from wild and cultured eels ranged from 67.3% to 71.1% and 66.1% to 68.2%, respectively. The protein contents of muscle from both eels were in the range of 18.3–19.4%. The lipid contents of muscle from wild and cultured eels ranged from 8.1% to 13.5% and 12.6% to 13.8%, respectively. The muscle lipid levels of wild eels seemed to be less than cultured eels, but both eels are fatty enough to be classified into typical fatty fish. In cases of American eel (*Anguilla rostrata*), the eels cultured with fish meal in earthen ponds for 18 months contained over 14.7% lipid in their muscle, whilst wild eels with similar body weight (166 g) to cultured eels averaged less than 3.4% lipid in their muscle (Ottwell & Rickards, 1981/1982). The difference of muscle lipid levels between wild Japanese and American eels may be influenced by feed composition.

Compared with the proximate compositions of muscle, the moisture contents of liver increased, whilst the lipid and protein contents of liver decreased (Table 1). The moisture contents of liver from wild and cultured eels ranged from 72.9% to 74.7% and 68.8% to 72.8%, respectively. The liver protein contents of wild eels were less than those of cultured eels, ranging 13.6% to 14.8% in the

**Table 1**  
Proximate compositions of muscle and liver from wild and cultured Japanese eels.

Component (g/100 g tissue)		Wild	Cultured
Muscle	Moisture	69.1 ± 1.9	67.4 ± 1.1
	Lipid	11.6 ± 3.1	13.1 ± 0.6
	Protein	19.0 ± 0.4	18.9 ± 0.5
Liver	Moisture	73.9 ± 1.0	70.8 ± 2.8
	Lipid	4.7 ± 1.1	7.6 ± 0.2
	Protein	14.2 ± 0.6	16.8 ± 3.3

former and 13.1% to 19.5% in the latter, respectively. The lipid contents of liver from wild and cultured eels ranged from 3.6% to 5.9% and 7.4% to 7.8%, respectively. The lipid contents of liver from cultured eels varied little and were higher than those of wild eel.

### 3.2. Lipid compositions of muscle, liver, and plasma from wild and cultured Japanese eels

The muscle and liver lipids consisted of non-phospholipid and phospholipid. Non-phospholipid included steryl ester, TG, free fatty acid, sterol, and partial glyceride, whilst phospholipid included phosphatidylethanolamine, phosphatidylserine, phosphatidylcholine, and sphingomyelin. A comprehensive analysis on fish tissues containing various amounts of lipids has revealed that phospholipid contents were nearly constant at 0.44–0.92% and 0.63–2.12% in the muscle and liver, respectively (Takama, Suzuki, Yoshida, Arai, & Anma, 1994; Takama, Suzuki, Yoshida, Arai, & Mitsui, 1999). Fig. 1 shows the relationship between total lipid, non-phospholipid, and phospholipid contents of muscle and liver from wild and cultured eels. Non-phospholipid contents of muscle from both eels increased with total lipid contents, whilst phospholipid contents of muscle were almost equal regardless of muscle total lipid contents. Similar relationships were found in the livers of wild and cultured eels, although, the livers contained more phospholipid than the muscles (1.86 g/100 g liver vs. 0.59 g/100 g muscle). Non-phospholipid comprised 95% of muscle total lipids from wild and cultured eels, and TG was the major lipid component of the muscles from both eels (Table 2). The lipid compositions of the liver differed between wild and cultured eels. Non-phospholipid comprised 55% and 78% of liver total lipids from wild and cultured eels, respectively. Compared with the liver lipid composition of cultured eels, TG level of the liver decreased in wild eels, whilst phosphatidylcholine and phosphatidylethanolamine levels of the liver increased in wild eels. The lipid profile in the liver, but not in the muscle, seemed to be useful for distinguishing between wild and cultured eels.

Table 3 summarises the protein and lipid contents of plasma from wild and cultured Japanese eels. The plasma lipid levels of both eels seemed not to be directly associated with their feeding behaviour since the plasma was prepared from the cultured eels starved for two days. Compared with the plasma profiles of cultured eels, protein and cholesteryl ester levels increased in wild eels, whilst TG, PL, and FC levels decreased in wild eels. The ratio of TG to CE decreased and CE to TC increased in the plasma from wild eels, reflecting their plasma lipid profiles. These ratios seemed to be useful criteria to distinguish between wild and cultured eels

**Table 2**

Lipid compositions of muscle and liver from wild and cultured Japanese eels.

Component (g/100 g lipid)	Muscle		Liver	
	Wild	Cultured	Wild	Cultured
Steryl ester	10.2 ± 2.2	9.5 ± 0.3	20.0 ± 2.7	20.9 ± 0.6
Triacylglycerol	68.2 ± 1.8	67.9 ± 1.6	16.8 ± 4.5	40.0 ± 0.1*
Free fatty acid	9.9 ± 1.0	11.2 ± 2.4	11.1 ± 1.4	9.4 ± 0.1
Sterol	4.5 ± 0.7	4.5 ± 0.5	5.4 ± 0.6	4.7 ± 0.2
Partial glyceride	2.2 ± 0.2	2.2 ± 0.1	2.2 ± 0.4	2.0 ± 0.2
Phosphatidylethanolamine	1.5 ± 0.2	1.2 ± 0.1	13.4 ± 2.7	6.5 ± 0.6
Phosphatidylserine	0.9 ± 0.2	0.8 ± 0.1	7.7 ± 0.6	3.2 ± 0.4
Phosphatidylcholine	2.4 ± 0.6	2.3 ± 0.3	20.0 ± 4.7	11.0 ± 0.9
Sphingomyelin	0.2 ± 0.1	0.4 ± 0.1	3.4 ± 0.6	1.2 ± 0.1*

\* Significant difference from wild eels at  $p < 0.05$ .

**Table 3**

Protein and lipid contents of plasma from wild and cultured Japanese eels.

Component (mg/ml plasma)	Wild	Cultured
Protein	54.2 ± 5.3	44.7 ± 4.0
Phospholipid	9.0 ± 0.9	13.8 ± 2.5*
Free cholesterol	2.1 ± 0.3	3.6 ± 0.9*
Triacylglycerol	7.9 ± 5.3	19.4 ± 9.0
Cholesteryl ester	8.4 ± 1.3	6.6 ± 0.4
TG/CE ratio <sup>A</sup>	1.2 ± 1.0	2.9 ± 1.4
CE/TC ratio <sup>B</sup>	0.68 ± 0.04	0.54 ± 0.06*

<sup>A</sup> The ratio of triacylglycerol to cholesteryl ester.


<sup>B</sup> The ratio of cholesteryl ester to total cholesterol.

\* Significant difference from wild eels at  $p < 0.05$ .

in a similar way to the lipid profile in the liver. The difference of plasma lipid profiles in wild and cultured eels affected their lipoprotein profiles. Both VLDL and LDL were the major carriers of TG in fish plasma and the low TG level in the plasma of wild eels seemed to be responsible for the low levels of these plasma lipoproteins (Ando, 1999).

### 3.3. Fatty acid compositions of muscle, liver, and plasma lipoprotein from wild and cultured Japanese eels

Table 4 shows fatty acid compositions of total lipid, non-phospholipid, and phospholipid in the muscle and liver from wild and cultured Japanese eels. The fatty acid compositions of muscle total lipid were similar to those of non-phospholipid, reflecting that non-phospholipid comprised 95% of muscle total lipid. The



**Fig. 1.** Relationship between total lipid, non-phospholipid, and phospholipid contents of muscle (A) and liver (B) from wild (open symbols) and cultured (closed symbols) Japanese eels.

**Table 4**

Fatty acid composition (weight %) in the lipid classes of muscle (A) and liver (B) from wild and cultured Japanese eels.

Fatty acid	Total lipid		Non-phospholipid		Phospholipid	
	Wild	Cultured	Wild	Cultured	Wild	Cultured
(A)						
14:0	3.5 ± 1.2	3.9 ± 0.2	3.4 ± 1.3	4.0 ± 0.3	0.5 ± 0.0	0.6 ± 0.1
16:0	18.1 ± 0.7	18.5 ± 1.4	17.5 ± 0.7	18.4 ± 1.1	17.3 ± 1.3	13.2 ± 1.2*
16:1	12.4 ± 0.8	8.3 ± 0.4*	12.3 ± 0.4	8.3 ± 0.6*	4.5 ± 0.2	2.3 ± 0.3*
18:0	2.4 ± 0.2	3.6 ± 0.5*	2.1 ± 0.1	3.4 ± 0.3*	6.9 ± 0.4	8.5 ± 1.0
18:1	35.5 ± 3.3	37.6 ± 2.7	35.3 ± 2.3	37.1 ± 2.3	20.4 ± 1.1	19.2 ± 0.6
18:2	5.8 ± 1.9	1.5 ± 1.0*	5.9 ± 2.0	1.6 ± 1.0*	4.2 ± 1.5	0.5 ± 0.3
18:3	3.3 ± 0.2	0.4 ± 0.0*	3.5 ± 0.4	0.5 ± 0.0*	1.5 ± 0.4	0.2 ± 0.1*
20:1	1.4 ± 0.7	5.9 ± 1.3*	1.6 ± 0.2	6.0 ± 1.4*	0.7 ± 0.2	2.7 ± 0.1*
20:2	0.6 ± 0.1	0.1 ± 0.0	0.6 ± 0.2	0.2 ± 0.1	0.4 ± 0.2	0.2 ± 0.1
20:4	2.1 ± 0.3	0.4 ± 0.0*	1.9 ± 0.5	0.4 ± 0.1*	9.8 ± 0.3	2.8 ± 0.3*
20:5	2.8 ± 0.4	4.0 ± 0.4*	2.8 ± 0.1	4.0 ± 0.3*	7.8 ± 1.5	8.4 ± 1.0
22:1	0.4 ± 0.5	2.3 ± 0.7	0.4 ± 0.0	2.3 ± 0.6	–	0.5 ± 0.1
22:5	2.4 ± 0.6	2.0 ± 0.2	2.5 ± 0.5	2.1 ± 0.2	4.6 ± 0.5	3.7 ± 0.3*
22:6	4.2 ± 2.7	7.2 ± 0.9	4.0 ± 2.5	6.7 ± 0.4	17.7 ± 5.4	34.2 ± 1.1*
(B)						
14:0	0.9 ± 0.2	2.1 ± 0.1	1.8 ± 0.4	2.6 ± 0.1	0.4 ± 0.0	0.5 ± 0.2
16:0	19.4 ± 1.6	17.5 ± 2.0	20.9 ± 0.4	17.7 ± 1.8	19.7 ± 2.7	18.1 ± 1.8
16:1	6.3 ± 2.6	6.3 ± 0.1	10.6 ± 0.8	7.2 ± 0.4*	3.6 ± 0.5	2.6 ± 0.3
18:0	6.8 ± 2.9	6.2 ± 1.6	3.0 ± 0.5	4.0 ± 1.3	8.6 ± 1.4	12.6 ± 2.1
18:1	21.7 ± 6.1	32.1 ± 1.6	37.1 ± 1.6	39.4 ± 1.6	13.5 ± 0.2	15.2 ± 0.2*
18:2	4.7 ± 0.6	1.3 ± 0.5*	4.7 ± 1.1	1.3 ± 0.7	4.0 ± 0.7	1.1 ± 0.6*
18:3	1.7 ± 0.3	0.4 ± 0.0*	1.9 ± 0.7	0.4 ± 0.1	1.4 ± 0.3	0.3 ± 0.1*
20:1	1.1 ± 0.2	3.9 ± 1.2	1.1 ± 0.2	4.3 ± 1.3	1.0 ± 0.2	1.8 ± 0.6
20:2	0.7 ± 0.0	0.4 ± 0.1	0.5 ± 0.2	0.2 ± 0.1	0.7 ± 0.1	0.5 ± 0.1
20:4	7.7 ± 0.5	1.4 ± 0.0*	3.8 ± 0.4	0.5 ± 0.1*	10.2 ± 1.4	4.2 ± 0.5*
20:5	4.3 ± 1.3	4.2 ± 0.0	1.7 ± 0.9	2.8 ± 0.2*	5.7 ± 0.4	8.6 ± 0.6*
22:1	–	0.8 ± 0.3	0.2 ± 0.0	1.0 ± 0.4	–	0.3 ± 0.0
22:5	2.7 ± 0.3	3.0 ± 0.5	2.1 ± 0.1	3.4 ± 0.7	3.1 ± 0.3	2.5 ± 0.4
22:6	16.8 ± 5.7	13.7 ± 0.3	7.6 ± 1.4	10.3 ± 0.1	22.4 ± 3.5	27.5 ± 2.4

\* Significant difference from wild eels at  $p < 0.05$ .

major fatty acids of total lipid and non-phospholipid were 18:1, 16:0, 16:1, 18:2, and 22:6 in the muscle of wild eels, whilst those of total lipid and non-phospholipid were 18:1, 16:0, 16:1, 22:6, and 20:1 in the muscle of cultured eels. Compared with the fatty acid compositions of total lipid and non-phospholipid in the muscle of cultured eels, wild eels contained high percentages of 16:1, 18:2, 18:3, and 20:4 and low percentages of 22:6, 20:1, 20:5, and 22:1. The fatty acid compositions of muscle phospholipid were different from those of total lipid and non-phospholipid. The major

fatty acids of phospholipid were 18:1, 22:6, 16:0, 20:4, and 20:5 in the muscle of wild eels, whilst those were 22:6, 18:1, 16:0, 18:0, and 20:5 in cultured eels. The phospholipid in the muscle of wild eels contained higher percentages of 16:0, 20:4, 16:1, and 18:2 and lower percentage of 22:6 than that of cultured eels.

The different fatty acid compositions of total lipid, non-phospholipid, and phospholipid were found in the livers from wild and cultured eels. The total lipid in the liver of wild eels contained higher percentages of 16:0, 22:6, 20:4, and 18:2 and lower percentages of 18:1 and 20:1 than that of cultured eels, although, the major fatty acids of total lipid were 18:1, 16:0, and 22:6 in the livers of both eels. The non-phospholipid contained low percentages of 22:6, 20:4, 18:0, and 20:5 and high percentages of 18:1 and 16:1 compared with the fatty acid composition of total lipid. The major fatty acids of liver phospholipid were similar to those of muscle phospholipid. The phospholipid in the liver of wild eels contained higher percentages of 20:4, 18:2, and 18:3 and lower percentages of 22:6, 18:0, and 20:5 than that of cultured eels.

The lipid contents of plasma lipoproteins from wild and cultured Japanese eels were far less than those of muscle and liver. This situation made it difficult to separate non-phospholipid and phospholipid fractions in plasma lipoproteins. As shown in Table 5, fatty acid compositions of total lipid were consequently determined for plasma lipoproteins of wild and cultured eels. The fatty acid compositions clearly differed in VLDL, LDL, HDL2, and HDL3. The percentages of 22:6, 20:4, and 20:5 increased with increasing the density of plasma lipoproteins, whilst those of 18:1 and 16:1 decreased with increasing the density of plasma lipoproteins. The main lipid components of VLDL and HDL were TG and PL, respectively. The proportion of PL increased with increasing the density of plasma lipoproteins and HDL3 was recognised as PL-rich lipoprotein (Ando, 1999; Ando & Matsuzaki, 1996). As shown in Table 4, phospholipid contained higher percentages of 22:6, 20:4, and 20:5 in HDL3 seemed to be derived from the PL of HDL3. The fatty acid compositions of plasma lipoproteins also differed between wild and cultured eels, similarly to those of muscle and liver. The wild eels contained high percentages of 16:1, 18:2, 20:4, and 18:3 and low percentages of 22:6, 18:0, 20:1, and 20:5 compared with the fatty acid compositions of plasma lipoproteins of cultured eels.

The dietary lipids have been recognised to modify the fatty acid composition of fish tissues, including muscle and liver (Bell et al.,

**Table 5**

Fatty acid composition (weight %) of plasma lipoproteins from wild and cultured Japanese eels.

Fatty acid	VLDL <sup>A</sup>		LDL <sup>B</sup>		HDL2 <sup>C</sup>		HDL3 <sup>D</sup>	
	Wild	Cultured	Wild	Cultured	Wild	Cultured	Wild	Cultured
14:0	1.7 ± 0.5	2.4 ± 0.1	1.2 ± 0.2	1.8 ± 0.2	0.9 ± 0.2	1.3 ± 0.1	0.7 ± 0.2	0.9 ± 0.1
16:0	20.3 ± 2.3	18.4 ± 1.7	24.6 ± 1.7	22.6 ± 1.6	24.9 ± 1.2	25.3 ± 2.2	24.8 ± 1.6	22.5 ± 1.7
16:1	10.7 ± 1.2	6.4 ± 0.7*	8.2 ± 0.3	4.6 ± 0.5*	6.2 ± 0.2	3.7 ± 0.7*	5.8 ± 0.6	3.0 ± 0.4*
18:0	3.1 ± 1.3	5.4 ± 1.3	4.3 ± 1.1	8.0 ± 1.0*	4.8 ± 1.4	9.2 ± 1.4*	4.9 ± 1.1	8.4 ± 1.4*
18:1	32.5 ± 2.8	35.3 ± 2.3	24.7 ± 0.4	26.8 ± 1.7	17.9 ± 0.6	21.4 ± 1.4*	16.2 ± 1.3	18.9 ± 0.6*
18:2	5.4 ± 1.1	0.8 ± 0.5*	4.8 ± 0.4	0.8 ± 0.2*	4.2 ± 0.6	0.7 ± 0.2*	4.2 ± 0.7	0.5 ± 0.6*
18:3	2.1 ± 0.6	0.2 ± 0.1*	1.4 ± 0.2	0.5 ± 0.2*	1.2 ± 0.2	0.4 ± 0.1*	1.2 ± 0.1	0.1 ± 0.1*
20:1	1.1 ± 0.1	5.5 ± 0.8*	0.9 ± 0.2	4.4 ± 0.6*	0.8 ± 0.3	2.9 ± 0.5*	0.7 ± 0.3	2.4 ± 0.5*
20:2	0.6 ± 0.2	0.2 ± 0.1	0.5 ± 0.1	0.3 ± 0.1	0.4 ± 0.2	0.2 ± 0.1	0.4 ± 0.1	0.2 ± 0.4
20:4	3.1 ± 0.7	0.6 ± 0.2*	5.9 ± 1.2	1.4 ± 0.4*	8.6 ± 1.3	1.7 ± 0.4*	9.4 ± 1.6	1.9 ± 0.2*
20:5	2.3 ± 0.2	4.0 ± 0.4*	3.9 ± 0.2	5.6 ± 0.8*	5.9 ± 0.4	7.3 ± 0.5*	6.4 ± 1.1	9.2 ± 1.1*
22:1	0.1 ± 0.0	2.4 ± 0.4*	–	1.7 ± 0.4	0.1 ± 0.0	1.0 ± 0.2	–	0.8 ± 0.2
22:5	2.9 ± 1.0	2.8 ± 0.2	2.7 ± 0.5	1.9 ± 0.4	3.2 ± 0.4	1.9 ± 0.2*	3.0 ± 0.2	2.8 ± 0.8
22:6	9.0 ± 2.8	11.0 ± 0.6	11.5 ± 2.4	13.0 ± 3.0	15.9 ± 2.3	17.3 ± 2.3	16.6 ± 1.6	23.0 ± 1.1*

<sup>A</sup> VLDL, very low density lipoprotein.<sup>B</sup> LDL, low density lipoprotein.<sup>C</sup> HDL2, high density lipoprotein 2.<sup>D</sup> HDL3, high density lipoprotein 3.\* Significant difference from wild eels at  $p < 0.05$ .



2002; Lie, Hemre, & Lambertsen, 1992; Mourente & Bell, 2006; Polvi & Ackman, 1992; Suzuki et al., 1986; Watanabe, 1982). A comparison of fatty acid composition between wild and cultured American eels has revealed that wild eels contained more 18:2, 18:3, and 20:4 and less 22:6 in their muscle than cultured eels, similar to this results (Otwell & Rickards, 1981/1982). Thus the fatty acid profiles of muscle, liver, and plasma lipoprotein were another criterion to distinguish between wild and cultured eels.

## References

- Ando, S. (1999). Occurrence of a very high density lipoprotein in the plasma from wild Japanese eel. *Fisheries Science*, *65*, 165–166.
- Ando, S., & Matsuzaki, M. (1996). A unique lipoprotein profile found in the plasma of cultured Japanese eel *Anguilla japonica*: Very low density lipoprotein, but not high density lipoprotein, is the main component of plasma. *Fish Physiology and Biochemistry*, *15*, 469–479.
- Ando, S., & Matsuzaki, M. (1997). Isolation of low density lipoprotein subfraction containing apolipoprotein B-like protein from Japanese eel *Anguilla japonica* plasma using dextran sulfate cellulose. *Comparative Biochemistry and Physiology*, *116B*, 191–196.
- Ando, S., Mori, Y., Nakamura, K., & Sugawara, A. (1993). Characteristics of lipid accumulation types in five species of fish. *Nippon Suisan Gakkaishi*, *59*, 1559–1564.
- AOAC (1984). Official methods of analysis. In *Association of official analytical chemists* (14th ed., pp.152–169). Arlington, VA.
- Bell, J. G., Henderson, R. J., Tocher, D. R., McGhee, F., Dick, J. R., Porter, A., et al. (2002). Substituting fish oil with crude palm oil in the diet of Atlantic salmon (*Salmo salar*) affects muscle fatty acid composition and hepatic fatty acid metabolism. *Journal of Nutrition*, *132*, 222–230.
- Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, *37*, 911–917.
- Lie, O., Hemre, G. I., & Lambertsen, G. (1992). Influence of dietary fatty acids on the glycerophospholipid composition in organs of cod (*Gadus morhua*). *Lipids*, *27*, 770–775.
- Mourente, G., & Bell, J. G. (2006). Partial replacement of dietary fish oil with blends of vegetable oils (rapeseed, linseed and palm oils) in diets for European sea bass (*Dicentrarchus labrax* L.) over a long term growth study: Effects on muscle and liver fatty acid composition and effectiveness of a fish oil finishing diet. *Comparative Biochemistry and Physiology*, *145B*, 389–399.
- Ohshima, T. (1985). Muscle lipids of cultured and wild fish. In M. Kayama (Ed.), *Muscle lipids of aquatic animals* (pp. 90–100). Tokyo: Koseisha-Koseikaku. in Japanese.
- Otwell, W. S., & Rickards, W. L. (1981/1982). Cultured and wild American eels, *Anguilla rostrata*: Fat content and fatty acid composition. *Aquaculture*, *26*, 67–76.
- Ozaki, Y., Koga, H., Takahashi, T., Adachi, S., & Yamauchi, K. (2008). Lipid content and fatty acid composition of muscle, liver, ovary and eggs of captive-reared and wild silver Japanese eel *Anguilla japonica* during artificial maturation. *Fisheries Science*, *74*, 362–371.
- Polvi, S. M., & Ackman, R. G. (1992). Atlantic salmon (*Salmo salar*) muscle lipids and their response to alternative dietary fatty acid sources. *Journal of Agricultural and Food Chemistry*, *40*, 1001–1007.
- Prevot, A. F., & Mordret, F. X. (1976). Utilisation des colonnes capillaires de verre pour l'analyse des corps gras par chromatographie en phase gazeuse. *Revue Française des Corps Grass*, *23*, 409–423.
- Sheridan, M. A. (1988). Lipid dynamics in fish: Aspects of absorption, transportation, deposition and mobilization. *Comparative Biochemistry and Physiology*, *90B*, 679–690.
- Sheridan, M. A., Allen, W. V., & Kerstetter, T. H. (1983). Seasonal variation in the lipid composition of steelhead trout, *Salmo gairdneri* Richardson, associated with parr-smolt transformation. *Journal of Fish Biology*, *23*, 125–134.
- Shibata, N., & Kayama, M. (1989). Lipids of some important fish in Japan. In M. Kayama (Ed.), *Comprehensive science of lipids* (pp. 831–844). Tokyo: Koseisha-Koseikaku. in Japanese.
- Suzuki, H., Okazaki, K., Hayakawa, S., Wada, S., & Tamura, S. (1986). Influence of commercial dietary fatty acids on polyunsaturated fatty acids of cultured freshwater fish and comparison with those of wild fish of the same species. *Journal of Agricultural and Food Chemistry*, *34*, 58–60.
- Tachiki, H. (1996). Sex control and induced maturation in the cultured eel. In O. Tabeta (Ed.), *Early Life history and prospects of seed production of the Japanese eel Anguilla japonica* (pp. 85–94). Tokyo: Koseisha-Koseikaku. in Japanese.
- Takama, K., Suzuki, T., Yoshida, K., Arai, H., & Anma, H. (1994). Lipid content and fatty acid composition of phospholipids in white-fleshed fish species. *Fisheries Science*, *60*, 177–184.
- Takama, K., Suzuki, T., Yoshida, K., Arai, H., & Mitsui, T. (1999). Phosphatidylcholine levels and their fatty acid compositions in teleost tissues and squid muscle. *Comparative Biochemistry and Physiology*, *124B*, 109–116.
- Wallert, C., & Babin, P. J. (1992). Effects of 17 $\beta$ -estradiol and starvation of trout plasma lipoproteins. *Lipids*, *27*, 1032–1041.
- Watanabe, T. (1982). Lipid nutrition in fish. *Comparative Biochemistry and Physiology*, *73B*, 3–15.