

# Ceramide-dependent Caspase 3 Activation is Prevented by Coenzyme Q from Plasma Membrane in Serum-deprived Cells

P. NAVAS<sup>a,\*</sup>, D.M. FERNANDEZ-AYALA<sup>a</sup>, S.F. MARTIN<sup>b</sup>, G. LOPEZ-LLUCH<sup>a</sup>, R. DE CABO<sup>a,c</sup>, J.C. RODRIGUEZ-AGUILERA<sup>a</sup> and J.M. VILLALBA<sup>b</sup>

<sup>a</sup>Laboratorio Andaluz de Biología, Universidad Pablo de Olavide, Carretera de Utrera, km 1, 41013 Sevilla, Spain; <sup>b</sup>Departamento de Biología Celular, Fisiología e Inmunología, Universidad de Córdoba, 14071 Córdoba, Spain; <sup>c</sup>Laboratory of Neurosciences, NIA/NIH, Baltimore, MD 21224, USA

Accepted by Professor H. Sies

(Received 30 April 2001; In revised form 10 October 2001)

Coenzyme Q (CoQ) is the key factor for the activity of the eukaryotic plasma membrane electron transport chain. Consequently, CoQ is essential in the cellular response against redox changes affecting this membrane. Serum withdrawal induces a mild oxidative stress, which produces lipid peroxidation in membranes. In fact, apoptosis induced by serum withdrawal can be prevented by several antioxidants including CoQ. Also, CoQ can maintain cell growth in serum-limiting conditions, whereas plasma membrane redox system (PMRS) inhibitors such as capsaicin, which compete with CoQ, inhibit cell growth and induce apoptosis. To understand how plasma membrane CoQ prevents oxidative stress-induced apoptosis we have studied the induction of apoptosis by serum withdrawal in CEM cells and its modulation by CoQ. Serum-withdrawal activates neutral sphingomyelinase (N-SMase), ceramide release and caspase-3-related proteases. CoQ addition to serum-free cultures inhibited a 60% N-SMase activation, an 80% ceramide release, and a 50% caspase-3 activity induced by serum deprivation. Caspase activation dependent on ceramide release since C<sub>2</sub>-ceramide was only able to mimic this effect in 10% foetal calf serum cultured cells but not in serum-free cultures. Also, *in vitro* experiments demonstrated that C<sub>2</sub>-ceramide and ceramide-rich lipid extracts directly activated caspase-3. Taken together, our results indicate that CoQ protects plasma membrane components and controls stress-mediated lipid signals by its participation in the PMRS.

**Keywords:** Coenzyme Q; Caspase; Plasma Membrane; Ceramide

## INTRODUCTION

Human plasma membrane contains significant amounts of CoQ<sub>10</sub>, which plays an essential role as antioxidant and electron carrier mediating the regeneration of other important antioxidants such as ascorbate and  $\alpha$ -tocopherol. These antioxidants are components of a plasma membrane-associated electron transport system (PMRS) which protects the membrane against damage caused by oxidative stress, where CoQ<sub>10</sub> is the central component.<sup>[1]</sup> Serum provides survival signals for cells and its depletion activates an intrinsic cell death program,<sup>[2,3]</sup> whose machinery is constitutively expressed.<sup>[4]</sup> There are evidences for an indirect generation of oxidative stress during apoptosis caused by growth factors withdrawal,<sup>[5]</sup> since this apoptotic program is inhibited by endogenous antioxidant enzymes<sup>[6,7]</sup> and other antioxidant molecules such as ascorbate,  $\alpha$ -tocopherol and CoQ<sub>10</sub>.<sup>[7–9]</sup> Furthermore, CoQ<sub>10</sub> is able to stimulate cell growth in the absence of serum,<sup>[10]</sup> and maintain growth in mitochondrial respiratory chain deficient cells.<sup>[9,11]</sup> In addition, CoQ<sub>10</sub> seems to be the key factor in preventing apoptosis induced by oxidative stress since antagonists of CoQ<sub>10</sub> trigger the apoptotic program.<sup>[12–14]</sup> The role of antioxidants in the prevention of apoptosis is keen in several

\*Corresponding author. Tel.: +34-95-434-9385. Fax: +34-95-434-9376. E-mail: pnavas@dex.upo.es.

systems, including regulation of antioxidant response mechanism by Bcl-2.<sup>[15–17]</sup>

Ceramide acts as a mediator of stress responses modulating enzymes of the eicosanoid pathway, protein kinases, nuclear factors, and gene expression.<sup>[18]</sup> Ceramide is also able to induce cell death after its intracellular accumulation,<sup>[9,19–22]</sup> by activating proteases of the caspase family.<sup>[23]</sup> Serum deprivation also induces a progressive increase of ceramide levels in Molt-4 and HL-60 cells.<sup>[9,20]</sup> Thus, the increase of ceramide levels appears to be an essential component in the stress response pathway triggered by serum withdrawal.<sup>[19]</sup>

We show here that CoQ<sub>10</sub> acts at the initiation phase of apoptosis by preventing the activation of neutral sphingomyelinase (N-SMase), consequently, decreasing the short-term ceramide accumulation and caspase-3 activation. Our results support the idea that CoQ<sub>10</sub> prevents apoptosis-induced by serum withdrawal through the inhibition of plasma membrane N-SMase and ceramide release.

## MATERIALS AND METHODS

### Cell Lines and Cultures

The human T-acute lymphatic leukaemia cell line CEM-C7H2 (CEM), a sub-line of CCRF-CEM, was grown in 5% CO<sub>2</sub> atmosphere at 37°C in RPMI-1640 medium (Sigma, Spain). Cultures were supplemented with 10% foetal calf serum (FCS) (PAA Labor, Austria), 100 U/ml penicillin, 100 µg/ml streptomycin, and 2.5 µg/ml amphotericin B (Sigma, Spain). For serum withdrawal experiments, cells were previously washed twice in serum free-RPMI 1640 medium and then seeded in tissue culture flasks in serum-free RPMI 1640 medium. CoQ<sub>10</sub> (Sigma, Spain) was dissolved in pure ethanol and added to serum-free culture media. Although no effect attributable to the vehicle was observed, 0.3% ethanol was added to control cultures.

### Apoptosis Determination

Cells were grown in 10% FCS supplemented RPMI medium. For treatments, cells were previously harvested, washed twice in serum-free medium and resuspended in 10% FCS-supplemented medium or in serum-free media supplemented or not with 30 µM CoQ<sub>10</sub>. After indicated times of incubation, cells were harvested, washed twice with ice cold PBS and fixed in 70% ethanol for at least 24 h at 4°C. After fixation, cells were washed with Hank's balanced salt solution (HBSS) and incubated for 30 min at 37°C with propidium iodide (PI) (Sigma, Spain) staining buffer [PBS pH 7.5, 0.1 mM EDTA, 0.1% Triton X-100, 50 U/ml DNase

free-RNase A (Sigma, Spain) and 50 µg/ml PI]. Cells were collected in a FACScan (Becton-Dickinson, USA) and data analysed using a Lysis II software (Becton-Dickinson, USA).

### Ceramide Quantification

Ceramide production was determined as previously described.<sup>[20]</sup> Briefly, cells were harvested at indicated times and washed twice with PBS. Lipids were then extracted by the method of Bligh and Dyer.<sup>[24]</sup> Lipids were dried and dissolved in 20 µl 1 mM diethylenetriamine pentaacetic acid, 7.5% octyl-β-D-glucopyranoside, and 5 mM cardiolipin (Sigma, Spain). After sonication for 15 s, 50 µl of 2 × reaction buffer (100 mM imidazole-HCl, pH 6.6, 100 mM NaCl; 25 mM MgCl<sub>2</sub> and 2 mM EGTA), DTT to a final concentration of 50 µM, 5 µg diacylglycerol-kinase (Sigma, Spain), and distilled water were added to a final volume of 90 µl. Reaction was started by the addition of a mixture of cold ATP and γ<sup>32</sup>P-ATP (specific radioactivity 100,000–500,000 cpm/nmol) (Amersham Iberica, Spain). After incubation for 20 min at 25°C, lipids were extracted with chloroform. Chloroform phase (500 µl) were dried with N<sub>2</sub> gas and dissolved in 100 µl of 5% methanol in chloroform. Ceramide-1-phosphate generated (25 µl) was resolved by TLC using chloroform/acetone/methanol/acetic acid/water (10:4:3:2:1) as solvent. Ceramide-phosphate spots were quantified using an automatic TLC-lineal analyser (Berthold, Germany). Ceramide was quantified using external standards and normalised to phosphate as indicator of total phospholipids.

### Assays for N-SMase and Caspase-3

Proteins (50–100 µg) from cell lysates, plasma membrane or purified N-SMase were mixed with assay buffer for N-SMase (50 mM Tris-HCl, pH 7.4; 0.5% Triton X-100; 5 mM DTT and 10 mM MgCl<sub>2</sub>). Then, 10 nmol of a mixture of cold sphingomyelin (SM) and [methyl-<sup>14</sup>C]-SM (specific radioactivity 10,000 cpm/nmol) (Amersham Iberica, Spain) were added. After incubation for 30–60 min at 37°C, the reaction was stopped by adding 1.5 ml chloroform/methanol (2:1) and 200 µl distilled water. Tubes were vortexed and then centrifuged at 1500g for 5 min to separate two phases. [<sup>14</sup>C]-phosphocholine present in the aqueous-phase was quantified using a liquid scintillation counter (Beckman, USA). SMase activity was expressed as nmol SM hydrolysed/mg protein/hour.

Caspase-3 activity was assayed by a colorimetric method using Ac-DEVD-pNa (Bachem, Switzerland) a specific substrate for caspase-3 and other related executioner caspases. The specific inhibitor Ac-DEVD-CHO (Bachem, Switzerland) was used to

TABLE I Effect of serum withdrawal on the ceramide-dependent pathway components of apoptosis and its inhibition by CoQ in CEM cells

Component	Growth conditions		
	+Serum	-Serum	-Serum + CoQ
Mg-dependent nSMase	60 ± 8	250 ± 21	105 ± 9*
Ceramide concentration	45 ± 6	89 ± 5	48 ± 5*
Caspase-3 activity	1100 ± 185	4200 ± 300	2000 ± 180*

N-SMase, ceramide levels and caspase-3 activity were determined in cells cultured in 10% FCS, serum-free and CoQ supplemented serum-free cultures. Cells were harvested after 1 h of incubation for both N-SMase and ceramide assays and after 2 h for caspase activity determination. Neutral SMase activity is expressed in cpm released/min/mg protein, ceramide concentration in pmol/nmol Pi, and caspase-3 activity in units (1 pmol pNa released/30 min/mg protein). \*Significant differences vs. serum-free conditions media,  $p \leq 0.05$  ( $n = 4$ ).

determine the specificity of the reaction. Protein (50 µg) were diluted in assay buffer (100 mM HEPES-KOH buffer, pH 7.5; 10% sucrose, 0.1% CHAPS, 0.1 mg/ml ovoalbumin, and 10 mM DTT). Activity was developed in 96 well plates at 37°C for 30–60 min, and paranitroaniline (pNa) released was determined spectrophotometrically at 405 nm using a multiwell reader. Activity was expressed in units (1 unit = 1 pmol pNa released/30 min/mg protein).

**Statistical Analysis**

Presented data represent the mean ± SD from three or four different experiments. Significant differences were determined using the one way ANOVA test.




FIGURE 1 Role of growth conditions on the activation of caspase-3 in CEM cells. Cells were incubated in the presence of serum or in serum free or 30 µM CoQ-supplemented serum-free media. C<sub>2</sub>-ceramide (50 µM) was also added to all the above-indicated conditions. After 2 h of incubation, cells were harvested and caspase activity determined. White bars: activity in the absence of C<sub>2</sub>-ceramide. Black bars: plus C<sub>2</sub>-ceramide. \*Significant differences vs. C<sub>2</sub>-ceramide free conditions ( $p \leq 0.01$ ).

**RESULTS**

CEM cells growing in a complete medium showed about 5% of apoptotic population at 48 h of culture but the culture in serum-free medium induced an increase of apoptotic cells reaching about 60% of the population at the same time. Addition of 30 µM CoQ<sub>10</sub> to serum-free cultures decreased apoptosis up to 35% of the population at the same time.

Different components of the ceramide-dependent apoptotic pathway were analysed in CEM cells and data are included in Table I. Two enzyme activities have been related to ceramide release in apoptotic events, N-SMase and acidic SMase (A-SMase). Latter activity was not affected in CEM cells after serum removal, but N-SMase activity increased four folds after the first hour of incubation in serum-free media. The presence of CoQ<sub>10</sub> in serum-free media clearly prevented N-SMase activation. Ceramide increased up to two folds in these cells after 1 h of culture in the absence of serum, but no differences in ceramide accumulation were observed when CoQ<sub>10</sub> was added to serum-free cultures. No changes in ceramide concentration occurred in 10% serum growing CEM cells at any time of incubation (data not shown). Further, caspase-3 proteases, able to recognise the aminoacidic sequence DEVD, were activated about four folds after 2 h of culture in CEM cells in the absence of serum, whereas the presence of CoQ decreased this activation by 50%.

TABLE II *In vitro* activation of caspase-3 by lipid extracts from CEM cells and C<sub>2</sub>-ceramide

Additions	Caspase-3 activity	Activity (%)
None	1164 ± 170	100
+CoQ 30 µM	1082 ± 110	93
10% FCS extract	1310 ± 264	113
0% FCS extract	2520 ± 350*	216*
0% FCS +CoQ extract	1480 ± 197	127
C <sub>2</sub> -ceramide 50 µM	3820 ± 540*	327*

Cytosol extracts from 10% FCS-cultured CEM cells were incubated *in vitro* with lipid extracts from 10% FCS, serum-free and 30 µM CoQ-supplemented serum-free cultured CEM cells. Controls were performed by incubation of cytosol extracts with ethanol, 30 µM CoQ and C<sub>2</sub>-ceramide (50 µM). Caspase-3 activity is expressed in units (1 pmol pNa released/30 min/mg protein). \*Significant differences vs. no additions,  $p \leq 0.05$  ( $n = 3$ ).

Free Radic Res Downloaded from informahealthcare.com by Library of Health Sci-Univ of Il on 11/24/11 For personal use only.




FIGURE 2 Inhibition of plasma membrane and purified N-SMase by reduced forms of CoQ<sub>0</sub> and CoQ<sub>10</sub>. (A) Plasma membrane from pig liver cells (60 µg) was incubated with vehicle, or 100 µM of CoQ<sub>10</sub>H<sub>2</sub> or CoQ<sub>0</sub>H<sub>2</sub> and N-SMase activity determined as indicated in "Material and methods" section. (B) Purified N-SMase (2 ng in 20 µl) from plasma membrane from pig liver cells was incubated in the presence or absence of 100 µM of CoQ<sub>10</sub>H<sub>2</sub> or CoQ<sub>0</sub>H<sub>2</sub>. Data represent the mean ± SD from three independent experiments. \*Significant differences vs. control, ( $p \leq 0.05$ ).

To determine if caspase-3 was present in the cytosol and ceramide was involved in its activation, C<sub>2</sub>-ceramide was added to CEM cells (Fig. 1). Clearly, caspase-3 was activated after the addition of C<sub>2</sub>-ceramide to cells growing in both the presence of serum and in the absence of serum plus CoQ<sub>10</sub>. However, C<sub>2</sub>-ceramide was unable to increase caspase-3 activity in cells growing in the absence of serum since these cells already reached the peak of activation of this caspase by incubation in serum-free media.

To determine if ceramide directly affected caspase-3 activity we performed an *in vitro* experiment by incubation of the cytosolic fraction of 10% FCS cultured CEM cells with C<sub>2</sub>-ceramide or lipid extracts from cells cultures in 10, 0% FCS or 0% FCS plus CoQ<sub>10</sub> supplemented-media (Table II). *In vitro* activation of cytosolic caspase-3 was significantly induced by lipid extracts from 0% FCS growing cells and by C<sub>2</sub>-ceramide but not by lipid extracts from both CEM cells growing in the presence of serum or in the absence of serum plus CoQ<sub>10</sub>. The effect of the lipid extracts from cells

grown in serum-free media was probably due to their higher concentration in ceramide. As control, CoQ<sub>10</sub> itself did not show any effect on *in vitro* cytosolic caspase-3 activation.

In order to study the mechanism of inhibition by CoQ<sub>10</sub> of ceramide-dependent apoptosis pathway, we analysed the effect of CoQ<sub>10</sub> on purified N-SMase from plasma membrane of pig liver hepatocytes (Fig. 2).<sup>[25]</sup> Plasma membranes purified from pig liver contain a N-SMase activity that was inhibited about 50% by CoQ<sub>10</sub>H<sub>2</sub>, but was not affected by the soluble form CoQ<sub>0</sub>H<sub>2</sub>. However, purified N-SMase was inhibited by CoQ<sub>0</sub>H<sub>2</sub> in a 60% but was not significantly affected by CoQ<sub>10</sub>H<sub>2</sub>. These results indicate lipid environment is important for the effective ubiquinone inhibition of N-SMase in plasma membrane.

## DISCUSSION

Data from the literature suggest that increased SMase activity and ceramide production are important factors in apoptosis induced by several agents including oxidative injury.<sup>[19,20,26]</sup> Ceramide accumulation induces cell death by activation of caspases.<sup>[23,27,28]</sup> In our study, caspase-3 was activated in CEM cells after ceramide release. The addition of C<sub>2</sub>-ceramide to cells increased caspase-3 activity to the same level in cultures grown with 10% FCS and in CoQ<sub>10</sub>-supplemented serum-free media. However, C<sub>2</sub>-ceramide was unable to further increase caspase activity in cells grown in serum-free media without CoQ<sub>10</sub>. These results suggest that CoQ<sub>10</sub>-dependent inhibition of ceramide release prevents caspase-3 activation. Exogenously added C<sub>2</sub>-ceramide restores ceramide levels for maximal activation of the caspases.

It is clear that growth factors or serum withdrawal induce a redox modulation of cellular activities.<sup>[5]</sup> Different antioxidants prevent both lipid hydroperoxides accumulation and apoptosis in serum-free conditions.<sup>[7-9,15]</sup> Then, lipid peroxidation seems to be a critical event in oxidative damage-dependent cell death and its prevention is one of the main functions of physiological antioxidants such as ascorbate, α-tocopherol and CoQ. Related to this apoptotic mechanism triggered by oxidative damage, phospholipase A<sub>2</sub> (PLA<sub>2</sub>) is activated by hydroperoxides in plasma membrane.<sup>[29]</sup> PLA<sub>2</sub> activation can be the responsible of subsequent activation of N-SMase initiating the ceramide-dependent apoptotic pathway.<sup>[30]</sup> In fact, inhibitors of PLA<sub>2</sub> delay cell death in other apoptotic systems linked to oxidative processes.<sup>[31]</sup> Thus, a putative relationship between PLA<sub>2</sub> and N-SMase activation is suggested. However, a direct activation of N-SMase by hydroperoxides can not be discarded

since intracellular antioxidants like glutathione are able to inhibit this lipase.<sup>[32]</sup> In our model, the antiapoptotic activity of CoQ<sub>10</sub> resides in the prevention of the lipid peroxidation chain reaction by itself or together with other antioxidants.<sup>[1,8,9]</sup>

Our results indicate that N-SMase localised in plasma membrane was inhibited by CoQ<sub>10</sub> but not by CoQ<sub>0</sub>, whereas the isolated enzyme was not affected by CoQ<sub>10</sub> but was inhibited by the soluble form CoQ<sub>0</sub>. These results clearly indicate a direct inhibition of N-SMase by ubiquinone and the importance of the lipid environment for the access of CoQ analogs to the enzyme.

Our data also suggest that ceramide is the factor responsible for the activation of caspase-3 related proteases after serum-deprivation. However, how ceramide activate these caspases remains controversial. It has been suggested that ceramide induce apoptosis through a mitochondrial factor, which is released once the mitochondrial transmembrane potential decreases and the mitochondrial pore complex is opened.<sup>[33]</sup> However, C<sub>2</sub>-ceramide is also able to induce nuclear apoptosis in isolated nuclei of CEM cells, and activate caspase-3 without the participation of mitochondria.<sup>[27,34]</sup> Our *in vitro* results confirm the caspase-3 activation in mitochondria-free fractions by C<sub>2</sub>-ceramide or ceramide-containing lipid extracts from serum-free grown cells. Thus, it seems that mitochondria are not absolutely required for ceramide activation of executioner caspases in our system.

In summary, in this model of oxidative injury induced by serum-withdrawal, CoQ may participate in the initiation phase of apoptosis preventing N-SMase activation, apparently by its direct inhibition, although an indirect effect through inhibition of hydroperoxide-induced PLA<sub>2</sub> activation can not be disregarded. In any case, plasma membrane CoQ<sub>10</sub> inhibits N-SMase activation and ceramide accumulation and thus, ceramide-mediated activation of executioner caspases, like caspase-3, independently of mitochondria. This model suggests that plasma membrane CoQ<sub>10</sub> is the first defensive level for the protection of cells against apoptosis induced by oxidative damage.

### Acknowledgements

This work has been partially supported by the Spanish D.G.E.S. grant number PB98-0329-C02-01.

### References

- [1] Navarro, F., Navas, P., Burgess, J.R., Bello, R.I., De Cabo, R., Arroyo, A. and Villalba, J.M. (1998) "Vitamin E and selenium deficiency induces expression of the ubiquinone-dependent antioxidant system at the plasma membrane", *FASEB J.* **12**, 1665–1673.
- [2] Raff, M.C. (1992) "Social controls on cell survival and cell death", *Nature* **356**, 397–400.
- [3] Raff, M.C., Barres, B.A., Burne, J.F., Coles, H.S., Ishizaki, Y. and Jacobson, M.D. (1993) "Programmed cell death and the control of the survival: lessons from the nervous system", *Science* **262**, 695–700.
- [4] Weil, M., Jacobson, M.D., Coles, H.S.R., Davies, T.J., Gradner, R.L., Raff, K.D. and Raff, M.C. (1996) "Constitutive expression of the machinery for programmed cell death", *J. Cell Biol.* **133**, 1053–1059.
- [5] Slater, A.F.G., Stefan, C., Nobel, I., Van der Dobbelen, D.J. and Orrenius, S. (1996) "Intracellular redox changes during apoptosis", *Cell Death Differ.* **3**, 57–62.
- [6] Greelund, L.J.S., Beckwert, T.L. and Johnson, E.M.J. (1995) "Superoxide dismutase delays neuronal apoptosis: a role for reactive oxygen species in programmed neuronal death", *Neuron* **14**, 303–314.
- [7] Tilly, J.L. and Tilly, K.I. (1995) "Inhibitors of oxidative stress mimic the ability of follicle-stimulating hormone to suppress apoptosis in cultured rat ovarian follicles", *Endocrinology* **136**, 242–252.
- [8] Barroso, M.P., Gómez-Díaz, C., López-Lluch, G., Malagón, M.M., Crane, F.L. and Navas, P. (1997) "Ascorbate and  $\alpha$ -tocopherol prevent apoptosis induced by serum removal independent of Bcl-2", *Arch. Biochem. Biophys.* **343**, 243–248.
- [9] Barroso, M.P., Gómez-Díaz, C., Villalba, J.M., Burón, M.I., López-Lluch, G. and Navas, P. (1997) "Plasma membrane ubiquinone controls ceramide production and prevents cell death induced by serum withdrawal", *J. Bioenerg. Biomemb.* **29**, 259–267.
- [10] Sun, I.L., Sun, E.E. and Crane, F.L. (1995) "Comparison of growth stimulation of HeLa cells, HL-60 cells, and mouse fibroblasts by coenzyme Q<sub>10</sub>", *Protoplasma* **184**, 214–219.
- [11] Martinus, R.D., Linnane, A.W. and Nagley, P. (1993) "Growth of human mammary cells lacking oxidative phosphorylation can be sustained by redox compounds ferricyanide or coenzyme Q putatively acting through the plasma membrane oxidase", *Biochem. Mol. Biol. Int.* **31**, 997–1005.
- [12] Wolvetang, E.J., Larm, J.A., Moutsoulas, P. and Lawen, A. (1996) "Effectors of the mammalian plasma membrane NADH-oxidoreductase system. Short-chain ubiquinone analogues as potent stimulators", *Cell Growth Differ.* **7**, 1315–1325.
- [13] Macho, A., Blázquez, M.V., Navas, P. and Muñoz, E. (1998) "Induction of apoptosis by vanilloid compounds does not require de novo gene transcription and activator protein 1 activity", *Cell Growth Differ.* **9**, 277–286.
- [14] Macho, A., Calzado, M.A., Muñoz-Blanco, J., Gómez-Díaz, C., Gajate, C., Mollinedo, F., Navas, P. and Muñoz, E. (1999) "Selective induction of apoptosis by capsaicin in transformed cells: the role of reactive oxygen species and calcium", *Cell Death Differ.* **6**, 155–165.
- [15] Hockenbery, D.M., Oltvai, Z.N., Yin, X.-M., Millman, C.L. and Korsmeyer, S.J. (1993) "Bcl-2 functions in an antioxidant pathway to prevent apoptosis", *Cell* **75**, 241–251.
- [16] Kane, D.J., Sarafin, T.A., Anton, R., Hahn, H., Gralla, E.B., Valentine, J.S., Ord, T. and Bredesen, D.E. (1993) "Bcl-2 inhibition of neuronal death: decreased generation of reactive oxygen species", *Science* **262**, 1274–1277.
- [17] Bruce-Keller, A.J., Begley, J.G., Fu, W., Butterfield, D.A., Bredesen, D.E., Hutchins, J.B., Hensley, K. and Mattson, M.P. (1998) "Bcl-2 protects isolated plasma and mitochondrial membranes against lipid peroxidation induced by hydrogen peroxide and amyloid-peptide", *J. Neurochem.* **70**, 31–39.
- [18] Hannun, Y.A. (1994) "The sphingomyelin cycle and the second messenger function of ceramide", *J. Biol. Chem.* **269**, 3125–3128.
- [19] Hannun, Y.A. (1996) "Functions of ceramide in coordinating cellular responses to stress", *Science* **274**, 1855–1859.
- [20] Jayadev, S., Liu, B., Bielawska, A.E., Lee, J.Y., Nazaire, F., Pushkareva, M.Y., Obeid, L. and Hannun, Y.A. (1995) "Role for ceramide in cell cycle arrest", *J. Biol. Chem.* **270**, 2047–2052.
- [21] Obeid, L.M., Linares, C.M., Karolak, L.A. and Hannun, Y.A. (1993) "Programmed cell death induced by ceramide", *Science* **259**, 1769–1771.
- [22] Jarvis, W.D., Kolesnick, R.N., Fornari, F.A., Traylor, R.S., Gewirtz, D.A. and Grant, S. (1994) "Induction of apoptotic

- DNA damage and cell death by activation of the sphingomyelin pathway", *Proc. Natl Acad. Sci. USA* **91**, 73–77.
- [23] Mizushima, N., Koike, R., Kohsaka, H., Kushi, Y., Handa, S., Yagita, H. and Miyasaka, N. (1996) "Ceramide induces apoptosis via CPP32 activation", *FEBS Lett.* **395**, 267–271.
- [24] Bligh, E. and Dyer, W. (1959) "A rapid method for total lipid extraction and purification", *Can. J. Biochem. Physiol.* **37**, 911–917.
- [25] Martin, S.F., Navarro, F., Forthoffer, N., Navas, P. and Villalba, J.M. (2001) "Neutral magnesium-dependent sphingomyelinase from liver plasma membrane: purification and inhibition by ubiquinone", *J. Bioenerg. Biomembr.*, **33**, 143–153.
- [26] Kim, M.-Y., Linardic, C., Obeid, L. and Hannun, Y.A. (1991) "Identification of sphingomyelin turnover as an effector for the action of tumor necrosis factor alpha and gamma-interferon. Specific role in cell differentiation", *J. Biol. Chem.* **266**, 484–489.
- [27] Martin, J., Newmeyer, D.D., Mathias, S., Farschon, D.M., Wang, H.-G., Reed, J.C., Kolesnik, R.N. and Green, D.R. (1995) "Cell-free reconstitution of Fas-UV radiation- and ceramide-induced apoptosis", *EMBO J.* **14**, 5191–5200.
- [28] Smyth, M.J., Perry, D.K., Zhang, J.D., Poirier, G.G., Hannun, Y.A. and Obeid, L.M. (1996) "Prl: a downstream target for ceramide-induced apoptosis and for the inhibitory action of Bcl-2", *Biochem. J.* **316**, 25–28.
- [29] Salgo, M.G., Corongiu, F.P. and Sevanian, A. (1993) "Enhanced interfacial catalysis and hydrolytic specificity of phospholipase A<sub>2</sub> toward peroxidized phosphatidylcholine vesicles", *Arch. Biochem. Biophys.* **304**, 123–132.
- [30] Jayadev, S., Hayter, H.L., Andrieu, N., Gamard, C.J., Liu, B., Balu, R., Hayakawa, M., Ito, F. and Hannun, Y.A. (1997) "Phospholipase A<sub>2</sub> is necessary for tumor necrosis factor alpha-induced ceramide generation in L929 cells", *J. Biol. Chem.* **272**, 17196–17203.
- [31] Neale, M.L., Fiera, A. and Matthews, N. (1988) "Involvement of phospholipase A<sub>2</sub> activation in tumor cell killing by tumour necrosis factor", *Immunology* **64**, 81–85.
- [32] Liu, B. and Hannun, Y.A. (1997) "Inhibition of the neutral magnesium-dependent sphingomyelinase by glutathione", *J. Biol. Chem.* **272**, 16281–16287.
- [33] Susin, S.A., Zamzami, N., Castedo, M., Daugas, E., Wang, H.-G., Gelei, S., Fassy, F., Reed, J.C. and Kroemer, G. (1997) "The central executioner of apoptosis: multiple connections between protease activation and mitochondria in Fas/APO-1/CD95- and ceramide-induced apoptosis", *J. Exp. Med.* **186**, 25–37.
- [34] Newmeyer, D.D., Farschon, D.M. and Reed, J.C. (1994) "Cell-free apoptosis in *Xenopus* eggs extracts: inhibition by Bcl-2 and requirement for an organelle fraction enriched in mitochondria", *Cell* **79**, 353–364.