

Green photochemistry: photo-Friedel–Crafts acylations of 1,4-naphthoquinone in room temperature ionic liquids

Brian Murphy,^a Peter Goodrich,^b Christopher Hardacre^b and Michael Oelgemöller^{*a,c}

Received 6th July 2009, Accepted 4th August 2009

First published as an Advance Article on the web 7th September 2009

DOI: 10.1039/b913252j

The photo-Friedel–Crafts acylation of 1,4-naphthoquinone with various aldehydes was investigated in a series of room temperature ionic liquids. High conversions and selectivities were achieved in [C₂mim]⁺-based ionic liquids with the highest isolated yields found in [C₂mim][NTf₂]. The developed procedure allowed for a replacement of hazardous solvents such as benzene and acetonitrile which are commonly used for this transformation.


Introduction

Ionic liquids (ILs) have been intensively studied over the last decade as non-flammable and replacement solvents for molecular systems.¹ The ease of product removal allows an efficient reuse of the ILs and this property, coupled with the lack of vapour pressure, has the potential to result in materials which are much less environmentally intrusive than conventional organic solvents. However, recent studies on the toxicity of ionic liquids have raised some concerns regarding their broader use.² In addition to their physical properties, ionic liquids can stabilise a wide range of highly reactive species and through the ionic solvent–solute interactions can direct the chemistry.³ For example, ionic liquids have been shown to stabilize radicals and radical ions in solution more efficiently than many conventional organic solvents.^{4,5} This coupled with their weak or missing absorption of light above 300 nm makes them an ideal solvent for photochemical reactions.⁶ Despite these advantages, photochemical transformations in ionic liquids are rare.⁷ This is surprising since photochemistry itself is regarded as a ‘green technology’,⁸ especially if conducted with sunlight.⁹ However, photochemical transformations conventionally use solvents that are toxic and/or flammable.

Over the last years we have intensively studied the photochemical acylation of 1,4-quinones with aldehydes as a mild and atom efficient alternative to Friedel–Crafts acylations or (photo-)Fries-rearrangements.^{10,11} The original procedure developed by Kraus and Kirihara involved the use of hazardous benzene or acetonitrile as solvents.¹² Supercritical carbon dioxide (SC-CO₂) has recently been reported as an alternative medium,¹³ but this technique suffers from technical limitations, *e.g.* high pressure or small reactor volume, especially for a large-scale application. We have, therefore, investigated the photochemical

acylation of 1,4-naphthoquinone in benzene and various room temperature ionic liquids, the latter either commercially available or readily synthesised by standard procedures.¹

The photoacylation of 1,4-naphthoquinone (**1**) with butyraldehyde (**2c**) was initially chosen as a model reaction to determine the optimum cation/anion combination for the ionic liquid (Scheme 1). A mixture of 1,4-naphthoquinone and excess of butyraldehyde dissolved in the IL was irradiated for 16 h while stirring. After work-up the crude product was analyzed by ¹H-NMR spectroscopy.¹⁴ The crude mixture generally consisted of unreacted 1,4-naphthoquinone (**1**), the desired photoacylation product (**3c**) and the reduction product 1,4-dihydroxynaphthalene (**4**). The ratio of these components critically depended on the structure of the ionic liquid employed (Table 1).



Scheme 1 Photoacylation of 1,4-naphthoquinone with butyraldehyde.

In many of the chosen ionic liquids the photoacylation product **3c** was formed in larger amounts than in the conventional solvent benzene. The best result was achieved in [C₂mim][OTf] and **3c** was obtained in 91% yield without any photoreduction product (**4**) present. The photoreaction in [C₂mim][NTf₂] also resulted in a high yield of **3c** at 81%; however, in this case, 1% of the photoreduction product (**4**) was formed as well. In both cases, the desired product **3c** was isolated by column chromatography.

For [C₂mim][OTf], the isolated yield of **3c** was low with only 10% obtained following work-up. This low yield may be associated with the strong hydrogen bonding interaction of the substrate with the ionic liquid anion. On changing the anion from [OTf]⁻ to [NTf₂]⁻ the isolated yield of **3c** improved to 40% (49% based on the amount of **1** consumed). This is still only modest and reflects the strong affinity of aromatics and carbonyl

^aSchool of Chemical Sciences and NCSR, Dublin City University, Glasnevin, Dublin 9, Ireland

^bSchool of Chemistry and Chemical Engineering/QUILL, Queen's University, Belfast, Northern Ireland, UK BT9 5AG

^cSchool of Pharmacy and Molecular Sciences, James Cook University, Townsville, Queensland, 4811, Australia.
E-mail: oelgemoller@jcu.edu.au; Fax: +61-7-4781-6078;
Tel: +61-7-4781-4535

Table 1 Product compositions for photoacylations of **1** with **2c**


Solvent ^a	Composition [%] ^b		
	1	3c	4
Benzene	49	44 (23 ^c /45 ^d)	7
[C ₂ mim][OTf]	9	91 (10 ^c /11 ^d)	—
[C ₂ mim][NTf ₂]	18	81 (40 ^c /49 ^d)	1
[C ₂ mim][BF ₄]	58	42	—
[C ₄ mim][NTf ₂]	29	25	46
[C ₆ mim][PF ₆]	44	13	43
[C ₈ mim][PF ₆]	39	10	51
[C ₄ Py][NTf ₂]	24	75	1
[C ₄ mPyrr][NTf ₂]	12	83	5

^a [C_nmim]⁺ = 1-alkyl-3-methylimidazolium, [C₄Py]⁺ = 1-butyl-pyridinium, [C₄mPyrr]⁺ = 1-butyl-1-methylpyrrolidinium, [NTf₂]⁻ = bis-(trifluoromethyl)sulfonyl imide, [OTf]⁻ = trifluoromethane sulfonate.

^b As determined by integration of characteristic peaks in the ¹H-NMR of the crude product. ^c Isolated yield. ^d Isolated yield based on conversion.

species in ionic liquids. Similarly poor isolation was found for other Friedel–Crafts processes.¹⁵ On extending the alkyl length of the imidazolium side-chain, photoreduction to **4** increased with the largest yield of **4** (51%) found in [C₈mim][PF₆]. The influence of the anion on the photoacylation was studied using [C₂mim]⁺-derived ionic liquids. The chemoselectivity remained high in all three cases studied; however, the conversion was significantly reduced in [C₂mim][BF₄]. This finding may be associated with the higher viscosity of the [BF₄]⁻ based ionic liquid.^{15,16} The higher viscosity, and thus the reduced diffusion, hinders the necessary approach of the two reactants during the irradiation. It was also noticed that the [PF₆]⁻ based ionic liquid was less favourable for the photoreaction and large amounts of the reduction product **4** were produced.

The photoacylation protocol was further applied to a range of aldehydes (Scheme 2). Due to the high selectivity obtained for the **1/2c** model pair in combination with the good isolated yield of **3c**, [C₂mim][NTf₂] was selected as the ionic liquid for this study. For comparison, selected transformations were also studied in [C₂mim][OTf].

**Scheme 2** Photoreaction of 1,4-naphthoquinone with various aldehydes.

The conversion rate was lower for long chain aldehydes (Table 2). With increasing chain-length of aliphatic aldehydes, a general increase in the photoreduction product **4** was also noticed.† This tendency may be best explained by the reduced solubility of long-chained aldehydes in the selected ionic liquids hence favouring hydrogen abstraction from the ionic liquid instead. A similar dependency of the outcome of a photoreaction

† In contrast, photoacylations in benzene give similar yields independent from the aldehyde.¹¹

Table 2 Product compositions for photoacylations of **1** with various aldehydes **2**

Entry	R	Composition [%] ^a					
		[C ₂ mim][OTf]			[C ₂ mim][NTf ₂]		
		1	3	4	1	3	4
a	CH ₃	— ^b	—	—	25	70	5
b	C ₂ H ₅	55	45	—	26	74	—
c	C ₃ H ₇	9	91	—	18	81	1
d	C ₄ H ₉	27	73	—	37	48	15
e	C ₅ H ₁₁	— ^b	—	—	26	66	8
f	C ₆ H ₁₃	30	70	—	35	47	18
g	C ₇ H ₁₅	50	50	—	20	70	10
h	C ₈ H ₁₇	39	56	5	69	23	8
i	C ₁₀ H ₂₁	62	32	6	36	51	13
j	C ₁₁ H ₂₃	— ^b	—	—	40	41	19
k	Ph	50	50	—	38	52	10
l	MeCH=CH	38	52	10	46	43	11


^a As determined by integration of characteristic peaks in the ¹H-NMR spectrum of the crude product. ^b Not studied.

on the solubility of the starting material in an ionic liquid has been reported for the attempted triplet-sensitised photolysis of myrcene.¹⁷ Likewise, benzaldehyde and crotonaldehyde gave the desired acylated hydroquinones **3k** and **l** in moderate yields and selectivities.

For recycling purposes, the purity of the ILs was investigated by NMR spectroscopy after irradiation and work-up.‡ Although the ¹H-NMR spectra of the recovered ILs did not reveal any significant changes, a strong yellow or brown colour often remained. Decolourisation was easily achieved by diluting the ionic liquid with methanol and stirring with activated charcoal. Subsequent filtration and removal of methanol gave almost entirely colourless ionic liquids. Experiments performed with recycled solvent showed no discernible loss in efficiency even after 3 cycles.

The mechanism of the photoacylation has been extensively studied by Bruce and Maruyama and both *in-cage* and *out-of-cage* scenarios have been described. Both mechanisms operate essentially simultaneously depending on the specific reaction conditions of the irradiation experiment, *i.e.* temperature, solvent, quinone/aldehyde applied.^{10a} The photoaddition is initiated by hydrogen-abstraction from the aldehyde function to the triplet excited quinone (Scheme 3, path **A**). The higher viscosity of ionic liquids compared with conventional solvents may prevent separation of the intermediary radical pair by diffusion, thus favouring the *in-cage* mechanism. A similar assumption was also made for the photoreduction of benzophenones in ionic liquids.¹⁸ Addition of the acyl radical to the delocalised semiquinone radical and subsequent tautomerisation yields the acylated hydroquinones **3**. For long alkyl-chain containing ionic liquids, which are highly viscous, hydrogen abstraction from the ionic liquid can operate the alternative pathway (path **B**) thus yielding the photoreduction product **4**. It is known that the triplet excited states of some molecules are capable

‡ For some common organic solvents, solvent recovery can be environmentally and economically harmful because of the large cumulative energy demand (CED) for the process.²² However, the relatively high price for common ionic liquids may justify recovery and reuse.



Scheme 3 Mechanistic scenario for the photoacylation and -reduction.

of abstracting hydrogen atoms from the side-chains of ionic liquids.¹⁹ The improved efficiency, *i.e.* higher conversion rates, of the photoacylation process in most ionic liquids may be best explained by an enhanced lifetime of the triplet excited state of the quinone as, for example, known for xanthone.⁴

In conclusion, the photoacylation of 1,4-naphthoquinone with a series of aldehydes has been carried out in ionic liquids. High conversions and chemoselectivities have been achieved in [C₂mim][OTf] and [C₂mim][NTf₂], respectively. The isolated yield of model compound **3c** was, however, better in [C₂mim][NTf₂]. The ionic liquids replaced previously used toxic solvents (benzene and acetonitrile) and, in comparison, showed increased activity and selectivity. In addition, the ionic liquid could be recovered and reused. Hence, this transformation can be regarded as a model reaction for 'Green Photochemistry'.²⁰ The simple protocol is currently being transferred to 'micro-photochemistry', *i.e.* photochemical transformations in micro-structured devices.²¹

Acknowledgements

This research project was financially supported by Dublin City University (Research Career Start Award 2006), Science Foundation Ireland (SFI, 07/RFP/CHEF817 and 06/RFP/CHO028), QUILL and a Portfolio Partnership grant from the EPSRC. BM thanks the EU Marie Curie Early Stage Training Site Fellowship programme for support, contract number HPMT-GH-00-00147-03.

References

- (a) T. Welton, *Chem. Rev.*, 1999, **99**, 2071; (b) J. Dupont, R. F. de Souza and P. A. Z. Suarez, *Chem. Rev.*, 2002, **102**, 3667; (c) N. Jain, A. Kumar, S. Chauhan and S. M. S. Chauhan, *Tetrahedron*, 2005, **61**, 1015; (d) N. V. Plechkova and K. R. Seddon, *Chem. Soc. Rev.*, 2008, **37**, 123.
- (a) D. Zhao, Y. Liao and Z. Zhang, *Clean*, 2007, **35**, 42; (b) A. S. Wells and V. T. Coombe, *Org. Process Res. Dev.*, 2006, **10**, 794.
- (a) V. Farmer and T. Welton, *Green Chem.*, 2002, **4**, 97; (b) S. Doherty, P. Goodrich, C. Hardacre, H. K. Luo, D. W. Rooney, K. R. Seddon and P. Styring, *Green Chem.*, 2004, **6**, 63; (c) E. Amigues, C. Hardacre, G. Keane, M. Migaud and M. O'Neill, *Chem. Commun.*, 2006, 72; (d) E. J. Amigues, C. Hardacre, G. Keane and M. E. Migaud, *Green*

- Chem.*, 2008, **10**, 660; (e) P. Goodrich, C. Hardacre, C. Paun, V. I. Parvulescu and I. Podolean, *Adv. Synth. Catal.*, 2008, **350**, 2473; (f) M. J. Earle, S. P. Katdare and K. R. Seddon, *Org. Lett.*, 2004, **6**, 707.
- M. Alvaro, B. Ferrer, H. García and M. Narayana, *Chem. Phys. Lett.*, 2002, **362**, 435.
- (a) C. Lee, T. Winston, A. Unni, R. M. Pagni and G. Mamantov, *J. Am. Chem. Soc.*, 1996, **118**, 4919; (b) A. Marcinek, J. Zielonka, J. Gebicki, C. M. Gordon and I. R. Dunkin, *J. Phys. Chem. A*, 2001, **105**, 9305.
- (a) G. S. Owens, A. Durazo and M. M. Abu-Omar, *Chem.-Eur. J.*, 2002, **8**, 3053; (b) A. Paul, P. K. Mandal and A. Samanta, *J. Phys. Chem. B*, 2005, **109**, 9148.
- For recent summarizing reviews, see: (a) R. M. Pagni, *Chim. Oggi*, 2007, **25**, 28; (b) R. M. Pagni and C. M. Gordon, in *CRC Handbook of Organic Photochemistry and Photobiology*, 2nd edn, W. M. Horspool and F. Lenci, (eds.), CRC Press, Boca Raton, 2004, Ch. 5, p. 1.
- A. Albin and M. Fagnoni, in *Green Chemical Reactions (NATO Science for Peace and Security Series C: Environmental Security)*, P. Tundo and V. Esposito, (eds.), Springer, Dordrecht, 2008, p. 173.
- (a) M. Oelgemöller, C. Jung, J. Ortner, J. Mattay and E. Zimmermann, *Green Chem.*, 2005, **7**, 35; (b) C. Schiel, M. Oelgemöller, J. Ortner and J. Mattay, *Green Chem.*, 2001, **3**, 224; (c) M. Oelgemöller, C. Jung and J. Mattay, *Pure Appl. Chem.*, 2007, **79**, 1939; (d) E. Haggiage, E. E. Coyle, K. Joyce and M. Oelgemöller, *Green Chem.*, 2009, **11**, 318.
- For a recent review on the photoacylation of quinones, see: (a) M. Oelgemöller and J. Mattay, in *CRC Handbook of Organic Photochemistry and Photobiology*, 2nd edn, W. M. Horspool and F. Lenci, (eds.), CRC Press, Boca Raton, 2004, Ch. 88, p. 1; for a general review on the photochemistry of quinones, see: (b) K. Maruyama and A. Osuka, in *The Chemistry of Quinonoid Compounds*, S. Patai and Z. Rappoport, (eds.), John Wiley & Sons, New York, 1988, Vol. 2, Ch. 13, p. 759.
- (a) F. Friedrichs, B. Murphy, D. Nayrat, T. Ahner, M. Funke, M. Ryan, J. Lex, J. Mattay and M. Oelgemöller, *Synlett*, 2008, 3137; (b) P. A. Waske, J. Mattay and M. Oelgemöller, *Tetrahedron Lett.*, 2006, **47**, 1329; (c) M. Oelgemöller, C. Schiel, R. Fröhlich and J. Mattay, *Eur. J. Org. Chem.*, 2002, 2465; (d) C. Schiel, M. Oelgemöller and J. Mattay, *Synthesis*, 2001, 1275; (e) C. Schiel, M. Oelgemöller and J. Mattay, *J. Inf. Rec.*, 1998, **24**, 257.
- (a) G. A. Kraus and M. Kirihara, *J. Org. Chem.*, 1992, **57**, 3256; (b) G. A. Kraus and P. Liu, *Tetrahedron Lett.*, 1994, **35**, 7723.
- R. Pacut, M. L. Grimm, G. A. Kraus and J. M. Tanko, *Tetrahedron Lett.*, 2001, **42**, 1415.
- General procedure for irradiation: In a typical photochemical experiment 63 mg (0.4 mmol) of 1,4-naphthoquinone and 2 mmol of aldehyde were dissolved by mild heating and vigorous stirring in 5 ml of ionic liquid in a Pyrex test tube (transmission ≥ 300 nm). The solution was degassed with argon and irradiated for 16 h using a Rayonet photochemical reactor (RMR-600; Southern New England Ultraviolet Company) equipped with 8 RPR-3000 Å lamps ($\lambda_{\max} = 300 \pm 25$ nm). The ionic liquid was then extracted with 7 × 10 ml portions of diethyl ether. The combined organic fractions were washed with saturated sodium bisulfite solution (3-times) and brine, dried over MgSO₄ and evaporated. The dried crude product was analysed by ¹H-NMR at this stage. Purified acylation products **3** were isolated by column chromatography (silica gel, ethyl acetate/*n*-hexane mixtures).
- (a) C. Hardacre, P. Nancarrow, D. W. Rooney and J. M. Thompson, *Org. Process Res. Dev.*, 2008, **12**, 1156; (b) C. Hardacre, S. P. Katdare, D. Milroy, P. Nancarrow, D. W. Rooney and J. M. Thompson, *J. Catal.*, 2004, **227**, 44.
- P. Hapiot and C. Lagrost, *Chem. Rev.*, 2008, **108**, 2238.
- S. C. Hubbard and P. B. Jones, *Tetrahedron*, 2005, **61**, 7425.
- J. L. Reynolds, K. R. Erdner and P. B. Jones, *Org. Lett.*, 2002, **4**, 917.
- (a) G. Zhu, J. Xu, G. Wu, H. Zhu, D. Long, S. Chen and S. Yao, *Int. J. Mol. Sci.*, 2006, **7**, 590; (b) M. J. Muldoon, A. J. McLean, C. M. Gordon and I. R. Dunkin, *Chem. Commun.*, 2001, 2364.
- For further examples, see: (a) R. A. Bourne, X. Han, A. O. Chapman, N. J. Arrowsmith, H. Kawanami, M. Poliakoff and M. W. George, *Chem. Commun.*, 2008, 4457; (b) N. Hoffmann, *Pure Appl. Chem.*, 2007, **79**, 1949; (c) M. Veerman, M. J. E. Resendiz and M. A. Garcia-Garibay, *Org. Lett.*, 2006, **8**, 2615; (d) A. V. Samoshin and V. V.

- Samoshin, *J. Undergrad. Chem. Res.*, 2006, **2**, 13; (e) A. G. Griesbeck, N. Maptue, S. Bondock and M. Oelgemöller, *Photochem. Photobiol. Sci.*, 2003, **2**, 450; (f) A. G. Griesbeck and A. Bartoschek, *Chem. Commun.*, 2002, 1594; (g) A. G. Griesbeck, W. Kramer and M. Oelgemöller, *Green Chem.*, 1999, **1**, 205.
- 21 (a) E. E. Coyle and M. Oelgemöller, *Photochem. Photobiol. Sci.*, 2008, **7**, 1313; (b) E. E. Coyle and M. Oelgemöller, *Highlights in Chemical Technology*, 2008, **5**, T95; (c) Y. Matsushita, T. Ichimura, N. Ohba, S. Kumada, K. Sakeda, T. Suzuki, H. Tanibata and T. Murata, *Pure Appl. Chem.*, 2007, **79**, 1959.
- 22 (a) C. Capello, U. Fischer and K. Hungerbühler, *Green Chem.*, 2007, **9**, 927; (b) K. Alfonsi, J. Colberg, P. J. Dunn, T. Fevig, S. Jennings, T. A. Johnson, H. P. Kleine, C. Knight, M. A. Nagy, D. A. Perry and M. Stefaniak, *Green Chem.*, 2008, **10**, 31.