

Reversible zwitterionic liquids, the reaction of alkanol guanidines, alkanol amidines, and diamines with CO₂

David J. Heldebrant,^{*a} Phillip K. Koech,^a M. Trisha C. Ang,^b Chen Liang,^b James E. Rainbolt,^a Clement R. Yonker^a and Philip G. Jessop^b

Received 25th November 2009, Accepted 22nd January 2010



First published as an Advance Article on the web 3rd March 2010

DOI: 10.1039/b924790d

Alkanolamidines, alkanolguanidines and diamines each react with CO₂ to form reversible zwitterionic liquids. CO₂ is chemically bound to the alcohol on alkanolamidines and alkanolguanidines as zwitterionic alkylcarbonates, while CO₂ is chemically bound on diamines as zwitterionic carbamates. All three classes of zwitterionic liquids could be reverted to their non-ionic forms by thermally stripping the CO₂ at temperatures near 50 °C, while one derivative was found to irreversibly form a cyclic carbonate.

Introduction



Carbon dioxide chemistry is increasingly relevant to real-world issues, thanks to global warming. Key chemistry issues currently being studied are the capture, storage, and utilization of CO₂. While the relevance of capture and storage are obvious, the relevance of CO₂ utilization is less clear. Although CO₂ utilization is unlikely to consume significant quantities of CO₂, it can be a significant strategy for the development of sustainable processes. As part of our research efforts towards CO₂ utilization, some of us invented switchable solvents, meaning solvents that can switch reversibly from one version to another.¹ Our original version was a mixture of an amidine and an alcohol (eqn (1), where B is a liquid amidine), but since then we and others have developed others such as guanidine/alcohol mixtures (eqn (1), where B is a liquid guanidine),² amidine/primary amine mixtures,³ and secondary amines (eqn (2)).⁴ Switchable solvents of these types all convert from a low-polarity non-ionic liquid to a high-polarity ionic liquid form upon exposure to an atmosphere of CO₂ and revert back to the low polarity form when the CO₂ is removed by heat or flushing with inert gas. We also demonstrated that these switchable solvents can be used as reversible CO₂-binding organic liquids (CO₂BOLs) for CO₂ capture, and are more energy-efficient in that role than aqueous solutions of ethanolamine.⁵



Because these switchable solvents consist of a CO₂-capturing group (an alcohol in eqn (1) and one equivalent of amine in eqn (2)) and a base (B in eqn (1) and the second equivalent of amine in eqn (2)), we conjectured that a single molecule containing both a CO₂-capturing group and a basic functional group

would also serve as a switchable solvent and CO₂BOL. These molecules could be an amidine-alcohol, a guanidine-alcohol, or a diamine (eqn (3) and (4)). The key differences in such switchable solvents compared to the earlier examples is that the polar, CO₂-containing form would be zwitterionic. If the reaction with CO₂ is reversible and the zwitterions are liquids, then these liquids could offer advantages over the previously-reported switchable solvents including the ease of use associated with having a single liquid component solvent rather than a mixture, and possibly an enhanced polarity difference between the ionic and non-ionic forms. There are very few reports of non switchable zwitterionic liquids in the literature,^{6–10} and none that we could find on their polarity, so it was difficult to predict the polarity of these proposed switchable zwitterionic liquids. The zwitterionic switchable solvents could also have advantages for CO₂ capture in that the gravimetric CO₂ capacity may be higher than in the two-component systems.

It is important not to confuse these proposed zwitterionic liquids with the mythical amidine-CO₂ zwitterionic adducts. Some groups have proposed zwitterionic adducts of amidines and guanidines with CO₂ in which the basic nitrogen atom binds directly to the CO₂ carbon. However such adducts have never been isolated or even observed spectroscopically; reports of spectroscopic detection of such adducts in the past have been erroneous due to confusion with the bicarbonate salts, a result of failure to rigorously remove water from the system before study.^{11–15} The zwitterions we report in this paper are entirely different in structure, having the CO₂ coordinated to an alkoxide oxygen (eqn (4)).



To our knowledge, the only known switchable zwitterionic liquid was developed in our laboratory.¹⁶ Our first generation switchable zwitterionic liquid was an alkanolamine, which reversibly captured SO₂ as an alkylsulfite salt. We hypothesized


^aPacific Northwest National Laboratory, Richland, WA, USA 99352.

E-mail: david.heldebrant@pnl.gov

^bDepartment of Chemistry, Queen's University, Kingston, ON, Canada K7L 3N6

Table 1 NMR chemical shifts of **1** in CDCl₃ with and without CO₂


¹ H	1 (OH)	2	3	4	5 (NH)	6	8, 10
1a	2.8 (s)	3.9 (t)	1.6 (m)	3.4 (t)	—	—	2.6 (s), 2.7 (s)
1b	—	4.2 (t)	1.8 (m)	3.7 (t)	6.0–7.0 (s)	—	2.8 (s)
¹³ C	1 (CO ₂)	2	3	4	5	6	8, 10
1a	—	62.0	23.3	42.4	—	158.9	35.7, 37.0, 39.8
1b	152.2	66.3	23.3	42.5	—	161.3	36.9, 39.7

**Fig. 1** Synthetic route to **1a** $n = 0$, **2a** $n = 4$, **3a** $n = 9$.

that a similar switchable zwitterionic liquid could be made with CO₂ if a sufficiently strong organic base such as a guanidine or amidine was paired with an alcohol in a single molecule (eqn (1), where “B” represents a basic group such as an amidine or guanidine). Alternatively, we also thought that a diamine, where one or both of the N atoms are secondary amines, would also react with CO₂ to give a zwitterionic liquid (eqn (3), where B’ represents a basic group such as an amine). Such systems could serve as either reversible CO₂-capturing agents or switchable solvents. We now report the preparation and study of these switchable zwitterionic liquids.

Results and discussion

We set out to synthesize a series of compounds (Scheme 1) that could react with CO₂ to make CO₂-activated switchable zwitterionic liquids.

**Scheme 1** Candidate precursors for preparing switchable zwitterionic liquids.


(a) Alkanol-guanidines

Our recent interest in CO₂BOLs^{1,2,5,17,18} led us to explore the functionalization of an amidine or guanidine with an alcohol to make a single component CO₂BOL. Combining the alcohol and the base components on one molecule (similar to alkanolamines) would simplify the binary alcohol/base solvent mixture into a unimolecular system, and potentially decrease component volatility through hydrogen bonding while increasing CO₂ weight capacities by using shorter alcohol chains. We chose

1,1,3,3-tetramethylguanidine (TMG) as our first substrate base because of its ease of functionalization.

Our first target molecules were alkanolguanidines (compounds **1a**, **2a**, and **3a** in Scheme 1) which were prepared from the reaction of 1,1,3,3-tetramethylguanidine with trimethylsilyl-protected bromoalkanol (Fig. 1). The resulting silyl-protected alkanolguanidines were then cleaved with K₂CO₃ in methanol.

Isolated compound **1a** was a clear and colorless non-viscous liquid. Spectroscopic characterization of **1a** was performed using both ¹H and ¹³C NMR in CDCl₃ (Table 1). The ¹H NMR of **1a** in CDCl₃ showed two inequivalent CH₃ peaks (carbons 8 and 10 using the numbering scheme in Fig. 2) at 2.6 and 2.7 ppm which we attribute to the lack of free rotation around the C=N double bond. Two triplets were observed at 3.9 and 3.4 ppm, which we assign to the CH₂ carbons 2 and 4 respectively. Carbon 3 was visible at 1.6 ppm as a multiplet and the alcohol O–H was observed at 2.8 ppm. The infrared (IR) spectrum of **1a** showed a broad O–H band at 3246 cm⁻¹ as well as strong (C=N) absorption bands at 1654 and 1618 cm⁻¹.

**Fig. 2** Numerical assignment of **1a**.

The reaction of anhydrous **1a** with CO₂ led to the formation of a highly viscous hazy liquid (**1b**), which we assigned the zwitterion structure (eqn (5)). The characteristic increase in viscosity has been observed when mixtures of amidines or guanidines and alcohols react with CO₂ to form their corresponding alkylcarbonate salts.^{1,2,4,5,17,18} The zwitterion **1b** was soluble in common organic solvents such as CDCl₃ and MeCN and was characterized by NMR. The ¹H NMR spectrum (CDCl₃) showed no evidence of the free alcohol OH. The spectrum also showed downfield shifting of the R–CH₂–O protons from 3.9 to

Table 2 NMR chemical shifts of **2a** in CDCl₃ with and without CO₂

¹ H	1 (OH)	2	3	4	5 (NH)	6	8, 10
2a	(s)	3.5 (t)	1.5, 1.7	3.1 (t)	—	—	2.5, (s) 2.7 (s)
2b	—	3.8 (t)	1.3, 1.5, 1.6	3.7 (t)	8.8 (s)	—	2.8 (s)


¹³ C	1 (CO ₂)	2	3	4	5	6	8, 10
2a	—	62.9	23.3	42.4	—	160.8	35.7, 37.0, 39.8
2b	158.6	65.1	23.3	42.5	—	163.2	36.9, 39.7

Table 3 NMR chemical shifts of **3a** in CDCl₃ with and without CO₂

¹ H	1 (OH)	2	3	4	5 (NH)	6	8, 10
3a	—	3.6 (t)	1.5 (m) 1.3 (s)	3.1 (t)	—	—	2.5, (s) 2.7 (s)
3b	—	3.8 (t)	1.2 (s) 1.5 (m) 1.6 (m)	3.1 (t)	—	—	2.9 (s) broad


¹³ C	1 (CO ₂)	2	3	4	5	6	8, 10
3a	—	62.8	26.0, 27.6, 29.7, 32.9	49.7	—	160.4	39.0, 39.8
3b	158.9	64.9	25.9–32.9	45.5	—	161.6	40.0

4.2 ppm indicating that CO₂ was chemically bound through the alcohol. The spectrum showed a broad singlet centered at 6.5 ppm corresponding to the guanidinium C=N–H. The viscous liquid could not be a bicarbonate salt of **1** because water was not present. If **1b** was a bicarbonate, there would be no downfield shifting of the alcohol CH₂O protons. The neat ¹³C spectrum of **1b** showed the guanidinium carbon at 161.3 ppm as well as the alkylcarbonate peak at 152.2 ppm. The guanidinium and carbonate peaks had the same integrated area, indicating stoichiometric absorption of CO₂. The ¹H and ¹³C chemical shifts are in agreement with our previous studies of amidinium and guanidinium alkylcarbonates.^{1,2,5,17,18,19}




The neat infrared (IR) spectrum of **1b** on NaCl plates showed strong bands at 1653 cm⁻¹ and 1615 cm⁻¹, which are comparable to other protonated amidine and guanidine imines [C=N–H]⁺ such as [DBUH]Cl (1645 cm⁻¹)¹ and [DBUH][OCO₂(CH₂)₅CH₃] (1648 cm⁻¹).² New strong bands appeared at 1590 cm⁻¹, 1403 cm⁻¹, and 1275 cm⁻¹ which we assign to the (C=O) of the carbonate. We also observed medium intensity N–H stretching bands at 3160 cm⁻¹. The free O–H of **1a** was undetectable when CO₂ was bound.

Gravimetrically, 0.502 grams (3 mmol) of **1a** absorbed 0.130 grams (3 mmol) of CO₂, corresponding to 20.6% CO₂ by weight. The zwitterion **1b** did not off-gas, rather it slowly cyclized into a cyclic carbonate. All attempts to thermally release CO₂ were unsuccessful because of condensation of the carboxylate to the activated imine carbon (6) to produce the cyclic ester (Fig. 3). The cyclic ester ¹³C spectrum (CDCl₃) showed a carboxylate peak at 153 ppm, the N₃C–O tetra-coordinated

**Fig. 3** The proposed structure of the cyclic carbonate product of **1b** heated above 25 °C.


carbon at 125 ppm, the CH₂–OCO₂ carbonate at 66 ppm, and the ¹H spectrum showed aliphatic CH₂–OCO₂– hydrogens at 4.1 ppm and CH₂–N hydrogens at 3.3 ppm. The CH₃ hydrogens were observed as an equivalent singlet at 2.7 ppm, and the internal R–CH₂–R hydrogens were observed as a pentet at 1.7 ppm. The cyclic carbonate is a non-viscous liquid that boils at 100 °C and does not release CO₂. The cyclic carbonate is likely forming because of the length of the propanol chain of the alcohol allowing the carboxylate to be in proximity to form a thermodynamically favorable 8-membered ring. Compound **1a** is not a useful reversible zwitterionic liquid because of its inability to release CO₂.

We hypothesized that a longer alkyl spacer between the alcohol and guanidine would inhibit cyclization, so we synthesized two longer chained TMG derivatives (compounds **2a** and **3a** in Scheme 1 and Fig. 4). Isolated compounds **2a** and **3a** were both hazy white non-viscous liquids. The ¹H NMR spectra of compounds **2a** and **3a** (Tables 2 and 3) were nearly identical to the compound **1a**, with the only difference being the

**Fig. 4** Numerical assignment of positions in compounds **2a** and **3a**.

6 or 9 aliphatic carbons of the alcohol chains. The infrared (IR) spectrum of **2a** and **3a** showed broad O–H bands at 3356 cm⁻¹ and 3381 cm⁻¹ as well as strong (C=N) absorption bands at 1604 cm⁻¹ and 1605 cm⁻¹ respectively.


Compound **3a** converts to a slightly yellow waxy solid in the presence of CO₂ (**3b**, eqn (6)). The solid was soluble in common organic solvents such as CDCl₃ and MeCN and was characterized by NMR. The ¹H and ¹³C spectra of this solid were similar to the propanol derivative **1b** except that only 74% CO₂ was absorbed.



The IR spectrum of the crude **3b** on NaCl plates showed what appeared to be a mixture of carboxylated and non-carboxylated TMG–(CH₂)₁₁–OH. New (C=N) bands at 1695 cm⁻¹, 1670 cm⁻¹, 1605 cm⁻¹, and (C=O) bands at 1591 cm⁻¹, 1402 cm⁻¹, 997 cm⁻¹, and 833 cm⁻¹ were observed for the carboxylated species. N–H stretching bands were not detected because of strong overlapping O–H bands at 3375 cm⁻¹. The solid does not melt, rather it decarboxylates completely at 65 °C. Attempts to crystallize the solid only yielded a waxy or gel like solid, which was not suitable for crystallographic analysis.

Our third target alkanolguanidine was a hexanol-functionalized TMG. The isolated TMG–(CH₂)₆–OH **2a** was a clear and colorless non-viscous liquid. The ¹H and ¹³C NMR spectra (CDCl₃) are presented in Table 3. The ¹H and ¹³C NMR spectra showed the same trends as the propanol and undecanol derivatives. The infrared (IR) spectrum of **2a** showed a broad O–H band at 3356 cm⁻¹ as well as strong (C=N) absorption bands at 1604 cm⁻¹.

The reaction of anhydrous **2a** with CO₂ leads to the formation of a highly viscous hazy liquid (**2b**, eqn (7)). The zwitterion was found to be soluble in CDCl₃ and MeCN as well as acetone. Nile red indicator dye was found to be insoluble in **2a** as well as **2b**, and no polarity data could be measured. No polarity measurements were performed on **1a,b** or **3a,b**. Table 3 lists the ¹H and ¹³C NMR spectra of the zwitterion in CDCl₃. The ¹H and ¹³C chemical shifts showed the same trends as the propanol and undecanol derivatives in the presence of CO₂.



The IR spectrum of **2b** on NaCl plates was a mixture of carboxylated and non-carboxylated material. We observed a new strong C=N band at 1618 cm⁻¹, strong C=O bands at 1600 cm⁻¹, 1403 cm⁻¹, and 1290 cm⁻¹ and broad O–H stretching bands at 3379 cm⁻¹. The N–H band was not detected because it was dwarfed under the very broad and strong O–H peak of **2a**.

Gravimetrically, 0.452 grams (2 mmol) of **2a** absorbed 0.046 grams (1 mmol) of CO₂, corresponding to 10.2% CO₂ by weight. The 0.5 molar equivalent of chemically bound CO₂ were thermally stripped from **2b** at 70 °C. The **2a** was then

re-contacted with CO₂ and then stripped for a total of three cycles. Each absorption cycle, **2a** bound 0.046 grams of CO₂, and each stripping released 0.046 grams of CO₂.


In our previous studies, dual component mixtures of TMG and linear alcohols in MeCN were unable to capture more than 0.47 molar equivalents of CO₂. Previous thermodynamic estimates of the energetics of CO₂ binding by mixtures of alcohols and TMG showed CO₂ binding was entropically unfavorable (on average $\Delta H = -160$ kJ mol⁻¹ and $\Delta S = -530$ J mol⁻¹).⁵

The single component showed slightly similar thermodynamic properties to our dual component mixtures of alcohols and guanidines. We estimate values from the Van't Hoff plot of the integrations of the hydrogens of the bound (CH₂–O–CO₂) and unbound (CH₂–OH) in CDCl₃ as a function of temperature to be $\Delta H = -87.0$ kJ mol⁻¹ and $\Delta S = -293$ J mol⁻¹ for **2a** and $\Delta H = -87$ kJ mol⁻¹ and $\Delta S = -291$ J mol⁻¹ for **3a**. In the zwitterionic experiments, *K* is defined as $[2b]/(P_{CO_2}[2a])$, which is different from the *K* defined in our dual component systems. The estimated thermodynamics from the single component systems are similar to estimates of the dual component systems in that the binding of CO₂ is entropically controlled. This suggests that alkanolguanidines bind CO₂ in a similar manner as dual component systems. The entropically unfavored binding of CO₂ makes TMG a poor choice for a base component for switchable zwitterionic liquids at STP, however they may be applicable in high pressure applications. We are investigating the functionalization of an amidine base like DBU with a hexanol chain, which we predict is likely to be more thermodynamically favored and may bind up to 1 molar equivalent of CO₂ at STP. The thermodynamic estimates to date suggest that the limitation of CO₂ binding is due to the thermodynamics of the alkylcarbonate guanidinium binding mode of CO₂BOLs and not the carbamate-like binding mode of alkanolamines.

CO₂ is bound through the alcohol moiety in alkanolguanidines in contrast to its binding to the amine of an alkanolamine because of two factors: the reduced nucleophilicity of guanidines compared to amines, and the higher basicity of amidines and guanidines compared to amines. To our knowledge, there are no published accounts of isolated carboguanidinium or carbamidinium zwitterions. We believe they are not isolable because amidines and guanidines are weak nucleophiles and weak hydrogen bond donors or acceptors²⁰ and are unable to react with CO₂ at standard temperature and pressure unless in the presence of a weak acid. The alcohol moiety of the alkanolguanidine is more nucleophilic than the delocalized guanidine component. Alcohols are known to react with CO₂ to form thermodynamically unstable alkylcarbonic acids.^{21,22} Alkylcarbonic acids have been stabilized by the addition of strong bases, such as DBU or TMG as is the case with our previous systems.^{1,2,5,17,19} We previously hypothesized that simple amines are not basic enough to accept the proton from alkylcarbonic acids and do not form alkylcarbonates⁵ like amidine and guanidine bases.

(b) Alkanol-Amidines

Amidines are slightly less basic than guanidines but are still sufficiently basic to serve as the basic group in a



zwitterionic liquid precursor. *N'*-(2-Hydroxyethoxy)ethyl-*N,N'*-dimethylacetamide (compound **4a** in Scheme 1 and Fig. 5) was prepared from aminoethoxyethanol by treatment with *N,N*-dimethylacetamide dimethylacetal and purified by vacuum distillation. The colorless liquid **4a** greatly increases in viscosity when CO₂ is bubbled through it. The product **4b** has been identified spectroscopically to be as shown in eqn (8).




Fig. 5 Numerical assignment of positions in compound **4**.

Upon the reaction of **4a** with CO₂, the mass of the liquid phase increases by 20%. The theoretical mass increase, according to the stoichiometry of the equation and neglecting dissolved CO₂, is 25%. This gravimetric evidence, as well as the NMR evidence, show that the reaction is incomplete. ¹³C NMR spectra of a solution of **4a** in DMSO (1 : 1 mixture by volume (Table 4)) show that, as CO₂ is bubbled through the solution, two small new peaks appear at 160.7 and 155.9 ppm, in addition to the peak for **4a** at 160.4 ppm. After further CO₂ treatment, the peaks have shifted to 161.6 and 156.2 and have become larger than the peak for **4a**. We attribute the new peaks to the protonated amidine group and the alkylcarbonate group of **4b**, respectively. In comparison, the alkylcarbonate peak of [DBUH][O₂COR] ionic liquid appears at 157–159 ppm depending on the alkyl group.² Heating the DMSO/**4b** mixture and bubbling argon through it restored the ¹³C-NMR spectrum of **4a**.

A dilute solution of **4a** in DMSO failed to react significantly with CO₂, suggesting that DMSO as a solvent inhibits the reaction. The same effect was observed for a dilute solution of **4a** in C₆D₆.

To determine the polarity change upon conversion from the amidine alcohol to the zwitterionic liquid, the spectrum of a solution of solvatochromatic dye Nile Red in liquid **4a** was obtained before and after treatment with CO₂. The wavelength of maximum absorbance (λ_{\max}) of a solution of Nile Red is indicative of the polarity of the solvent.²³ The λ_{\max} value of

Nile Red dissolved in neat **4a** was 544 nm before exposure to CO₂, indicating a polarity comparable to 1-octanol. After CO₂ was bubbled through the sample for 20 min at room temperature, the absorbance shifted to 554 nm, indicating a polarity that is greater than that of methanol. This amount of polarity change is comparable to that observed with the DBU/*n*-propanol switchable solvent,² showing that the amidine–alcohol single-component switchable solvent is just as effective as the DBU/*n*-propanol two-component switchable solvent and that there is no inherent advantage, in terms of increasing the size of the polarity change, to using a switchable zwitterionic liquid rather than a switchable ionic liquid.

(c) Diamines containing two secondary amines

Liquid secondary amines represent another kind of switchable solvent, wherein CO₂ is bound directly to the nitrogen atom in a carbamate salt rather than to an oxygen atom in an alkylcarbonate as is the case for alkanolguanidines and alkanolamidines.⁴ Amines such as *N*-ethyl-*N*-butylamine, *N*-ethyl-*N*-propylamine, dipropylamine, and benzylmethylamine, are low-polarity non-ionic liquids that are switched, by exposure to CO₂, into carbamate ionic liquids of increased polarity (eqn (2)). By heating the viscous liquid to 60 °C and purging it with N₂, the liquid can switch back to the low polarity form. In most cases, especially for primary amines, solid carbamate salts are formed following the addition of CO₂.

We speculated that a diamine in which both amines were secondary would function as a switchable solvent and as a potential CO₂-capturing liquid (eqn (9)). However, only a liquid diamine that gave a liquid zwitterion would be acceptable for these applications. We screened the liquid diamines of the structure RHN(CH₂)_{*n*}NHR where *n* = 2 (R = Me), *n* = 3 (R = Me, Et, *i*-Pr), and *n* = 6 (Me, Et, Bu). All gave solids upon reaction with CO₂. Note that in every case the diamines were used as received and not dried in any way. It is quite possible, for reasons explained below, that thoroughly dried amines might give liquids upon reaction with CO₂, but this possibility was not pursued because, for practical reasons, it is preferable to have a system that is water tolerant. The reaction

Table 4 NMR chemical shifts of **4** in DMSO-d₆ with and without CO₂

¹ H	1 (OH)	2–5	6 (NH)	7	8	10
4a	4.7 (br)	3.23 (t), 3.4–3.5 (m)	—	—	1.82 (s)	2.78 (s)
4b	7.4 ^a (br)	3.3–3.5 ^a (m)	—	—	2.03 ^a (br)	2.95 ^a (br)
4a +HCl	5.6 (br)	3.45 (m), 3.48 (m), 3.56 (m)	9.19 (s)	—	2.31 (s)	3.17 (s), 3.22 (s)
¹³ C	1 (CO ₂)	2–5	6	7	8	10
4a	—	51.1, 62.4, 73.0, 73.3	—	159.8 ^b	12.7	38.0
4b	155.9	47.0, 62.8, 70.2, 70.7	—	160.7	13.3	38.6
4a +HCl	—	43.8, 60.0, 69.1, 72.3	—	164.7	15.0	41.0, 39.8

^a These peaks are broadened due to high viscosity. ^b In very concentrated solution, this peak is at 160.4 ppm.

Table 5 NMR chemical shifts of **5** in CDCl₃ with and without CO₂

¹ H	1	2 (NH)	3	4	5	6	7
5a	2.3 (s)	1.0 (br)	2.5 (t)	1.6 (quint)	—	—	—
5b	2.6 (br)	10.6 (br)	2.9 (br) ^a	1.9 (br)	3.4 (br)	2.8 (br) ^a	—
¹³ C	1	2 (NH)	3	4	5	6	7
5a	36.7	—	50.7	30.2	—	—	—
5b	32.1	—	44.7 ^a	23.8	44.7 ^a	34.2	163.9

^a Overlapped.

of MeHN(CH₂)₃NHMe (**5a**) with CO₂ is described here, and is representative of the behaviour of these amines.

The viscosity of neat liquid **5a** increased as CO₂ was bubbled through it. Upon completion of the reaction, the liquid turned into a white solid, to which we assign the zwitterionic structure [MeH₂N⁺CH₂CH₂CH₂N(Me)CO₂⁻] (eqn (9)). The ATR-IR spectrum of the solid showed a band at 1545 cm⁻¹, which is characteristic of a carbamate anion.²⁴ No peak was observed at 835 cm⁻¹, which would have indicated the presence of a bicarbonate anion. The ¹H NMR spectrum of **5b** shows only minor changes in the chemical shifts of most carbon-bound protons except at positions 5 and 6 (see numbering scheme in Fig. 7), where the peaks have shifted downfield by 0.9 and ~0.6 ppm, respectively (Table 5). The N-bound proton has shifted from 1.0 ppm to 10.6 ppm, consistent with the change in charge on the N atom from neutral to positive. In the ¹³C NMR spectrum of **5b**, a new peak appears at 163.9 ppm, which corresponds to the carbamate central carbon. Crystals of the product were grown by slow evaporation of a solution of the compound in hexanes under a CO₂ atmosphere. The crystal structure confirms the formation of a zwitterionic salt co-crystallized with one water molecule (Fig. 6). The water molecule is an integral part of the hydrogen-bonding network in the structure, which suggested to us that if the water had been absent, the salt might not have solidified. However, treating thoroughly-dried **5a** with CO₂ also created a solid.




Fig. 6 Crystal structure of zwitterionic [MeH₂N⁺CH₂CH₂CH₂N(Me)CO₂⁻] \cdot H₂O (**5b**). Displacement ellipsoids are drawn at the 50% probability level.





Fig. 7 Numerical assignment of positions in **5a** and **5b**.



(d) Diamines containing one secondary amine and one tertiary amine

Because all diamines having two secondary amines gave solids, a diamine containing one secondary and one tertiary amine was tested. Bubbling CO₂ through neat liquid *N,N'*-trimethyl-1,3-propanediamine (compound **6a**) increased the viscosity of the liquid greatly, but no solidification was observed. We assign the zwitterionic structure [Me₂HN⁺CH₂CH₂CH₂N(Me)CO₂⁻] (**6b**) to the liquid product (eqn (10)). The ATR-IR spectrum of the solid showed the disappearance of the N–H band at 3298 cm⁻¹ and the appearance of a band at 1545 cm⁻¹, which is characteristic of a carbamate anion.²⁴ No peak was observed at 835 cm⁻¹, which would have indicated the presence of bicarbonate anion. The ¹H NMR spectrum of **6b** shows only minor changes in the chemical shifts of most carbon-bound protons except at positions 5 and 7 (see numbering scheme in Fig. 8), where the peaks have shifted downfield by 0.6 and 0.4 ppm, respectively. The N-bound proton peak for the secondary amine disappears and a new peak at 10.6 ppm appears, consistent with a dialkylammonium cation (Table 6). In the ¹³C NMR spectrum, a new peak appears at 163.9 ppm, which corresponds to the carbamate central carbon. High resolution ESI mass-spec (positive mode) of [Me₂HN⁺CH₂CH₂CH₂N(Me)CO₂⁻] was measured and matched the expected M + 1 peak of 161.13 m/z.




Fig. 8 Numerical assignment of positions in **6a** and **6b**.

The mass of liquid **6a** increased by 28 wt% due to uptake of CO₂. The theoretical increase in mass, assuming complete conversion but neglecting the mass of dissolved CO₂, should be 38%. The lower observed uptake suggests that the conversion of eqn (9) is incomplete, which may be the reason why the liquid did

Table 6 NMR chemical shifts of **6** in CDCl₃ with and without CO₂

¹ H	1	2 (NH)	3	4	5	6	7
6a	2.2 (s)	—	2.3 (t)	1.6 (quint)	2.6 (t)	1.4 (br)	2.4 (s)
6b	2.3 (s)	10.5 (br)	2.4 (t)	1.8 (m)	3.2 (br)	—	2.8 (br)
¹³ C	1	2 (NH)	3	4	5	6 (CO ₂)	7
6a	45.4	—	57.9	27.9	50.5	—	36.5
6b	44.8	—	56.7	25.1	47.2	161.1	34.0

not solidify. Removal of CO₂ and reversion of the liquid back to the neutral diamine form was achieved by bubbling argon through the viscous zwitterionic liquid at 60 °C. The reversion back to the diamine was confirmed by ¹H NMR spectroscopy.

To determine the polarity change upon conversion from the diamine to the zwitterionic liquid, the spectrum of the solvatochromatic dye Nile Red was obtained while the dye was in liquid **6** before and after treatment with CO₂. The λ_{max} value of Nile Red dissolved in neat **6a** was 523 before exposure to CO₂, indicating a polarity comparable to benzene. After CO₂ was bubbled through the sample with CO₂ for 20 min at room temperature, the absorbance shifted to 534 nm, indicating a polarity that is comparable to that of THF. This amount of polarity change is comparable to that observed with simple secondary amines such as ethyl butyl amine,⁴ showing that there is no inherent advantage, in terms of increasing size of the polarity change, to using switchable zwitterionic liquids rather than switchable ionic liquids.

Experimental

All reactions were run under an atmosphere of argon, unless otherwise indicated. Flasks were oven-dried and cooled under vacuum. All secondary diamines, solvatochromatic dye, CO₂ (supercritical fluid chromatography grade), argon (99.98%), chlorotrimethylsilane, triethylamine, 3-bromo-1-propanol, 6-bromo-1-hexanol, 11-bromo-1-undecanol and 1,1,3,3-tetramethylguanidine were purchased from Sigma-Aldrich and used as received. High-resolution mass spectra (HRMS) ESI for compounds **2Si–3Si**, **1a–3a** were obtained on a Thermo Scientific Exactive (tm) Mass Spectrometer from Thermo Scientific Company as *m/z* (relative intensity). High-resolution mass spectrum (HRMS) ESI for **6b** was obtained on a Qstar XL QqTOF from Applied Biosystems/MDS Sciex. Accurate masses are reported for the molecular ion (*M* + 1) or a suitable fragment ion. IR measurements for compounds **1Si–3Si**, **1a–3b** were performed on NaCl disks using a Nicolet Magna-750 spectrometer running on OMNIC software. The IR spectra for compounds **5a**, **5b**, **6a**, and **6b** were obtained on a Varian Scimitar 1000 FTIR equipped with the Pike MIRacle™ ZnSe ATR accessory. Proton nuclear magnetic resonance (¹H NMR) spectra for compounds **1Si–3Si**, **1a–3b** were recorded with a Varian (300 MHz) spectrometer. Chemical Shifts are reported in delta (δ) units, parts per million (ppm) downfield from tetramethylsilane. Coupling constants are reported in Hertz (Hz). Carbon-13 nuclear magnetic resonance (¹³C{¹H})

NMR) spectra for compounds **1Si–3Si**, **1a–3b** were recorded with a Varian 300 (75.5 MHz) spectrometer. Chemical shifts are reported in delta (δ) units, parts per million (ppm) relative to the center of the triplet at 77.00 ppm for deuteriochloroform. ¹H and ¹³C NMR spectra for compounds **4a–6b** were obtained with a Bruker AVANCE-400 MHz NMR spectrometer where each sample is referenced to tetramethylsilane. The NMR spectra of the CO₂ treated amines were obtained by bubbling CO₂ into a septa capped NMR tube that had 0.1 mL amine in dry CDCl₃. Data collection for the crystal of **5b** was done on a Bruker SMART APEX II X-ray diffractometer.

N,N,N',N'-Tetramethyl-*N''*-(3-trimethylsilyloxypropyl)-guanidine (**1Si**)

1,1,3,3-Tetramethylguanidine (TMG) (33.6 mL, 260.4 mmol) was charged into a 250 mL round bottom flask and then xylenes (25 mL) was added, this solution was heated to reflux. While maintaining the temperature (3-bromopropoxy)trimethylsilane²⁵ 25 g (118.39 mmol) was added. The resulting white suspension was refluxed overnight. After which the reaction was allowed to cool to room temperature and the resulting precipitate was filtered off, and the filtrate was distilled under vacuum, collecting xylenes followed by a second fraction compound **1a** as a colorless viscous liquid (9 g, 38%). ¹H NMR (300 MHz, CDCl₃) δ 3.67 (t, *J* = 5.4 Hz, 2H), 3.46 (t, *J* = 5.4 Hz, 2H), 2.00 (quin, *J* = 5.3 Hz, 2H), 0.09 (s, 9H). ¹³C{¹H} NMR (75.5 MHz, neat) δ 160.4, 61.1, 45.9, 39.7, 39.0, 35.6, -0.3. FTIR (NaCl, film): 2940, 1661, 1653, 1495, 1366, 1249, 1091, 839 cm⁻¹

N''-(3-Hydroxypropyl)-*N,N,N',N'*-tetramethylguanidine (**1a**)

1.3 g of compound **1Si** was charge into a clean dry 100 mL RB flask, and then anhydrous MeOH 20 mL was added. The solution was degassed by bubbling argon for 10 min and then K₂CO₃ 1.46 g was added, this mixture was stirred at RT for 1 h. The reaction mixture was then filtered and the resulting filtrate was distilled under vacuum to afford compound **1a** as a colorless liquid (0.73 g 80% yield). ¹H NMR (300 MHz, CDCl₃) δ 3.85 (t, *J* = 5.0 Hz, 2H), 3.33 (t, *J* = 5.0 Hz, 2H), 2.72 (br s, 1H), 2.19 (s, 6H), 2.58 (s, 6H) 1.64 (m, 2H). ¹³C{¹H} NMR (75.5 MHz, neat) δ 158.9, 62.0, 42.4, 39.8, 39.0, 35.7, 23.3 FTIR (NaCl, film): 3246, 2927, 1653, 1611, 1496, 1455, 13609, 1330, 1178, 1132, 1062 cm⁻¹ HRMS ESI (positive mode *M* + *H*⁺) calculated for C₈H₁₉N₃O 174.1600, found 174.1596.

N,N,N',N'-Tetramethyl-*N''*-(6-trimethylsilyloxyhexyl)-guanidine (**2Si**)

Following the same procedure as for **1Si** using, (6-bromohexoxy)trimethylsilane²⁷ (3.50 g, 13.90 mmol), TMG (3.80 mL, 30.55 mmol) and xylenes (5 mL) affords **2a** (1.8 g, 45%) as a pale yellow liquid. ¹H NMR (300 MHz, CDCl₃) δ 3.53 (t, *J* = 7.0 Hz, 2H), 3.06 (t, *J* = 7.0 Hz, 2H), 2.70 (s, 6H), 2.62 (s, 6H), 1.47 (m, 2H) 1.24 (m, 6H), 0.08 (s, 9H). ¹³C{¹H} NMR (75.5 MHz) δ 160.1, 62.9, 49.6, 39.8, 39.0, 32.9, 27.4, 25.9, -0.3. FTIR (NaCl, film): 2931, 2859, 1663, 1494, 1453, 1363, 1243, 1095, 840 cm⁻¹ HRMS ESI (positive mode M + H⁺) calculated for C₁₄H₃₃N₃O₂Si 288.2465, found 288.2453.

N''-(6-Hydroxyhexyl)-*N,N,N',N'*-tetramethylguanidine **2a**

Following the same procedure as for compound **1a** using compound **2Si** (1.70 g, 5.92 mmol), K₂CO₃ (1.63 g, 11.83 mmol), and MeOH (15 mL) provides **2a** (1.0 g, 78%) as a pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 3.58 (t, *J* = 6.7 Hz, 2H), 3.06 (t, *J* = 6.7 Hz, 2H), 2.70 (s, 6H), 2.61 (s, 6H), 1.51 (m, 4H) 1.32 (m, 4H). ¹³C{¹H} NMR (75.5 MHz) δ 160.8, 62.9, 42.4 39.8, 37.0, 35.7, 23.3. FTIR (NaCl, film): 3355, 2995, 2928, 1618, 1501, 1454, 13.74, 1238, 1137, 1102, 751 cm⁻¹ HRMS ESI (positive mode M + H⁺) calculated for C₁₁H₂₅N₃O 216.2070, found 216.2061.

N,N,N',N'-Tetramethyl-*N''*-(11-trimethylsilyloxyundecyl)-guanidine (**3Si**)

Following the same procedure as for **1Si** using, (11-bromoundecanoxy)trimethylsilane²⁶ (8.1 g, 25.10 mmol), TMG (7.0 mL, 55.30 mmol) and xylenes (10 mL) provide compound **3Si** (3.70 g, 41%) as a pale yellow liquid. ¹H NMR (300 MHz, CDCl₃) δ. 3.56 (t, *J* = 7.2 Hz, 2H), 3.04 (t, *J* = 7.2 Hz, 2H), 2.69 (s, 6H), 2.61 (s, 6H), 1.49 (quintet, *J* = 7.2 Hz, 4H), 1.23 (unresolved s, 14H), 0.24 (s, 9H). ¹³C{¹H} NMR (75.5 MHz, neat) δ 62.8, 49.7, 39.8, 39.0, 32.9, 32.8, 29.7, 27.6, 25.9, 2.1 FTIR (NaCl, film): 2925, 2853, 1604, 1502, 1456, 1375, 1252, 1136, 1059, 845 cm⁻¹. HRMS ESI (positive mode M + H⁺) calculated for C₁₉H₄₃N₃O₂Si 358.3248, found 358.3256.

N''-(11-Hydroxyundecyl)-*N,N,N',N'*-tetramethylguanidine (**3a**)

Following the same procedure as for compound **1a** using compound **3Si** (2.80 g, 7.83 mmol), K₂CO₃ (2.12 g, 16.79 mmol), and MeOH (20 mL) provides **3a** (2.16 g, 95%) as a pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ. 3.57 (t, *J* = 7.5 Hz, 2H), 3.04 (t, *J* = 7.5 Hz, 2H), 2.85 (s, 1H), 2.70 (s, 6H), 2.62 (s, 6H), 1.49 (quintet, *J* = 6.8 Hz, 4H), 1.23 (unresolved s, 14H). ¹³C{¹H} NMR (75.5 MHz, neat) δ 62.8, 49.6, 39.7, 38.9, 32.9, 37.7, 29.6 27.6, 25.9. FTIR (NaCl, film): 3381, 2921, 2852, 1605, 1501, 1455, 1375, 1145, 1063 cm⁻¹. HRMS ESI (positive mode M + H⁺) calculated for C₁₆H₃₅N₃O 286.2853, found 286.2855.

Preparation of *N'*-(2-hydroxyethoxy)ethyl)-*N,N*-dimethylacetamide (**4a**)

Aminoethoxyethanol (6.1 g) was charged into a round bottom flask. *N,N*-Dimethylacetamide dimethylacetal (8.0 g) was added to reaction mixture, which was then heated to 75 °C, stirred for

30 min, cooled, and concentrated in a rotary evaporator. A clear colorless liquid was recovered after vacuum distillation (yield 82%). ESI-MS (positive mode, M + H⁺) observed: 175.1447, calculated for C₈H₁₉N₂O₂: 175.1441. IR (neat, cm⁻¹): 3365 m, 2864 s, 1612 s, 1390 m, 1128 m, 1063 m, 1013 w, 891 w, 574 w.

Treatment of *N'*-(2-hydroxyethoxy)ethyl)-*N,N*-dimethylacetamide (**4a**) with CO₂

Carbon dioxide was bubbled, *via* a syringe needle, through a sample of **4a** in a test tube. Although the liquid became quite viscous after only 0.5 h, the bubbling was continued for another 3 h to ensure maximum possible conversion. IR (neat, cm⁻¹): 3366 m, 2871 m, 1646 s, 1396 m, 1287 m, 1126 m, 1066 m, 1013 m, 836 w, 621 w. NMR data given in Table 4. ESI-MS (negative mode, M+H⁺) observed: 217.1200, calculated for C₉H₁₇N₂O₄: 217.1188.

IR spectroscopic data for the bases before and after CO₂ treatment

N,N'-Dimethyl-1,3-propanediamine (**5a**). (neat, cm⁻¹) Before CO₂ treatment: 3280 m, 2929 m, 2879 m, 2841 m, 2786 s, 2694 w, 1473 s, 1446 s, 1124 s, 739 s. After CO₂ treatment: 3289 w, 2928 m, 2840 m, 2787 m, 2405 m, 2361 m, 1545 s, 1471 s, 1372 s, 1034 s, 1255 s, 1038 s, 807 s, 757 s.

N,N,N'-Trimethyl-1,3-propanediamine (**6a**). (neat, cm⁻¹) Before CO₂ treatment: 3298 w, 2940 s, 2857 m, 2814 s, 2783 s, 2764 s, 1739 w, 1460 s, 1376 w, 1263 w, 1150 m, 1122 m, 1041 s, 839 s, 743 s. After CO₂ treatment: 2941 m, 2814 m, 2762 m, 2462 w, 1683 w, 1624 w, 1545 w, 1460 s, 1376 s, 1305 m, 1262 s, 1040 s, 808 m.

Gravimetric uptake of CO₂ by *N,N,N'*-trimethyl-1,3-propanediamine (**6a**)

Compound **6a** (0.3 mL, 2.05 mmol) was added to a flame-dried 1 dram vial sealed with a rubber septa and massed. CO₂ was bubbled into the amine at room temperature and 1 bar to form the zwitterion. Mass was taken at 10 min intervals for 100 min although no further mass increase was observed after the first 10 min.

Polarity measurement of *N,N,N'*-trimethyl-1,3-propanediamine (**6a**)

Approximately 3 mL of the amine was placed in a 1 cm quartz cuvette where it was dotted with Nile red on a needle tip to give an absorbance between that of 0.5 and 1.0. The cuvette was then bubbled with atmospheric CO₂ for a period of 20 min or until the viscosity of the amine changed. A UV/visible spectrum was measured again. The experiment was done in triplicate, where the average of all three is the reported measurement.

Conclusions

Compounds **2a**, **4a**, and **6a** are switchable solvents, in that they react reversibly with CO₂ to form zwitterionic liquids. Any of these three compounds can serve as switchable solvents and as organic CO₂-capturing agents (CO₂BOLs). The

liquid zwitterionic salts were all highly viscous and polar. The change in polarity, upon conversion of the neutral molecule to the zwitterion, was comparable to that observed with non-zwitterionic switchable solvents. In terms of gravimetric CO₂-capturing capacity, these liquids are comparable to the two-component CO₂-binding liquids reported previously.

For alkanolguanidines, the chain length of the alkyl group between the alcohol and guanidine has an effect on the physical properties and regeneration of the alkanolguanidines. Short alkyl chains promoted cyclization of the carbonate to form the cyclic carbonate. Longer alkyl chains produced a reversible zwitterionic liquid, while very long alkyl chains produced reversible zwitterionic wax-like materials.

Di-secondary diamines and a secondary/tertiary diamine were found to form reversible zwitterionic carbamate salts, although only the secondary/tertiary diamine gave a liquid salt.

Acknowledgements

DJH, PKK and CRY gratefully acknowledge internal funding from Pacific Northwest National Laboratory (PNNL) Directed Research and Development. PNNL is proudly operated by Battelle for the U. S. Department of Energy. PGJ, TA, and CL thank the Natural Sciences and Engineering Research Council of Canada (NSERC), CFI/OIT, and the Canada Research Chairs program for funding the research, the equipment, and PGJ's salary, respectively. The authors thank Dr Rui Zhang for the high resolution mass spec. (HRMS) analysis and Dr Ruiyao Wang for the X-ray crystallography.

References

- 1 P. G. Jessop, D. J. Heldebrant, X. W. Li, C. A. Eckert and C. L. Liotta, *Nature*, 2005, **436**, 1102–1102.
- 2 L. Phan, D. Chiu, D. J. Heldebrant, H. Huttenhower, E. John, X. W. Li, P. Pollet, R. Y. Wang, C. A. Eckert, C. L. Liotta and P. G. Jessop, *Ind. Eng. Chem. Res.*, 2008, **47**, 539–545.
- 3 T. Yamada, P. J. Lukac, T. Yu and R. G. Weiss, *Chem. Mater.*, 2007, **19**, 4761–4768.
- 4 L. Phan, J. R. Andreatta, L. K. Horvey, C. F. Edie, A. L. Luco, A. Mirchandi, D. J. Darensbourg and P. G. Jessop, *J. Org. Chem.*, 2008, **73**, 127.
- 5 D. J. Heldebrant, C. R. Yonker, P. G. Jessop and L. Phan, *Energy Environ. Sci.*, 2008, **1**, 156–493.
- 6 D. Fang, Z.-h. Fei and Z.-l. Liu, *Monatsh. Chem.*, 2008, **139**, 799–803.
- 7 A. Narita, W. Shibayama and H. Ohno, *J. Mater. Chem.*, 2001, **11**, 1057.
- 8 H. Ohno, *Electrochemical Aspects of Ionic Liquids*, 1st edn, John Wiley & Sons, Hoboken, NJ, USA, 2005.
- 9 H. Ohno, M. Yoshizawa and W. Ogihara, *Electrochim. Acta*, 2003, **48**, 2079–2083.
- 10 M. Yoshizawa and H. Ohno, *Chem. Lett.*, 2004, **33**, 1594–1595.
- 11 B. Ochiai, K. Yokota, A. Fujii, D. Nagai and T. Endo, *Macromolecules*, 2008, **41**, 1229–1236.
- 12 T. Endo, D. Nagai, T. Monma, H. Yamaguchi and B. Ochiai, *Macromolecules*, 2004, **37**, 2007–2009.
- 13 F. S. Pereira, E. Ribeiro de Azevedo, E. F. da Silva, T. J. Bonagamba, D. L. d. S. Agostini, A. Magalhaes, A. E. Job and E. R. P. Gonzalez, *Tetrahedron*, 2008, **64**, 10097–10106.
- 14 E. R. Perez, M. O. da Silva, V. C. Costa, U. P. Rodrigues and D. W. Franco, *Tetrahedron Lett.*, 2002, **43**, 4091–4093.
- 15 E. R. Perez, R. H. A. Santos, M. T. P. Gambardella, L. G. M. de Macedo, U. P. Rodrigues, J. C. Launay and D. Franco, W., *J. Org. Chem.*, 2004, **69**, 8005–8011.
- 16 D. J. Heldebrant, P. K. Koech and C. R. Yonker, *Energy Environ. Sci.*, 2010, **3**, 111.
- 17 P. G. Jessop, D. J. Heldebrant, X. W. Li, J. Lu, J. P. Hallett, R. S. Jones, P. Pollet, C. A. Thomas, C. A. Eckert and C. L. Liotta, *Abstr. Pap. Am. Chem. Soc.*, 2005, **229**, U971–U971.
- 18 Y. X. Liu, P. G. Jessop, M. Cunningham, C. A. Eckert and C. L. Liotta, *Science*, 2006, **313**, 958–960.
- 19 D. J. Heldebrant, P. G. Jessop, C. A. Thomas, C. A. Eckert and C. L. Liotta, *J. Org. Chem.*, 2005, **70**, 5335–5338.
- 20 W. Galezowski, A. Jarczewski, M. Stanczyk, B. Brzezinski, F. Bartl and G. Zundel, *J. Chem. Soc., Faraday Trans.*, 1997, **93**, 2515–2518.
- 21 J. P. Hallett, P. Pollet, C. L. Liotta and C. A. Eckert, *Acc. Chem. Res.*, 2008, **41**, 458–467.
- 22 R. R. Weikel, J. P. Hallett, C. L. Liotta and C. A. Eckert, *Top. Catal.*, 2006, **37**, 75–80.
- 23 J. F. Deye, T. A. Berger and A. G. Anderson, *Anal. Chem.*, 1990, **62**, 615.
- 24 C. Hisatsune, *Can. J. Chem.*, 1984, **62**, 945–948.
- 25 K. Takeda, H. Haraguchi and Y. Okamoto, *Org. Lett.*, 2003, **5**, 3705.
- 26 S. P. Gido and J. Penelle, *Polymer*, 2002, 3007.
- 27 H. Mattes and C. Benzra, *J. Org. Chem.*, 1988, **53**, 2732.