


This article was downloaded by:

On: 15 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Green Chemistry Letters and Reviews

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t748292817>

A green synthesis of 2-phenyl/2-styrylchromones under solvent-free conditions using grinding technique

Dinesh Sharma^a; J. K. Makrandi^a

^a Department of Chemistry, Maharshi Dayanand University, Rohtak, Haryana, India

To cite this Article Sharma, Dinesh and Makrandi, J. K.(2009) 'A green synthesis of 2-phenyl/2-styrylchromones under solvent-free conditions using grinding technique', *Green Chemistry Letters and Reviews*, 2: 3, 157 – 159

To link to this Article: DOI: 10.1080/17518250903241966

URL: <http://dx.doi.org/10.1080/17518250903241966>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

RESEARCH LETTER

A green synthesis of 2-phenyl/2-styrylchromones under solvent-free conditions using grinding technique

Dinesh Sharma and J.K. Makrandi*

Department of Chemistry, Maharshi Dayanand University, Rohtak 124001, Haryana, India

(Received 2 June 2009; final version received 6 August 2009)

An efficient eco-friendly synthesis of flavones and 2-styrylchromones via cyclodehydration of corresponding 1-(2-hydroxyaryl)-3-aryl/styryl-1,3-propanediones is described under solvent-free conditions using grinding technique.


Keywords: flavones; 2-styrylchromones; cyclodehydration; grinding technique; eco-friendly reaction

Introduction

Flavones (2-phenylchromones) and 2-styrylchromones constitute an important class of naturally occurring compounds belonging to flavonoid group (1–4). These compounds have been reported to exhibit a variety of pharmacological properties (5–10) including anti-cancer (1) and antiHIV (11).

There are a number of methods available for the synthesis of chromone derivatives including the Allan–Robinson synthesis (12) and the oxidative cyclization of 2'-hydroxychalcones/2-hydroxycinnamylideneacetophenones (13). However, the cyclodehydration of 1-(2-hydroxyaryl)-3-aryl/styryl-1,3-propanediones obtained by the Baker–Venkataraman rearrangement of 2-aryloxy/cinnamoyloxy acetophenones remains the most practical method for their preparation (14). The cyclodehydration of the ensuing 1,3-propanediones requires heating under strongly acidic conditions using acetic acid (15), hydrochloric acid (16), sulfuric acid (17), and p-toulenesulphonic acid (18). Recently, the cyclodehydration of these 1,3-propanediones has been reported with CuCl_2 in ethanol (19) under microwave irradiation.

Today much emphasis is on the development of synthetic procedures which avoid toxic and hazardous chemicals and solvents (20). In continuation of our work to develop simple and eco-friendly procedures for the synthesis of organic compounds, we wish to report a high yield synthesis of flavones and 2-styrylchromones which involves the grinding of the intermediate 1,3-propanediones with phosphorous pentoxide in a mortar and pestle at room temperature in the absence of any solvent (Scheme 1). The moisture absorbed by the reaction mixture during the grinding



Scheme 1. Synthesis of flavone and 2-styrylchromone.

seems to be sufficient for the formation of a homogeneous mixture and the product is isolated by diluting the reaction mixture with ice cold water.

The possibility of the cyclodehydration of 1,3-propanediones occurring by phosphoric acid which could have formed during the grinding by absorption of water by phosphorous pentoxide, was investigated by grinding the 1,3-propanediones with phosphoric acid alone. No reaction was found to have taken place even after grinding for 30 minutes.

The validity of the procedure was checked by preparing differently substituted flavones and 2-styrylchromones. The identity of the compounds (Table 1) was confirmed from their IR, H^1 -NMR spectra and melting point comparison with literature value.

Experimental

All the chemicals were purchased from Aldrich and Fluka. Melting points were determined in open capillary tubes. IR (KBr) spectra were recorded in a Perkin–Elmer spectrum BX series FT-IR

*Corresponding author. Email: jagdish_chem2000@rediffmail.com

Table 1. Physical data of the compounds synthesized.

Compound	R ₁	R ₂	R ₃	Time (min) ^a	Melting point (°C)	Literature melting point (°C)	Yield (%) ^b
Ia	H	H	H	10	98–100	97 (21)	80
Ib	H	CH ₃	H	10	121	122 (21)	85
Ic	H	H	OCH ₃	10	160	157–158 (22)	90
Id	H	CH ₃	OCH ₃	10	166–168	170 (23)	95
Ie	OCH ₃	H	OCH ₃	10	147	145 (24)	90
If	OCH ₃	H	H	10	103	103–105 (25)	80
IIa	H	H	H	15	130	131 (26)	75
IIb	H	CH ₃	H	15	135	137 (27)	80
IIc	H	H	OCH ₃	15	170	167 (27)	95
IId	H	CH ₃	OCH ₃	15	157	160 (27)	85
IIe	OCH ₃	H	H	15	187	189–190 (28)	90

^aGrinding time.^bYield after crystallization.

spectrophotometer and ¹H NMR on Bruker Avance II 400 MHz instrument using tetramethyl-silane as an internal standard.

General procedure

Synthesis of flavones/2-styrylchromones

The substituted 1-(2-hydroxyphenyl)-3-phenyl/styryl-1,3-propanedione (2.1 mmol) was ground with phosphorous pentoxide (2.1 mmol) by a pestle in a mortar for 10–15 minutes. The completion of the reaction was checked by thin layer chromatography (benzene). The reaction mixture was diluted with ice-cold water and the solid that separated out was filtered at vacuum, washed with water, and recrystallized from methanol to afford flavones/2-styrylchromones.

4'-Methoxyflavone (Table 1, Compound Ic)

IR (KBr) 1646 (C=O); ¹H NMR (CDCl₃) δ 3.90 (s, 3H, OCH₃); 6.75 (s, 1H, H-3); 7.0 (d, *J*=9.0 Hz, 2H, H-3', H-5'); 7.25–7.55 (m, 3H, H-6, H-7, H-8); 7.80 (d, *J*=9.0 Hz, 2H, H-2', H-6'); 8.20 (d, *J*=9.0 Hz, 1H, H-5).

4'-Methoxy-6-methylflavone (Table 1, Compound Id)

IR (KBr) 1640 (C=O); ¹H NMR (CDCl₃) δ 2.30 (s, 3H, CH₃); 3.75 (s, 3H, OCH₃); 6.60 (s, 1H, H-3); 6.70 (d, *J*=9.0 Hz, 2H, H-3', H-5'); 7.30 (s, 2H, H-7, H-8); 7.75 (d, *J*=9.0 Hz, 2H, H-2', H-6'); 7.90 (s, 1H, H-5).

6-Methyl-2-styrylchromone (Table 1, Compound IIb)

IR (KBr) 1629 (C=O); ¹H NMR (CDCl₃) δ 2.45 (s, 3H, CH₃); 6.30 (s, 1H, H-3); 6.78 (d, *J*=16.0 Hz, 1H, H-α); 6.82–7.42 (m, 8H, C₆H₅, H-β, H-7, H-8); 7.90 (s, 1H, H-5).

6-Methyl-2-(4-methoxystyryl)chromone (Table 1, Compound IId)

IR (KBr) 1640 (C=O); ¹H NMR (CDCl₃) δ 2.40 (s, 3H, CH₃); 3.80 (s, 3H, OCH₃); 6.20 (s, 1H, H-3); 6.50 (d, *J*=16.0 Hz, 1H, H-α); 6.70–7.80 (m, 8H, C₆H₅, H-β, H-7, H-8); 7.90 (s, 1H, H-5).

Conclusion

In conclusion, it can be said that the present method developed for the synthesis of flavones and 2-styrylchromones is simple, highly efficient and eco-friendly, and avoids the use of organic solvent during the reaction and working up of the reaction mixture.

Acknowledgements

The authors are grateful to M.D. University, Rohtak for financial assistance to provide research fellowship to Dinesh Sharma.

References

- (1) Martens, S.; Mithofer, A. *Phytochemistry* **2005**, *66*, 2399–2407.
- (2) Malikov, V.M.; Yuldashev, M.P. *Chem. Nat. Compd.* **2002**, *38*, 358–406.
- (3) Huck, C.W.; Huber, C.G.; Ongania, K.H.; Bonn, G.K. *Chromatogr. A* **2000**, *870*, 453–462.
- (4) Nagao, T.; Abe, F.; Kinjo, J.; Okabe, H. *Biol. Pharm. Bull.* **2002**, *25*, 875–879.
- (5) Welton, A.F.; Tobias, L.D.; Fiedler-Nagy, C.; Anderson, W.; Hope, K.; Meyers, K.; Coffey, J.W. In *Plant Flavonoids in Biology and Medicine: biochemical, pharmacological, and structure-activity relationships*, Proceedings of a symposium held in Buffalo, New York, July 22–26, 1985; Cody, V., Middleton, E., Harborne, J.B., Eds.; A.R. Liss: New York, 1986; pp 231–242.

- (6) Xu, H.X.; Lee, S.F. *Phytother. Res.* **2001**, *15*, 39–43.
- (7) Goker, H.; Boykin, D.; Yildiz, S. *Bioorg. Med. Chem.* **2005**, *13*, 1707–1714.
- (8) Kim, H.P.; Son, K.H.; Chang, H.W.; Kang, S.S. *J. Pharmacol. Sci.* **2004**, *96*, 229–245.
- (9) Chu, H.; Wu, H.; Lee, Y. *Tetrahedron* **2004**, *60*, 2647–2655.
- (10) Silva, A.M.S.; Pinto, D.C.G.A.; Cavaleiro, J.A.S.; Levai, A.; Patonay, T. *ARKIVOC* **2004**, 2004; pp 106–123.
- (11) Wu, J.; Wang, X.; Yi, Y.; Lee, K. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1813–1815.
- (12) Allan, J.; Robinson, R.J. *J. Chem. Soc.* **1924**, *125*, 2192–2195.
- (13) Linuma, M.; Iwashima, K.; Matsuura, S. *Chem. Pharm. Bull.* **1984**, *32*, 4935–4941.
- (14) Varma, R.S.; Saini, R.K.; Kumar, D. *J. Chem. Res.(S)* **1998**, No. 1, 348–349.
- (15) Kumar, P.E.; Prashad, K.J.R. *Ind. J. Chem.* **1999**, *38B*, 1277–1279.
- (16) Jung, J.C.; Min, J.P.; Park, O.S. *Synth. Commun.* **2001**, *31*, 1837–1845.
- (17) Ares, J.J.; Outt, P.E.; Kakodkar, S.V.; Buss, R.C.; Geiger, J.C. *J. Org. Chem.* **1993**, *58*, 7903–7905.
- (18) Jain, P.K.; Makrandi, J.K.; Grover, S.K. *Curr. Sci.* **1981**, *50*, 857–858.
- (19) Kabalka, G.W.; Mereddy, A.R. *Tetrahedron Lett.* **2005**, *46*, 6315–6317.
- (20) Makrandi, J.K.; Lamba, M.S.; Kumar, S. *Green Chem. Lett. Rev.* **2008**, *1*, 123–125.
- (21) Dann, O.; Mylius, G. *Libigs Ann. Chem.* **1954**, *587*, 1–15.
- (22) Swapnil, S.R.; Pathan, M.Y.; Paike, V.V.; Pachmase, P.R.; Jadhav, W.N.; Pawar, R.P. *ARKIVOC* **2006**, 2006; pp 43–48.
- (23) Fitzgerald, D.M.; Sullivan, J.F.; Philbin, E.M.; Wheeler, T.S. *J. Chem. Soc.* **1955**, No. 1, 860–862.
- (24) Banerji, A.; Goomer, N. *Synthesis* **1980**, 1980; pp 874–875.
- (25) Lee, J.I.; Son, H.S.; Park, H. *Bull. Korean Chem. Soc.* **2004**, *25*, 1945–1947.
- (26) Cheema, U.S.; Gulati, K.C.; Venkataraman, K. *J. Chem. Soc.* **1932**, No. 1, 925–933.
- (27) Gaggad, H.L.; Wadodkar, K.N.; Ghiya, B.J. *Indian J. Chem.* **1984**, *24B*, 1244–1247.
- (28) Gulati, K.C.; Seth, S.R.; Venkataraman, K. *J. Chem. Soc.* **1934**, No. 1, 1765–1767.