

287. High Asymmetric Induction in Conjugate Additions of $\text{RCu} \cdot \text{BF}_3$ to Chiral Enoates

Preliminary Communication¹⁾

by **Wolfgang Oppolzer** and **Heinz J. Löher**

Département de Chimie Organique, Université de Genève, CH-1211 Genève 4

(11.XI.81)

Summary

1,4-Additions of $\text{PhCu} \cdot \text{BF}_3$, $n\text{-Bu} \cdot \text{BF}_3$ and $\text{MeCu} \cdot \text{BF}_3$ to the *trans*-8-phenylmenthyl enoates **1** proceeded with high chiral induction. Saponification of the resulting esters **2** gave the corresponding enantiomerically pure β -substituted alkanolic acids **3** and the recovered (-)-8-phenylmenthol in good overall yields. Analogous additions to the *cis*-crotonate **1** led preferentially to the acids **3** enantiomeric to those obtained from the *trans*-crotonate **1**, although with lower selectivity. A stereochemical model is proposed consistent with the observed results (*Scheme 2, Table*).

The highly enantioselective formation of C,C-bonds is of fundamental importance in organic synthesis²⁾. Elegant efforts to achieve asymmetric induction in *Michael* reactions involve the use of chiral basic catalysts [2], cosolvents [3] and nucleophiles [4]. However, enantioselective C,C-bond closure β to a carbonyl group has been accomplished more efficiently by 1,4-addition of organometallic reagents to α,β -unsaturated oxazolines [5], *t*-leucine-, *t*-butylester-aldimines [6], oxazepines [7], carboxylic amides derived from *l*-ephedrine [8] and α -carbonyl- α,β -ethylenic sulfoxides [9]³⁾. All these methods rely on a rigid substrate conformation owing to a chelation with the metal. We report here an entirely different concept which in terms of flexibility and efficiency offers an advantageous alternative.


Recently, the BF_3 -mediated 1,4-addition of organocopper reagents to α,β -unsaturated carbonyl compounds has been reported by *Yamamoto et al.* [12] (*Scheme 1*). The particularly smooth reaction may be attributed to a transition state **A** [12] or **B** implying either a synplanar or antiplanar $\text{C}=\text{C}/\text{C}=\text{O}$ -arrangement. In context with asymmetric ene [13] and cycloaddition reactions [14-16] we assume that enoates of secondary alcohols prefer a conformation where the $\text{C}=\text{O}$ is anti-

¹⁾ Presented by one of us (*W.O.*) at the 11th Northeast Regional Meeting of the American Chemical Society, Rochester, Oct. 19, 1981.

²⁾ Reviews [1].

³⁾ For further methods to generate a chiral center in a β -position to a carbonyl group see [10] [11].

Scheme 1



planar with the olefinic C,C-, and synplanar with the alkoxy-C,H-bonds; furthermore, in enoates derived from (-)-8-phenylmenthol [14] **1** or from other secondary alcohols [16] the appropriately positioned aryl group may shield selectively one enantiotopic face of the enoate. The conformational rigidity of such enoates as well as the π, π -orbital overlap should be enhanced by the coordination of the C=O group with a Lewis acid.

Accordingly we anticipated that a reagent $\text{R}^3\text{Cu} \cdot \text{BF}_3$ could add to the 8-phenylmenthyl enoates **1** preferentially at the olefinic face opposite to the phenyl ring via a transition state **B** (Scheme 2). The experimental results in support of this hypothesis are summarized in Scheme 2 and the Table.

Esterification⁴⁾ of *trans*-crotonic acid and *trans*-2-heptenoic acid [18] furnished the *trans*-enoates **1** ($\text{R}^1 = \text{H}$, $\text{R}^2 = \text{Me}$) and **1** ($\text{R}^1 = \text{H}$, $\text{R}^2 = n\text{-Bu}$), respectively whilst the *cis*-crotonate **1** ($\text{R}^1 = \text{Me}$, $\text{R}^2 = \text{H}$) was prepared by esterification⁴⁾ of tetrolic




Table. 1,4-Additions of $\text{R}^3\text{Cu} \cdot \text{BF}_3$ to chiral enoates **1**→**2** and enantiomeric purity of the thus obtained β -substituted alkanolic acids **3**

Entry	R ¹	R ²	R ³	Yield of 2 %	e.e. of 3 %
a	H	Me	C ₆ H ₅	76	> 99 ^{a)} 99.3 ^{b)}
b	H	Me	<i>n</i> -C ₄ H ₉	75.4	> 99 ^{c)} 99.5 ^{b)}
c	Me	H	C ₆ H ₅	36 (72) ^{d)}	24 ^{a)}
d	Me	H	<i>n</i> -C ₄ H ₉	76	70 ^{c)}
e	H	<i>n</i> -C ₄ H ₉	Me	28 (48) ^{d)}	78 ^{c)}

a) ¹H-NMR. (360 MHz) and HPLC. (SiO₂, hexane/EtOAc/MeOH 3:7:0.24) of amide derived from **3** and L-phenylalaninol.

b) **3a**: $[\alpha]_D^{25} = -56.8^\circ$ (*c* = 9, C₆H₆); **3b**: $[\alpha]_D^{25} = -4.18$ (neat).

c) HPLC. (SiO₂ hexane/EtOAc/MeOH 3:7:0.24) of amide derived from **3** and D-phenylglycinol.

d) Yield in parenthesis based on recovered **1**.

4) The corresponding acid was treated with (-)-8-phenylmenthol [14] (0.5 mol-equiv.), dicyclohexylcarbodiimide (1 mol-equiv.) and *p*-N,N-dimethylaminopyridine [17] in CH₂Cl₂ at room temperature for 16 h to give the esters **1** in 83–95% yield.

acid [19] and subsequent partial hydrogenation in presence of *Lindlar*-catalyst. In analogy to the described procedure [12], PhLi (2 mol-equiv.) was added to purified CuI (2 mol-equiv.) in ether at $-40^{\circ} \rightarrow -20^{\circ}$. Successive slow addition of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (2 mol-equiv.) and of the enoate **1a** (1 mol-equiv.) to the dark solution at -70° , stirring of the mixture for 10 min at -70° , raising the temperature slowly to $+25^{\circ}$, quenching with sat. aq. NH_4Cl -solution and work-up yielded the 1,4-adduct **2a**⁵ in 76% yield. Saponification⁶) of **2a** allowed the recovery of the auxiliary (-)-8-phenylmenthol (98% yield) and afforded (*R*)-(-)-3-phenylbutyric acid (**3a**) in over 99% enantiomeric purity. This dramatically high induction was assigned by chiroptic measurements [20] and further confirmed by conversion of **3a** into the amide of L-phenylalaninol⁷) and subsequent analysis of the latter [21] by HPLC. and ¹H-NMR. in comparison with the amide obtained⁷) from racemic 3-phenylbutyric acid and L-phenylalaninol. Under analogous reaction conditions as described for the conversion **1a** \rightarrow **2a** *n*-BuCu \cdot BF₃ was added to the *trans*-crotonate **1** giving the expected ester **2b** in 75.4% yield. Saponification⁶) of **2b** gave optically pure (-)-(*S*)-3-methylheptanoic acid (**3b**). The absolute configuration of **3b** was determined by its optical rotation [22] which indicated an enantiomeric purity of 99.5% in agreement with the HPLC. analysis [21] of the corresponding D-phenylglycinol amide⁷). Similar 1,4-addition of PhCu \cdot BF₃ and of *n*-BuCu \cdot BF₃ to the *cis*-crotonate **1** gave after saponification⁶) (+)-(*S*)-3-phenylbutyric acid (**3c**, 24% e.e.) and (+)-(*R*)-3-methylheptanoic acid (**3d**, 70% e.e.), respectively, which were analyzed as described above. Although the sign of asymmetric induction in the reaction *cis*-**1** \rightarrow **2c** and *cis*-**1** \rightarrow **2d** agrees with our stereochemical model its is definitely lower. *Cis/trans*-isomerization of the *cis*-crotonate **1** prior to 1,4-addition seems unlikely since *cis*-**1** was recovered unchanged after partial conversion to **2c**. Nevertheless, by reversing the order of group introduction, either enantiomer of a β -substituted carboxylic acid **3** should be accessible in high enantiomeric excess *via* the *trans*-esters **1**. Thus, addition of MeCu \cdot BF₃⁸) to the *trans*-2-heptenoate **1** and saponification⁶) of the adduct **2e** yielded (+)-(*R*)-3-methylheptanoic acid (**3d** \equiv **3e**) in over 72% chiral purity.

Further options include the use of new, either 'si-face' or 're-face' directing auxiliary alcohols, which are presently being developed in this laboratory⁹). Combination of these possibilities with a plethora of known stereocontrolled routes to α,β -conjugated esters [23] offers considerable promise. We are exploring systematically the scope and limitations of this asymmetric 1,4-addition reaction as well as its applicability to the synthesis of natural products.

⁵) All new compounds were characterized by IR., ¹H-NMR. (360 MHz) and mass spectroscopy.

⁶) Compound **2** (0.66 mmol) was heated in a mixture of 11.6N NaOMe in MeOH (7.5 ml) and water (4 mol-equiv.) at 75° for 24 h. Subsequent work-up and separation of neutral and acidic products furnished (-)-8-phenylmenthol and the corresponding acid **3**.

⁷) The corresponding acid **3** was heated in an excess of oxalyl chloride at 40° for 2 h. After evaporation a solution of the residue in dioxane was added slowly to a mixture of the chiral amine (1.1 mol-equiv.), NEt₃ (2 mol-equiv.) in dioxane at $+5^{\circ}$. Stirring of the mixture at RT. for 2 h, work-up and rapid chromatography (SiO₂, hexane/EtOAc, 2:3) gave the corresponding amide in over 70% yield.

⁸) 10 Mol-equiv., -65° for 10 h \rightarrow $+20^{\circ}$.

⁹) See for example the preceding communication [16].

Financial support of this work by the *Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung*, Sandoz SA, Basel, and Givaudan SA, Vernier, is gratefully acknowledged. We also thank Mr. J. P. Saulnier, Mr. A. Pinto and Mrs. D. Clément for NMR. and MS. measurements.

REFERENCES

- [1] a) J. D. Morrison & H. S. Mosher, 'Asymmetric Organic Reactions', Prentice-Hall: New Jersey 1971; b) H. B. Kagan & J. C. Fiaud, *Top. Stereochem.* 1978, 10, 175; c) D. Valentine jr. & J. W. Scott, *Synthesis* 1978, 329; d) J. W. ApSimon & R. P. Seguin, *Tetrahedron* 1979, 35, 2797.
- [2] a) H. Wynberg & R. Helder, *Tetrahedron Lett.* 1975, 4057; b) K. Hermann & H. Wynberg, *J. Org. Chem.* 44, 2238 (1979).
- [3] W. Langer & D. Seebach, *Helv. Chim. Acta* 63, 1710 (1979); and references therein.
- [4] a) J. S. Zweig, J. L. Luche, E. Barreiro & P. Crabbé, *Tetrahedron Lett.* 1975, 2355; b) L. Colombo, C. Gennari, G. Resnati & C. Scolastico, *Synthesis* 1981, 74; c) T. Takeda, T. Hoshiko & T. Mukaiyama, *Chemistry Lett.* 1981, 797.
- [5] A. I. Meyers, R. K. Smith & Ch. E. Whitten, *J. Org. Chem.* 44, 2250 (1979).
- [6] a) S. Hashimoto, S. Yamada & K. Koga, *J. Am. Chem. Soc.* 98, 7450 (1976); b) S. Hashimoto, N. Komeshima, S. Yamada & K. Koga, *Chem. Pharm. Bull.* 27 (10), 2437 (1979).
- [7] T. Mukaiyama, T. Takeda & K. Fujimoto, *Bull. Chem. Soc. Jpn.* 51, 3368 (1978).
- [8] T. Mukaiyama & N. Iwasawa, *Chemistry Lett.* 1981, 913.
- [9] G. H. Posner, J. P. Mallamo & K. Miura, *J. Am. Chem. Soc.* 103, 2886 (1981).
- [10] H. Kumobayashi, S. Akutagawa, S. Otsuka, *J. Am. Chem. Soc.* 100, 3949 (1978).
- [11] D. Valentine jr., K. K. Johnson, W. Priester, R. C. Sun, K. Toth & G. Saucy, *J. Org. Chem.* 45, 3698 (1980).
- [12] Y. Yamamoto & K. Maruyama, *J. Am. Chem. Soc.* 100, 3240 (1978).
- [13] W. Oppolzer, C. Robbiani & K. Bättig, *Helv. Chim. Acta* 63, 2015 (1980).
- [14] E. J. Corey & H. E. Ensley, *J. Am. Chem. Soc.* 97, 6908 (1975).
- [15] W. Oppolzer, M. Kurth, D. Reichlin & F. Moffatt, *Tetrahedron Lett.* 1981, 2545, and references cited therein.
- [16] W. Oppolzer, M. Kurth, D. Reichlin, C. Chapuis, M. Mohnhaupt & F. Moffatt, *Helv. Chim. Acta* 64, 2802 (1981).
- [17] B. Neises & W. Steglich, *Angew. Chem.* 90, 556 (1978); *Angew. Chem. Int. Ed.* 17, 522 (1978).
- [18] P. Coutrot, M. Snoussi & P. Savignac, *Synthesis* 1978, 133.
- [19] F.-M. Simmross & P. Weyerstahl, *Synthesis* 1981, 72.
- [20] V. Prelog & H. Scherrer, *Helv. Chim. Acta* 42, 2227 (1959).
- [21] G. Helmchen, G. Nill, D. Flockerzi, W. Schühle & M. S. K. Youssef, *Angew. Chem.* 91, 64 (1979); *Angew. Chem. Int. Ed. Engl.* 18, 62 (1979); G. Helmchen & G. Nill, *Angew. Chem.* 91, 66 (1979); *Angew. Chem. Int. Ed. Engl.* 18, 65 (1979).
- [22] P. A. Levene & R. E. Marker, *J. Biol. Chem.* 95, 153 (1932).
- [23] G. H. Posner, *Org. React.* 19, 1 (1972); N. Miyaura, N. Sasaki, M. Itoh & A. Suzuki, *Tetrahedron Lett.* 1977, 3369; Y. Yamamoto, H. Yatagai & K. Maruyama, *J. Org. Chem.* 44, 1744 (1979); J. Boutagy & R. Thomas, *Chem. Rev.* 74, 87 (1974); K. Tanaka, N. Yamagishi, R. Tanikaga & A. Kaji, *Bull. Chem. Soc. Jpn.* 52, 3619 (1979); K. Tamao, T. Kakiu & M. Kumada, *Tetrahedron Lett.* 1979, 619; N. Miyaura & A. Suzuki, *Chem. Lett.* 1981, 879; G. Friour, G. Cahiez, A. Alexakis & J. F. Normant, *Bull. Soc. Chim. Fr.* 1979, II-515; B. Bogdanović & B. Wermeckes, *Angew. Chem.* 93, 691 (1981); *Angew. Chem. Int. Ed. Engl.* 20, 684 (1981).