

141. Fluxional Behaviour of Isocyanide Derivatives of Dodecacarbonyltetrairidium

by Alessandra Orlandi¹), Renzo Ros²), and Raymond Roulet*

Institut de Chimie Minérale et Analytique de l'Université, 3, place du Château, CH-1005 Lausanne

(29. VII.91)

The fluxional behaviour of $[\text{Ir}_4(\text{CO})_{11}(t\text{-BuNC})]$ (**1**) and $[\text{Ir}_4(\text{CO})_{11}(\text{ArNC})]$ (Ar = 4-methoxyphenyl, 4-tolyl, phenyl, 4-chlorophenyl, 4-nitrophenyl; **2–5**) has been investigated by 2D and variable-temperature ^{13}C -NMR. These techniques give new evidence of the processes responsible for CO exchange, namely two successive merry-go-rounds causing *Cotton*-like scrambling and a CO rotation about one metal center.

Introduction. – The synthesis of two series of isocyanide derivatives of $[\text{Ir}_4(\text{CO})_{12}]$, namely $[\text{Ir}_4(\text{CO})_{12-x}\text{L}_x]$ ($x = 1\text{--}4$, L = *t*-BuNC, MeNC) has been reported by *Stunz* and *Shapley* [1]. Carbonyl scrambling in $[\text{Ir}_4(\text{CO})_{11}(t\text{-BuNC})]$ (**1**) has been studied by these authors through variable-temperature ^{13}C -NMR without simulation. Two mechanisms were proposed for the dynamic processes of lowest activation energy, consisting of two merry-go-rounds taking place, simultaneously, on two different faces of the Ir_4 tetrahedron.

We have re-examined the fluxionality of **1**, together with that of five new isocyanide derivatives $[\text{Ir}_4(\text{CO})_{11}(\text{ArNC})]$ (Ar = 4-methoxyphenyl, **2**; 4-tolyl, **3**; phenyl, **4**; 4-chlorophenyl, **5**; 4-nitrophenyl, **6**) by ^{13}C -NMR. The distinction between possible mechanisms is now based on dynamic connectivities derived from 2D-NOESY spectra and backed up by simulation of the variable-temperature ^{13}C -NMR spectra up to complete coalescence. The ArNC ligands in **2–6** were selected for their different χ parameters [2] to examine a possible effect of ligand basicity on the fluxional processes.


Results and Discussion. – The direct reaction of $[\text{Ir}_4(\text{CO})_{12}]$ with RNC (R = *t*-Bu, Me) in the presence or absence of Me_3NO in THF gives low yields of $[\text{Ir}_4(\text{CO})_{12-x}(\text{RNC})_x]$ ($x = 1, 2$). We have found that the reaction of $[\text{Ir}_4(\text{CO})_{11}\text{I}][\text{NEt}_4]$ with AgBF_4 and 1 mol-equiv. of isocyanide in CH_2Cl_2 gives better yields of $[\text{Ir}_4(\text{CO})_{11}(\text{RNC})]$ (88–96%) under milder conditions and without need of prep. TLC.

Cluster **1** adopts an unbridged C_s structure in the solid state [3]. This structure is also adopted by compounds **1–6** in solution as only IR absorptions in the region (2140–2200 cm^{-1}) appropriate for terminal isocyanide ligands and in the range (2000–2100 cm^{-1}) suitable for terminal CO ligands are observed.

The ^{13}C -NMR spectra of **1–6** display in each case one single carbonyl resonance around 310 K. The limiting low-temperature ^{13}C -NMR spectra show five resonances,

¹) Taken from the doctoral dissertation of A. O., Université de Lausanne.

²) Permanent address: Istituto di Chimica Industriale dell'Università, Via Marzolo 9, I-35131 Padova.



each of relative intensity two, and one resonance of unit relative intensity, all of which appear in the terminal CO region (see *Exper. Part*). The signal of relative intensity one is assigned to the unique carbonyl ligand *f* in pseudo-*trans*-position with respect to the isonitrile ligand. Inspection of the COSY ¹³C-NMR spectra of samples enriched with ¹³CO (ca. 30%; e.g. *Fig. 1* for ¹³C-enriched **1**) leads to the assignment of the signal of




Fig. 1. COSY ¹³C-NMR spectrum of ¹³C-enriched **1** in CD₂Cl₂ at 180 K

relative intensity two, and devoid of couplings to carbonyl ligands *b*. It is indeed well known that in carbonyltetrairidium clusters, the only CO's which couple significantly are in relative pseudo-*trans*-positions (in particular, geminal $^2J(\text{CO},\text{CO})$ are very small) [1]. The remaining four signals with cross peaks have identical vicinal $^3J(\text{CO},\text{CO})$ coupling constants (11.4 ± 0.8 Hz) and cannot be individually assigned; hence they are labelled 1, 2, 4, and 5 in order of decreasing chemical shift. Among the 24 virtually possible assignments, only eight are consistent with the couplings 1,4 and 2,5.





Fig. 2. NOESY ^{13}C -NMR spectrum of ^{13}C -enriched **1** in CD_2Cl_2 a) at 215 K (mixing time 50 ms); and b) at 225 K (mixing time 65.5 ms)

The NOESY ^{13}C -NMR spectrum of **1** at 215 K (Fig. 2a) gives the dynamic connectivities $1 \leftrightarrow b \leftrightarrow 4$ and corresponds to the exchange matrix of a first merry-go-round of 6 CO's around either the Ir(1)-Ir(2)-Ir(3) face (for 4 of the 8 possible assignments) or the Ir(2)-Ir(3)-Ir(4) (for the remaining ones). The NOESY spectrum at 225 K (Fig. 2b) shows 4 additional cross peaks which lead to the dynamic connectivities $1 \leftrightarrow 2 \leftrightarrow b \leftrightarrow 5 \leftrightarrow 4$ which corresponds to a second merry-go-round taking place on any of the equivalent Ir(1)-Ir(3)-Ir(4) or Ir(1)-Ir(2)-Ir(4) faces. The eight possible assignments are in agreement with such a mechanism. A merry-go-round of 5 CO's has never been observed when the ground state of the Ir_4 cluster contains a face bearing a radial L ligand, 3 edge-bridging, and two terminal CO's [4]. The proposed Scheme shows that the intermediate of the second merry-go-round in $[\text{Ir}_4(\text{CO})_{11}(\text{RNC})]$ contains a radial RNC ligand. However, as

Scheme



the ground state has here all terminal ligands, the only condition to be satisfied in the fluxional process is to end up with a state of equal energy with the same pseudo-*trans* relationship between RNC and carbonyl ligand *f* and without constraints on the choice of the face for the merry-go-round. This condition doesn't necessarily imply the participation of the RNC ligand in the CO site exchanges.

At ca. 270 K, a third process involving carbonyl ligand *f* in a 3 CO's exchange around one metal center (Ir(4)) is taking place.

Line-shape analysis [5] of the variable-temperature ^{13}C -NMR spectra was carried out using the following *Kubo-Sack* matrix elements: $(1,1) = (4,4) = -k_1/2 - k_2/2$, $(1,2) = (2,b) = (b,5) = (4,5) = k_2/2$, $(1,b) = (b,4) = k_1/2$, $(2,2) = -k_2$, $(b,b) = -k_1 - k_2$, $(5,5) = -k_2 - k_3/2$, $(5,f) = k_3/2$, $(f,f) = -k_3$, $(f,5) = k_3$, with $k_1, k_2, k_3 =$ rate constant in s^{-1} of the first and the second merry-go-round processes and of the rotation of 3 CO's at one metal center, respectively. An example of the obtained and simulated spectra is given in Fig. 3.

A simulation using matrix elements derived from the mechanism proposed by *Stunz et al.* [1] failed to reproduce variable-temperature ^{13}C -NMR spectra of **1**. However, other possible CO site exchange mechanisms should also be considered. First, one should take into account a process involving bridging of the isocyanide ligand, since edge-bridging RNC geometries have been observed in di- and trinuclear complexes [6]. Bridging of RNC on the Ir(1)-Ir(2) or Ir(1)-Ir(3) edge involved in merry-go-rounds around the Ir(1)-Ir(2)-Ir(4) or Ir(1)-Ir(3)-Ir(4) faces, respectively, gives an exchange matrix different from that corresponding to the observed connectivities of the second merry-go-round (Fig. 2b). This hypothetical matrix failed to reproduce experimental spectra.

Second, we have seen that the mechanism proposed for the lowest-energy process (Fig. 2a) is a merry-go-round through a CO-bridged intermediate having RNC either in axial or in apical position. When the ground state of Ir₄ clusters has 3 edge-bridging CO's defining a basal plane, as in $[\text{Ir}_4(\text{CO})_{11}(\text{PET}_3)]$ [7], the relative energies of the 3 possible isomers follow the sequence $L(\text{axial}) > L(\text{radial}) > L(\text{apical})$. If we assume that the same sequence of energy levels is applicable to the postulated intermediates of the first merry-go-round in $[\text{Ir}_4(\text{CO})_{11}(t\text{-BuNC})]$ (**1**), this same process could be due to a merry-go-round on the Ir(1)-Ir(2)-Ir(3) face, followed by another on the Ir(2)-Ir(3)-Ir(4) face. This alterna-




Fig. 3. a) Experimental and b) simulated variable-temperature ^{13}C -NMR spectra of ^{13}C -enriched **1** in CD_2Cl_2

tive mechanism agrees with the observed NOESY ^{13}C -NMR spectra of **1** at 225 K (Fig. 2b), if second-order peaks are taken into account. Therefore, a quantitative treatment of the 2D spectrum was needed. Recently, Orrell and coworkers [8] have proposed a general method for evaluating rate constants in complex exchange networks from two-dimensional NOESY NMR spectra. We have used their D2DNMR computer program capable of performing signal intensity to exchange rate calculation to distinguish between first- and second-order peaks. Including, as input data, the number of exchanging sites (6), their relative populations, and the experimental 2D signal intensities based on integration of all 2D peaks, a kinetic matrix was derived (see *Supplementary Material*) whose elements are the rate constants of the first-order exchange process. This matrix was equivalent to that of the proposed second merry-go-round, but not to that of the above alternative mechanism.

Table. Free Enthalpies of Activation ΔG^\ddagger (± 0.4 kJ mol⁻¹) of the 3 Fluxional Processes in $[\text{Ir}_4(\text{CO})_{11}(\text{RNC})]$ at 298 K

R	ΔG_1^\ddagger	ΔG_2^\ddagger	ΔG_3^\ddagger	R	ΔG_1^\ddagger	ΔG_2^\ddagger	ΔG_3^\ddagger
1 <i>t</i> -Bu	47.5	48.6	60.7	4 phenyl	49.4	49.9	60.8
2 4-methoxyphenyl	48.8	49.8	59.3	5 4-chlorophenyl	50.6	51.6	60.6
3 4-tolyl	49.0	50.4	59.9	6 4-nitrophenyl	49.8	53.2	62.4

The observed fluxional behaviour of cluster compounds **2–6** is the same as that of **1**, and can be viewed as a CO site exchange through two consecutive merry-go-rounds, followed by the rotation of 3 CO's around one metal center. The energy barriers are given in the *Table*.

The basicity of the non-CO ligand has no significant effect on the ΔG^\ddagger_{298} of the exchange processes, as also observed in $[\text{Ir}_4(\text{CO})_{11}\text{X}]^-$ (X = Br, I) and $[\text{Ir}_4(\text{CO})_{11}(\text{PAr}_3)]$ systems [9]. The rotation of apical CO's is also the highest activation energy process in the halogeno and phosphine clusters which have a different ground-state geometry.

We thank the *Swiss National Science Foundation* for financial support, and Dr. *Orrell* for a copy of the D2DNMR program.

Experimental Part

1. *General*. IR spectra: *Perkin-Elmer 880* spectrophotometer equipped with a data station; 0.1-mm CaF₂ cells previously purged with N₂. ¹³C-NMR spectra: *Bruker WH 360* (90.55 MHz); in CD₂Cl₂ at normal atmospheric pressure; δ in ppm, $J(\text{C,C})$ in Hz from COSY spectra. 2D ¹³C-NMR spectra: COSY experiment in CD₂Cl₂ at 180 K, 256 t_1 increments with 2-K transients, spectral width 900.90 Hz in the F_2 domain and 450.45 Hz in the F_1 domain; NOESY experiments in CD₂Cl₂ at 215 and 225 K, mixing time 50 and 65.5 ms, resp., 512 increments with 2-K transients, spectral width 1199.0 Hz in the F_2 domain and 599.52 Hz in the F_1 domain; a squared sine bell was applied in both domains prior to *Fourier* transformation.

2. *Isocyanide Ligands*. Commercially available *t*-BuNC was used (*Merck*) without further purification. The other ligands were synthesised starting from the corresponding formamides and following the method proposed by *Ugi* and coworkers [10]. When formamides were not commercially available (4-chloro-, 4-nitro-, and 4-methoxyphenyl-substituted ones), they were obtained from the corresponding amines according to [11].

3. *Monosubstituted Complexes*. A soln. of $[\text{Ir}_4(\text{CO})_{11}\text{I}][\text{NEt}_4]$ [12] (200 mg, 0.15 mmol) and the isocyanide ligand (0.15 mmol) was vigorously stirred, after addition of AgBF₄ (32.1 mg, 0.165 mmol) for 1 h at 263 K. The mixture containing solid AgI was filtered through a 15 × 3 cm column packed with silica gel (70–230 mesh) and eluted with cold CH₂Cl₂. The yellow filtrate was evaporated and the residue crystallised from CH₂Cl₂/hexane 1:1 at 248 K.

(*tert*-Butyl isocyanide)undecacarbonyltetrairidium ($[\text{Ir}_4(\text{CO})_{11}(\textit{t}\text{-BuNC})]$; **1**): For spectral characteristics, see [1].

Undecacarbonyl(4-methoxyphenyl isocyanide)tetrairidium ($[\text{Ir}_4(\text{CO})_{11}(4\text{-MeOC}_6\text{H}_4\text{NC})]$; **2**): IR (CH₂Cl₂, 298 K): 2091*m*, 2060*vs*, 2038*s*, 2018*s*, 2009*m* (CO); 2149*m* (CN). ¹³C-NMR (CD₂Cl₂, 190 K): 158.7 (*J* = 11.4, 1); 156.9 (*J* = 11.2, 2); 155.7 (*b*); 155.3 (*J* = 11.4, 4); 155.0 (*J* = 11.6, 5); 154.4 (*f*). Anal. calc. for C₁₉H₇Ir₄NO₁₂ (1210.14): C 18.86, H 0.58, N 1.16; found: C 19.20, H 0.59, N 1.17.

Undecacarbonyl(4-tolyl isocyanide)tetrairidium ($[\text{Ir}_4(\text{CO})_{11}(4\text{-MeC}_6\text{H}_4\text{NC})]$; **3**): IR (CH₂Cl₂, 298 K): 2092*m*, 2058*vs*, 2037*s*, 2016*s*, 2007*m* (CO); 2165*m* (CN). ¹³C-NMR (CD₂Cl₂, 190 K): 158.6 (*J*); 156.9 (2); 155.7 (*b*); 155.3 (4); 154.9 (5); 154.3 (*f*); same *J*'s as in **2**. Anal. calc. for C₁₉H₇Ir₄NO₁₁ (1194.1): C 19.11, H 0.59, N 1.17; found: C 19.34, H 0.6, N 1.18.

Undecacarbonyl(phenyl isocyanide)tetrairidium ($[\text{Ir}_4(\text{CO})_{11}(\text{PhNC})]$; **4**): IR (CH₂Cl₂, 298 K): 2092*m*, 2058*vs*, 2037*s*, 2016*s*, 2008*m* (CO); 2164*m* (CN). ¹³C-NMR (CD₂Cl₂, 190 K): 158.5 (*J*); 156.9 (2); 155.7 (*b*); 155.3 (4); 154.9

(5); 154.4 (f); same J 's as in 2. Anal. calc. for $C_{18}H_5Ir_4NO_{11}$ (1180.1): C 18.32, H 0.43, N 1.19; found: C 18.70, H 0.44, N 1.19.

Undecacarbonyl(4-chlorophenyl isocyanide)tetrairidium ($[Ir_4(CO)_{11}(4-ClC_6H_4NC)]$; 5): IR (CH_2Cl_2 , 298 K): 2092m, 2059vs, 2038s, 2016m (CO); 2161s, 2008m (CN). ^{13}C -NMR (CD_2Cl_2 , 190 K): 158.3 (1); 156.9 (2); 155.6 (b); 155.1 (4); 154.9 (5); 154.3 (f); same J 's as in 2. Anal. calc. for $C_{18}H_4ClIr_4NO_{11}$ (1214.6): C 17.8, H 0.33, N 1.16; found: C 18.5, H 0.34, N 1.16.

Undecacarbonyl(4-nitrophenyl isocyanide)tetrairidium ($[Ir_4(CO)_{11}(4-NO_2C_6H_4NC)]$; 6): IR (CH_2Cl_2 , 298 K): 2093m, 2059vs, 2036s, 2014s, 2006m (CO); 2183m (CN). ^{13}C -NMR (CD_2Cl_2 , 190 K): 157.8 (1); 156.9 (2); 155.4 (b); 155 (4); 154.8 (5); 154.4 (f); same J 's as in 2. Anal. calc. for $C_{18}H_4Ir_4N_2O_{13}$ (1225.1): C 17.65, H 0.33, N 2.29; found: C 17.88, H 0.34, N 2.27.

Supplementary Material. – The kinetic matrices for the D2DNMR program is available from *R. Roulet* upon request.

REFERENCES

- [1] G. F. Stunz, J. R. Shapley, *J. Organomet. Chem.* **1981**, 213, 329.
- [2] G. R. Van Hecke, W. De W. Horrocks, Jr., *Inorg. Chem.* **1966**, 5, 1960; C. A. Tolman, *Chem. Rev.* **1970**, 77, 313.
- [3] M. R. Churchill, J. P. Hutchinson, *Inorg. Chem.* **1979**, 9, 2451.
- [4] A. Strawczynski, R. Ros, R. Roulet, *Helv. Chim. Acta* **1988**, 71, 1885.
- [5] 'EXCHANGE', program library, Computing Center, University of Lausanne.
- [6] K. K. Joshi, O. S. Mills, P. L. Pauson, B. W. Shaw, W. H. Stubbs, *J. Chem. Soc., Chem. Commun.* **1965**, 181; F. A. Cotton, B. A. Frenz, *Inorg. Chem.* **1974**, 13, 253.
- [7] B. E. Mann, C. M. Spencer, A. K. Smith, *J. Organomet. Chem.* **1983**, 244, C17.
- [8] E. W. Abel, T. P. J. Coston, K. G. Orrell, V. Sik, D. Stephenson, *J. Magn. Reson.* **1986**, 70, 34.
- [9] G. Suardi, doctoral dissertation, University of Lausanne, 1991.
- [10] R. Obrecht, R. Herrmann, I. Ugi, *Synthesis* **1985**, 400.
- [11] W. Werner, *J. Chem. Res.* **1981**, 120.
- [12] P. Chini, G. Ciani, L. Garlaschelli, M. Manassero, S. Martinengo, A. Sironi, F. Canziani, *J. Organomet. Chem.* **1978**, 152, C35.