

146. 8-Azaadenine 2',3'-Dideoxyribonucleosides: Synthesis *via* 1,2,3-Triazolo[4,5-*d*]pyrimidinyl Anions

by Frank Seela* and Karin Mersmann


Laboratorium für Organische und Bioorganische Chemie, Fachbereich Biologie/Chemie, Universität Osnabrück,
Barbarastr. 7, D-4500 Osnabrück

(3. VII. 92)


The 8-azaadenine 2',3'-dideoxyribonucleosides 1–3 were synthesized *via* glycosylation of the 7-methoxy- or 7-amino-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidinyl anion with 5-*O*-[(*tert*-butyl)dimethylsilyl]-2,3-dideoxy- β -*D*-glycero-pentofuranosyl chloride (6). As 6 was an anomeric mixture, anomeric glycosylation products were formed. Moreover, three regioisomers (N^1 , N^2 , and N^3) were obtained in the case of the methoxy compound, but only two (N^1 and N^2) using 8-azaadenine. They were characterized by ^{13}C -NMR and CD spectra. The N-glycosylic bond of 8-aza-2',3'-dideoxyadenosine (1) was *ca.* 10 times more stable against acid than that of 2',3'-dideoxyadenosine. Compound 1 was deaminated by adenosine deaminase and showed inhibitory activity on HIV reverse transcriptase in form of its *O*-5'-triphosphate.

Introduction. – The 8-azaadenine nucleosides received considerable attention as purine antagonists in biological systems including viruses and cancer [1–3]. Analogs of the bases of common nucleosides were prepared [4]. The first azapurine, which was synthesized by Traube, was 1,3-dimethyl-7*H*-8-azapurine-2,6(1*H*,3*H*)-dione [5]. The chemistry of 8-azapurines (= 1*H*-1,2,3-triazolo[4,5-*d*]pyrimidines) is reviewed by Albert [6]. Whithin the series of 8-azaadenine derivatives, ribonucleosides [7–10], arabinonucleosides [11] [12] and 2'-deoxyribonucleosides [13] [14] were synthesized by common glycosylation techniques. Recently, we reported on the synthesis of 8-aza-2'-deoxyadenosine and related 2'-deoxyribonucleosides *via* nucleobase-anion glycosylation [15].


Purine 2',3'-dideoxy- β -*D*-ribonucleosides are highly active compounds against the human immunodeficiency virus (HIV), thereby inhibiting the HIV reverse transcriptase in form of their triphosphates [16]. Recently, we synthesized various deazapurine 2',3'-dideoxyribonucleosides and tested their activity [17–19]. We now focussed our interest on 8-azapurine 2',3'-dideoxyribonucleosides. In the following, we report on the synthesis, stability, and activity of the regioisomeric 8-azaadenine 2',3'-dideoxyribonucleosides 1–3 [20] related to $\text{A}_{\text{d}}^{2',3'}$ (4).




1



2



3 (systematic numbering)




4 (purine numbering)

Results and Discussion. – In 1974, the preparation of 8-azaadenine 2',3'-dideoxyribonucleoside (**1**) from the corresponding ribonucleoside was reported without substance characterization [21]. Compound **1** was also mentioned in another patent, again without any characterization [22]. Microbial preparation from 8-azaadenine and 2',3'-dideoxypyrimidine nucleosides was reported *via* transglycosylation [23] [24]. As the chemical synthesis of base-modified 2',3'-dideoxyribonucleosides from corresponding ribo-, or 2'-deoxyribonucleosides is laborious, we decided to use the direct glycosylation of the 8-azapurinyl anion, carrying a 6-amino or 6-methoxy group, by the halogenose **6**. According to the nucleophilicity of the triazole moiety, *N*-regioisomers were expected, as found in the case of 2'-deoxyribonucleosides [15].

The glycosylation of 7-methoxy-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidine (**5**) with 5-*O*-[(*tert*-butyl)dimethylsilyl]-2,3-dideoxy-D-*glycero*-pentofuranosyl chloride (**6**) was carried out in MeCN in the presence of a three-fold excess of powdered KOH and the cryptand tris[2-(2-methoxyethoxy)ethyl]amine (TDA-1), as described for the synthesis of 3-deazapurine and other base-modified 2',3'-dideoxyribonucleosides [17] [18]. The halogenose **6**


Scheme 1



was prepared from the corresponding lactol [25]. After workup, the reaction mixture was separated by repeated flash chromatography (FC; see *Exper. Part*), giving the six glycosylation products **7a–12a** in 60% overall yield. The ratio of the regioisomers was *ca.* 1:1:1, comparable with that of the corresponding 2'-deoxyribonucleosides [15]. A small excess of the β -D-anomers **7a–9a** was formed. Compounds **7a–12a** were desilylated with Bu_4NF in THF (\rightarrow **7b–12b**) and then treated with MeOH/NH_3 , to yield the crystalline β -D-anomers **1–3** and the corresponding α -D-anomers (see below **18–20**). Structural assignment of the glycosylation products are discussed below.

As the sugar halide **6** was an anomeric mixture, anomeric glycosylation products were formed under conditions (MeCN) which proceeded stereoselectively in the case of 2'-deoxyribonucleosides [15]. Therefore, non-stereoselective conditions (DMF) could be used as well. This allowed the use of 8-azaadenine (**13**) as base, thus circumventing the MeO/NH_2 displacement of the former reaction route (*Scheme 1*) and reducing the number of regioisomers to only two. The reaction of **13** with **6** was carried out in the presence of K_2CO_3 and TDA-1 (*Scheme 2*). After chromatographic workup (see *Exper. Part*; 32% overall yield), only **14** (faster migrating), **16** (slower migrating), and the anomers **15/17** (slowest migrating; not separated) were isolated. Neither an N^1 - nor an N^4 -isomer was obtained. Deprotection with Bu_4NF in THF afforded compounds **1, 18,**

Scheme 2



and **2/19**, respectively, in 80–90% yield. Contrary to **15/17**, the deprotected nucleosides **2/19** could be separated. The anomeric ratios of the amino derivatives were about the same as those found for the methoxy compounds. In opposition to the synthesis of 8-aza-2'-deoxyadenosine [15] which makes use of 7-methoxy-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidine (**5**), the synthesis of 8-aza-2',3'-dideoxyadenosine (**1**) gives better results with 8-azaadenine (**13**) because of a shorter synthetic sequence and less regioisomers formed, even if the total glycosylation yield is lower.

During the work on 8-azapurine 2'-deoxyribonucleosides, the ¹³C-NMR chemical shifts of the nucleobase moieties were already assigned [15]. We now measured gated-decoupled spectra of all 2',3'-dideoxyribonucleosides described above (Table 1). Moreover,

Table 1. ¹³C-NMR Chemical Shifts of 8-Aza-2',3'-dideoxyadenosine (**1**) and Related 1,2,3-Triazolo[4,5-*d*]pyrimidine 2',3'-Dideoxyribofuranosides in (D₆)DMSO at 23°

	C(3a) ^a	C(5)	C(7a)	C(7)	C(1')	C(2')	C(3')	C(4')	C(5')
1	148.5	156.8	123.9	156.2	85.9	30.6	27.0	82.7	63.8
2	157.5	157.1	125.5	156.9	94.4	31.9	26.5	83.8	63.9
3	160.9	154.5	113.6	151.9	89.5	29.4	25.3	81.8	63.0
7a	150.3	156.4	125.3	161.4	86.6	30.8	26.5	82.7	65.0
b	150.3	156.4	125.3	161.4	86.5	30.9	27.0	83.2	63.7
8a	159.2	155.9	126.1	162.3	95.2	32.0	25.8	83.7	65.0
b	159.1	155.9	126.1	162.3	95.3	32.4	26.4	84.3	63.8
9a^b	161.7	153.7	114.5	157.3	89.3	30.9	26.2	82.9	65.0
b^b	161.7	153.8	114.7	157.4	89.3	31.4	26.6	83.5	63.6
10a	150.2	156.4	125.4	161.4	87.2	30.6	26.2	81.5	64.9
b^c	150.2	156.2	125.4	161.4	87.2	30.4	26.4	82.1	63.2
11a	159.2	155.9	126.2	162.3	95.7	31.8	25.3	82.2	64.8
b^c	159.2	155.9	126.2	162.3	95.7	31.7	25.5	82.8	63.1
12a^b	161.6	153.7	114.3	157.2	89.9	30.8	25.8	81.3	64.8
b^b	161.7	153.9	114.4	157.5	90.0	31.0	26.1	81.9	63.2
14	148.6 ^d	156.8	123.9	156.1 ^d	85.7	30.4	26.6	82.2	65.1
15^e	157.5 ^d	157.1	125.5	156.9 ^d	94.3	31.6	25.9	83.2	65.2
16	148.5 ^d	156.9	124.0	156.1 ^d	86.3	30.2	26.3	81.1	64.8
17^e	157.5 ^d	157.1	125.5	156.9 ^d	94.7	31.4	25.4	81.7	64.7
18^e	148.5 ^d	156.9	124.0	156.1 ^d	86.3	30.1	26.5	81.7	63.1
19^e	157.5 ^d	157.1	125.5	156.7 ^d	94.7	31.3	25.6	82.3	63.1
20^e	160.9 ^d	154.5	113.7	151.9 ^d	89.1	30.2	25.4	83.3	63.0

^a) Systematic 1,2,3-triazolo[4,5-*d*]pyrimidine numbering. ^b) Aglycone signals according to the gated-decoupled spectra of 7-methoxy-1-[2'-deoxy-3',5'-di-*O*-(4-toluoyl)- β -D-*erythro*-pentofuranosyl]-1*H*-1,2,3-triazolo[4,5-*d*]pyrimidine [12]. ^c) According to gated-decoupled ¹³C-NMR spectra. ^d) Tentative. ^e) Taken from the anomeric mixture **15/17**.

the ¹³C,¹H-coupling constants were obtained from the gated-decoupled spectra of a set regioisomeric 7-amino- and 7-methoxy-1,2,3-triazolo[4,5-*d*]pyrimidine 2',3'-dideoxyribonucleosides (Table 2). It became apparent that in all cases, the coupling between C(3a) to the anomeric proton is low or not detectable. This is different to purine nucleosides, indicating that 8-azapurine 2'-3'-dideoxyribonucleosides prefer a high-*anti* conformation around the N-glycosylic bond, similar to other 8-azapurine nucleosides [26]. The lack of $J(C(3a),H-C(1'))$ makes the assignment of C(3a) vs. C(7) difficult in the

Table 2. $J(C,H)$ Values [Hz] of 1,2,3-Triazolo[4,5-d]pyrimidine 2',3'-Dideoxyribofuranosides^{a)}

	10b	11b	18	19	20
$J(C(3a),H-C(5))$	13	13	13	13	12
$J(C(3a),H-C(1'))$	1	–	–	–	–
$J(C(5),H-C(5))$	209	207	201	199	200
$J(C(7),H-C(5))$	11	11	11	11	12
$J(C(7),MeO)$	4	3	–	–	–
$J(C(7a),NH_2)$	–	–	4	<i>m</i>	<i>m</i>
$J(C(1'),H-C(1'))$	171	174	169	172	175
$J(C(2'),H-C(2'))$	135	135	135	139	135
$J(C(3'),H-C(3'))$	133	134	133	131	133
$J(C(4'),H-C(4'))$	147	148	148	147	147
$J(C(5'),H-C(5'))$	140	140	140	142	138

^{a)} Systematic numbering.

case of the NH_2 compounds; the MeO compounds, however, allow the assignment as two *q* appear for C(7). If the chemical shifts of C(7) of the N^1 -, N^2 -, and N^3 -regioisomers of the 7- NH_2 compounds are compared with those of the 7-MeO derivatives, almost the same chemical-shift differences are observed within the series of regioisomers. However, the NH_2 compounds resonate at higher field (5 ppm), which is in line with the change of *ipso*-substituents (Table 1). According to this, the chemical shifts of C(3a) vs. C(7) for the 8-azaadenine N^1 -(2',3'-deoxyribofuranosides) have to be reversed in [15].

The assignments of the anomeric configuration of the 2',3'-dideoxyribonucleosides are based on data summarized in Table 3. The NOE of $H-C(4')$ upon irradiation of $H-C(1')$ can be used for the assignment of the β -D-configuration [27]. In the case of 2',3'-dideoxynucleosides, this NOE is observed, apart from a smaller NOE in the case of the α -D-isomers. The small NOE of the α -D-anomers is due to the three-spin effect [28]. In

Table 3. Spectroscopic Data of Anomeric 8-Azaadenine 2',3'-Dideoxyribonucleosides

	NOE [%] at $H-C(4')$ ^{a)}	$\delta(C(4'))$	$R_f^c)$		$\delta(C(4'))$	$\Delta\delta(CH_3Si)$
1 (N^3,β -D)	1.5	82.7	0.55	7a (N^3,β -D)	82.7	0.044
2 (N^2,β -D)	1.6	83.8	0.30	8a (N^2,β -D)	83.7	0.052
3 (N^1,β -D)	2.0	83.3	0.25	9a (N^1,β -D)	82.9	0.040
18 (N^3,α -D)	0.8	81.7	0.40	10a (N^3,α -D)	81.5	0.016
19 (N^2,α -D)	0.9	82.3	0.25	11a (N^2,α -D)	82.2	0.017
20 (N^1,α -D)	0.9	81.8	0.25	12a (N^1,α -D)	81.3	0.015

^{a)} Upon irradiation of $H-C(1')$. ^{b)} Measured in (D_6)DMSO. ^{c)} TLC (silica gel, $CH_2Cl_2/MeOH$ 9:1).

the case of the amino nucleoside **3**, an additional NOE at the NH_2 group (2.4%) is detected upon irradiation of $H-C(1')$. This confirms N^1 as the glycosylation site (systematic numbering). The N^3 -glycosylation site is deduced from the typical up- and downfield shifts (10 ppm) of C(3a) and C(7a) induced by the substitution of the N-atom in α -position [29]. The assignment of the N^2 -isomers resulted from the absence of NOE's at $H-C(5)$, which should be present if the sugar is attached to N^4 or N^6 . The anomeric

assignments are also supported by other data summarized in *Table 3*, e.g. the chemical shifts of C(4') of the protected MeO compounds being always 1–1.5 ppm upfield shifted in the case of the α -D-anomers compared to the β -D-compounds. Furthermore, the ¹H-NMR chemical-shift differences of the Me signals of the (*t*-Bu)Me₂Si groups was 0.04 for the β -D-anomers and 0.02 for the α -D-anomers. Moreover, the deprotected MeO compounds were converted into the corresponding NH₂ nucleosides confirming these assignments. From the 8-azapurine 2',3'-dideoxyribonucleosides, only compound **1** can be converted into the corresponding inosine derivative (data not shown), which is an additional structural proof, at least within the series of the *N*³-regioisomers.

Ulbricht and coworkers [30] proposed that the anomeric configuration of nucleosides can be correlated to the CD spectra. Pyrimidine β -D-nucleosides should have a positive *Cotton* effect at 260 nm, while the α -D-anomers show the opposite behaviour. For purine nucleosides, the situation is different; β -D-purine nucleosides exhibit negative *Cotton* effects in the region of 260 nm and *vice versa*. However, it was already recognized that this rule depends on the electronic state and the torsion angle of the nucleobases [31]. We measured the CD spectra of compounds **1–3** and **18–20**. The *Figure* shows that the empirical CD rules cannot be transferred from purine to 8-azapurine nucleosides. Opposite CD spectra are observed in the anomeric pair of **3** and **20**. The situation becomes more difficult in the case of **1** and **18**. Assignment of the anomers **2** and **19** does not follow these rules.

Ribo- and 2'-deoxyribonucleosides can be cleaved at the N-glycosylic bond within cells by the action phosphorylases thereby loosing their activity. Hydrolysis depends on the activity of the enzyme but also on the chemical stability of the N-glycosylic bond. In particular, 2',3'-dideoxyribonucleosides are extremely labile [32]. It was shown that 8-azaadenine 2'-deoxyribonucleosides undergo acid-catalysed hydrolysis *via* cyclic glycosyl oxocarbenium ions as rate-limiting step with the *N*¹-regioisomers (systematic numbering) as the most labile and the *N*³-isomers as the most stable compounds [33]. *Table 4*

Table 4. Kinetic Data of *N*-Glycosylic-Bond Hydrolysis of Adenine and 8-Azaadenine 2',3'-Dideoxy-D-ribofuranosides

	<i>T</i> [°C]	<i>t</i> _{1/2} [min]	<i>c</i> _{HCl} [mol l ⁻¹]		<i>T</i> [°C]	<i>t</i> _{1/2} [min]	<i>c</i> _{HCl} [mol l ⁻¹]
1 (<i>N</i> ³ , β -D)	25	12.3	0.1	3 (<i>N</i> ¹ , β -D)	25	5.4	0.0005
18 (<i>N</i> ³ , α -D)	25	19.7	0.1	20 (<i>N</i> ¹ , α -D)	25	8.5	0.0005
2 (<i>N</i> ² , β -D)	25	11.8	0.1	4 (<i>N</i> ³ , β -D)	25	1.6	0.1
19 (<i>N</i> ² , α -D)	25	16.4	0.1				

summarizes the stability of the regioisomeric 8-azaadenine 2',3'-dideoxyribonucleosides. Again, the *N*¹-isomers are the most labile compounds, whereas the *N*²- and the *N*³-isomers are comparably stable. The β -D-anomers are hydrolysed slightly faster than the α -D-compounds by a factor of *ca.* 1.5 which is opposite to the 8-azaadenine 2'-deoxyribonucleosides [15] as well as to parent 2'-deoxyadenosine and its α -D-anomer [32]. In comparison to A_{43',3'} (**4**), compound **1** is approximately ten-fold more stable (*Table 4*).

Next, compounds **1** and **2** were phosphorylated in a one-pot reaction following a protocol originally developed for phosphorylation of purine nucleosides [34]. The nu-






Figure. CD Spectra of 8-Azaadenine 2',3'-dideoxyribonucleosides **1–3** and **18–20** measured at 5° in 0.06 M Na-cacodylate, pH 7.0

cleosides were dissolved in PO(MeO)₃ and treated at 0° with 2.7 equiv. of POCl₃ resulting in the formation of an activated dichlorophosphate which was directly condensed with [Bu₄N]₄P₂O₇. The triphosphates **21** and **22** were purified by DEAE-cellulose column chromatography and isolated as solid triethylammonium salts. Their inhibitory activity



21



22

was tested against HIV reverse transcriptase [35]: dideoxyadenosine analogue **21** showed an IC_{50} value of 42 μM against HIV reverse transcriptase, whereas the regioisomeric **22** was inactive ($> 100 \mu\text{M}$). Compared to the parent $\text{pppA}_{4,3',3'}$ [18], the *in vitro* activity of **21** is *ca.* 2 magnitudes lower, but it is in the range of the activity of 3-deaza-2',3'-dideoxyadenosine triphosphate ($IC_{50} = 79 \mu\text{M}$) [35].

We thank Dr. B. König from the *Boehringer Mannheim GmbH* for the inhibitory studies on HIV-reverse transcriptase and Dr. H. Rosemeyer for the discussion of the NMR spectra.

Experimental Part

General. See [15]. CD Spectra: *Jasco-600* spectropolarimeter; thermostatically controlled 1-cm cuvettes. Anal. TLC: glass plates coated with a 0.25-mm layer of silica gel *Sil G-25* with fluorescent indicator *UV₂₅₄* (*Merck*, Germany). Flash chromatography (FC): silica gel *60 H* (*Merck*, Germany); at 0.8 bar. Ion-exchange chromatography: *DEAE-Sephadex A25* (HCO_3^- form; *Kabi Pharmacia*, Sweden). Solvent systems: *A* = light petroleum ether/*AcOEt* 3:2, *B* = CH_2Cl_2 /acetone 95:5, *C* = light petroleum ether/*AcOEt* 7:3, *D* = CH_2Cl_2 /MeOH 95:5, *E* = CH_2Cl_2 /MeOH 9:1, *F* = CH_2Cl_2 /MeOH 8:2, *G* = light petroleum ether/*AcOEt* 1:4, *H* = *i*-PrOH/ NH_3 / H_2O 3:1:1.

Glycosylation of 7-Methoxy-3H-1,2,3-triazolo[4,5-d]pyrimidine (5) with 2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]-D-glycero-pentofuranosyl Chloride (6). To a suspension of KOH (0.75 g, 13.4 mmol) in MeCN (50 ml) were added TDA-1 (40 μl , 0.12 mmol) and **5** (0.9 g, 6 mmol) [36]. After stirring for 10 min at r.t., a freshly prepared cold (-80°) THF soln. (30 ml) of **6** (12 mmol; calculated on the basis of 100% yield of **6**) [17] [25] was added in portions of 5 ml within 30 min. Stirring was continued for another 30 min, insoluble material filtered off, and the filtrate poured into 20% aq. NaHCO_3 soln. (100 ml). The aq. layer was extracted twice with CH_2Cl_2 and the combined org. phase dried (Na_2SO_4) and evaporated. FC (column $50 \times 3 \text{ cm}$, *A*) gave five fractions. *Fr. 2*, which contained **11a** and some non-nucleoside by-products, was rechromatographed (FC, column $15 \times 3 \text{ cm}$, *B*) to give **11a**. Rechromatography of *Fr. 3* (FC, column $20 \times 3 \text{ cm}$, *B*) led to **7a** and **8a**.

3-{2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-glycero-pentofuranosyl}-7-methoxy-3H-1,2,3-triazolo[4,5-d]pyrimidine (**7a**). The faster-migrating part of *Fr. 3* yielded a colourless gum (265 mg, 12%). TLC (*B*): R_f 0.65. UV (MeOH): 250 (9900). $^1\text{H-NMR}$ (D_6)DMSO): -0.19 , -0.15 (2s, Me_2Si); 0.72 (s, *t*-Bu); 2.14–2.39 (m, $\text{CH}_2(3'')$); 2.59, 2.81 (2m, $\text{CH}_2(2'')$); 3.55 (dd, $J = 11.0, 5.9$, $\text{CH}_2(5'')$); 4.21 (s, MeO); 4.26 (m, H-C(4'')); 6.67 (dd, $J = 7.1, 1.7$, H-C(1'')); 8.79 (s, H-C(5)). Anal. calc. for $\text{C}_{16}\text{H}_{27}\text{N}_5\text{O}_3\text{Si}$ (365.51): C 52.58, H 7.45, N 19.16; found: C 52.53, H 7.44, N 18.92.

2-{2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-glycero-pentofuranosyl}-7-methoxy-2H-1,2,3-triazolo[4,5-d]pyrimidine (**8a**). The slower-migrating part of *Fr. 3* gave a colourless syrup (230 mg, 10.6%). TLC (*B*): R_f 0.55. UV (MeOH): 260 (9800). $^1\text{H-NMR}$ (D_6)DMSO): -0.16 , -0.11 (2s, Me_2Si); 0.72 (s, *t*-Bu); 2.17 (m, $\text{CH}_2(3'')$); 2.65, 2.56 (2m, $\text{CH}_2(2'')$); 3.66 (m, $\text{CH}_2(5'')$); 4.18 (s, MeO); 4.31 (m, H-C(4'')); 6.59 (d, $J = 5.6$, H-C(1'')); 8.75 (s, H-C(5)). Anal. calc. for $\text{C}_{16}\text{H}_{27}\text{N}_5\text{O}_3\text{Si}$ (365.51): C 52.58, H 7.45, N 19.16; found: C 52.52, H 7.51, N 19.10.

1-{2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-glycero-pentofuranosyl}-7-methoxy-1H-1,2,3-triazolo[4,5-d]pyrimidine (**9a**). *Fr. 4* yielded a colourless gum (240 mg, 11.0%). TLC (*A*): R_f 0.5. UV (MeOH): 270

(5500), 233 (4100). ¹H-NMR ((D₆)DMSO): 0.24, -0.20 (2s, Me₂Si); 0.67 (s, *t*-Bu); 2.15 (m, H-C(3')); 2.87, 2.60 (2m, CH₂(2')); 3.40, 3.59 (2m, CH₂(5')); 4.19 (s, MeO); 4.28 (m, H-C(4')); 6.68 (d, *J* = 6.9, H-C(1')); 8.80 (s, H-C(5)). Anal. calc. for C₁₆H₂₇N₃O₃Si (365.51): C 52.58, H 7.45, N 19.16; found: C 52.62, H 7.46, N 19.02.

3-(2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]-β-D-glycero-pentofuranosyl)-7-methoxy-3H-1,2,3-triazolo[4,5-d]pyrimidine (**10a**). Evaporation of the fast-migrating *Fr. 1* gave **10a** (225 mg, 10.3%). Colourless syrup. TLC (*C*): R_f 0.6. UV (MeOH): 250 (9800). ¹H-NMR ((D₆)DMSO): 0.03, 0.05 (2s, Me₂Si); 0.86 (s, *t*-Bu); 1.97 (m, H-C(3')); 2.6 (m, CH₂(2')); 3.66 (m, CH₂(5')); 4.21 (s, MeO); 4.40 (m, H-C(4')); 6.72 (dd, *J* = 6.8, 3.3, H-C(1')); 8.76 (s, H-C(5)).

2-(2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]-α-D-glycero-pentofuranosyl)-7-methoxy-2H-1,2,3-triazolo[4,5-d]pyrimidine (**11a**). From the slower-migrating part of *Fr. 2*, **11a** (180 mg, 8.3%) was obtained. Colourless oil. TLC (*C*): R_f 0.5. UV (MeOH): 260 (9600). ¹H-NMR ((D₆)DMSO): -0.06, 0.07 (2s, Me₂Si); 0.86 (s, *t*-Bu); 1.96 (m, H-C(3')); 2.5 (m, CH₂(2')); 3.68 (m, CH₂(5')); 4.18 (s, MeO); 4.50 (m, H-C(4')); 6.67 (dd, *J* = 5.9, 2.7, H-C(1')); 8.76 (s, H-C(5)).

1-(2,3-Dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]-α-D-glycero-pentofuranosyl)-7-methoxy-1H-1,2,3-triazolo[4,5-d]pyrimidine (**12a**). Evaporation of *Fr. 5* gave **12** (170 mg, 7.8%). Colourless oil. TLC (*A*): R_f 0.35. UV (MeOH): 270 (5500), 234 (4100). ¹H-NMR ((D₆)DMSO): 0.05, 0.06 (2s, Me₂Si); 0.87 (s, *t*-Bu); 1.98 (2m, CH₂(3')); 2.75, 2.6 (2m, CH₂(2')); 3.66 (m, CH₂(5')); 4.20 (s, MeO); 4.33 (m, H-C(4')); 6.77 (dd, *J* = 6.9, 2.9, H-C(1')); 8.80 (s, H-C(5)).

3-(2,3-Dideoxy-β-D-glycero-pentofuranosyl)-7-methoxy-3H-1,2,3-triazolo[4,5-d]pyrimidine (**7b**). To a soln. of **7a** (510 mg, 1.4 mmol) in THF (20 ml) was added 1.1M Bu₄NF in THF (2 ml). The mixture was stirred for 1 h at r.t. FC (column 20 × 3 cm, *D*) afforded **7b** (295 mg, 84%). Colourless solid. TLC (*E*): R_f 0.70. UV (MeOH): 251 (10300). ¹H-NMR ((D₆)DMSO): 2.24 (m, H-C(3')); 2.63, 2.74 (2m, CH₂(2')); 3.42 (m, CH₂(5')); 4.22 (m, H-C(4'), MeO); 4.70 (*t*, *J* = 5.7, OH-C(5')); 6.66 (dd, *J* = 7.2, 2.2, H-C(1')); 8.80 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 48.11, H 5.18, N 27.74.

2-(2,3-Dideoxy-β-D-glycero-pentofuranosyl)-7-methoxy-2H-1,2,3-triazolo[4,5-d]pyrimidine (**8b**). As described for **7b**, **8a** (580 mg, 1.59 mmol) was treated with Bu₄NF/THF. After chromatography (column 20 × 3 cm, *E*), **8b** (330 mg, 83%) was obtained. Colourless solid. TLC (*E*): R_f 0.65. UV (MeOH): 260 (10200). ¹H-NMR ((D₆)DMSO): 2.17 (m, H-C(3')); 2.57 (m, CH₂(2')); 3.50 (m, CH₂(5')); 4.16 (s, MeO); 4.28 (m, H-C(4')); 4.76 (*t*, *J* = 5.6, OH-C(5')); 6.60 (dd, *J* = 4.8, 3.2, H-C(1')); 8.75 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 47.90, H 5.22, N 27.76.

1-(2,3-Dideoxy-β-D-glycero-pentofuranosyl)-7-methoxy-1H-1,2,3-triazolo[4,5-d]pyrimidine (**9b**). Deprotection of **9a** (285 mg, 0.78 mmol) was carried out as described for **7b**. FC (column 20 × 3 cm, *E*) afforded **9b** (140 mg, 72%). Colourless solid. TLC (*E*): R_f 0.6. UV (MeOH): 271 (6000). ¹H-NMR ((D₆)DMSO): 1.97 (m, H-C(3')); 2.71, 2.6 (2m, CH₂(2')); 3.47 (m, CH₂(5')); 4.21 (s, MeO); 4.29 (m, H-C(4')); 4.81 (*t*, *J* = 5.7, OH-C(5')); 6.79 (dd, *J* = 7.4, 3.3, H-C(1')); 8.75 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 47.97, H 5.20, N 27.70.

3-(2,3-Dideoxy-α-D-glycero-pentofuranosyl)-7-methoxy-3H-1,2,3-triazolo[4,5-d]pyrimidine (**10b**). As described for **7b**, **10a** (420 mg, 1.15 mmol) was treated with TBAF/THF (2 h). Evaporation and FC (column 20 × 3 cm, *E*) gave **10b** (220 mg, 76%). Colourless solid. TLC (*E*): R_f 0.55. UV (MeOH): 251 (10300). ¹H-NMR ((D₆)DMSO): 1.97 (m, H-C(3')); 2.63 (m, CH₂(2')); 3.47 (m, CH₂(5')); 4.21 (s, MeO); 4.36 (m, H-C(4')); 4.81 (*t*, *J* = 5.7, OH-C(5')); 6.74 (dd, *J* = 7.0, 3.4, H-C(1')); 8.79 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 48.02, H 5.33, N 27.71.

2-(2,3-Dideoxy-α-D-glycero-pentofuranosyl)-7-methoxy-2H-1,2,3-triazolo[4,5-d]pyrimidine (**11b**). As described for **7b**, **11a** was prepared from **11a** (280 mg, 0.77 mmol; 75 min). FC (column 20 × 3 cm, *D*) yielded **11b** (140 mg, 73%). Colourless solid. TLC (*E*): R_f 0.60. UV (MeOH): 260 (10100). ¹H-NMR ((D₆)DMSO): 1.94, 2.34 (2m, CH₂(3')); 2.56 (m, CH₂(2')); 3.48 (m, CH₂(5')); 4.18 (s, MeO); 4.46 (m, H-C(4')); 4.85 (*t*, *J* = 5.8, OH-C(5')); 6.67 (dd, *J* = 6.2, 2.5, H-C(1')); 8.76 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 47.93, H 5.21, N 27.68.

1-(2,3-Dideoxy-α-D-glycero-pentofuranosyl)-7-methoxy-1H-1,2,3-triazolo[4,5-d]pyrimidine (**12b**). Deprotection of **12a** (350 mg, 0.96 mmol) was carried out as described for **7b** (30 min). FC (column 20 × 3 cm, *E*) yielded **12b** (185 mg, 77%). Colourless solid. TLC (*E*): R_f 0.55. UV (MeOH): 271 (5800). ¹H-NMR ((D₆)DMSO): 2.19 (m, CH₂(3')); 2.6, 2.79 (2m, CH₂(2')); 3.33 (m, CH₂(5')); 4.20 (s, MeO); 4.23 (m, H-C(4')); 4.64 (*t*, *J* = 5.6, OH-C(5')); 6.69 (d, *J* = 5.7, H-C(1')); 8.75 (s, H-C(5)). Anal. calc. for C₁₀H₁₃N₃O₃ (251.24): C 47.81, H 5.22, N 27.88; found: C 47.75, H 5.33, N 27.73.

7-Amino-3-(2,3-dideoxy-β-D-glycero-pentofuranosyl)-3H-1,2,3-triazolo[4,5-d]pyrimidine (1). From **7b**: For 12 h **7b** (250 mg, 1 mmol) was stirred in MeOH (saturated with NH₃ at 0°; 20 ml). The mixture was evaporated, the residue dissolved in CH₂Cl₂/MeOH 9:1, and the soln. filtered over a 5-cm layer of silica gel. The filtrate was evaporated, and crystallization of the residue from MeOH afforded **1** (210 mg, 89%). Colourless needles. M.p. 182°. TLC (*E*): R_f 0.55. UV (MeOH): 278 (11000). ¹H-NMR ((D₆)DMSO): 2.24 (*m*, H-C(3')); 2.6, 2.70 (2*m*, CH₂(2')); 3.46 (*m*, CH₂(5')); 4.21 (*m*, H-C(4')); 4.81 (*t*, *J* = 5.7, OH-C(5')); 6.55 (*dd*, *J* = 7.1, 2.7, H-C(1')); 8.14, 8.47 (2*s*, NH₂); 8.33 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 45.89, H 5.30, N 35.64.

From **14**: A soln. of **14** (250 mg, 0.7 mmol) in THF (10 ml) was treated with 1.1*M* Bu₄NF in THF (2 ml) and stirred for 30 min at r.t. FC (column 10 × 3 cm, *E*) afforded **1** (140 mg, 83%).

7-Amino-2-(2,3-dideoxy-β-D-glycero-pentofuranosyl)-2H-1,2,3-triazolo[4,5-d]pyrimidine (2). From **8b**: As described for **1**, **8b** (340 mg, 1.35 mmol) was treated with NH₃/MeOH (8 h). FC (column 10 × 3 cm, *E*) and crystallization from AcOEt yielded **2** (265 mg, 83%). Colourless crystals. M.p. 122° (dec.). TLC (*F*): R_f 0.6. UV (MeOH): 295 (8700), 264 (sh, 2400), 255 (sh, 1800). ¹H-NMR ((D₆)DMSO): 2.17 (*m*, CH₂(3')); 2.54 (*m*, CH₂(2')); 3.48 (*t*, *J* = 5.6, CH₂(5')); 4.24 (*m*, H-C(4')); 4.75 (*t*, *J* = 5.6, OH-C(5')); 6.47 (*d*, *J* = 4.4, H-C(1')); 8.10 (*s*, NH₂); 8.30 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 45.86, H 5.13, N 35.60.

From **15/17**: As described for **1**, 350 mg (1 mmol) of **15/17** were dissolved in THF and treated with Bu₄NF (1 h). FC afforded **2** (110 mg, 46.6%) from the faster migrating zone.

7-Amino-1-(2,3-dideoxy-β-D-glycero-pentofuranosyl)-1H-1,2,3-triazolo[4,5-d]pyrimidine (3). As described for **1**, **9b** (200 mg, 0.8 mmol) was treated with NH₃/MeOH (36 h). FC (column 110 × 3 cm, *E*) and crystallization from acetone yielded **3** (125 mg, 66.5%). Colourless needles. M.p. 148° (dec.). TLC (*F*): R_f 0.55. UV (MeOH): 289 (7500). ¹H-NMR ((D₆)DMSO): 1.91, 2.13 (2*m*, CH₂(3')); 2.5, 3.09 (2*m*, CH₂(2')); 3.51 (*m*, CH₂(5')); 4.15 (*m*, H-C(4')); 4.89 (*t*, *J* = 5.7, OH-C(5')); 6.74 (*dd*, *J* = 6.3, 1.7, H-C(1')); 7.76 (*s*, NH₂); 8.33 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 45.83, H 5.24, N 35.64.

7-Amino-3-(2,3-dideoxy-α-D-glycero-pentofuranosyl)-3H-1,2,3-triazolo[4,5-d]pyrimidine (18). From **10b**: As described for **1**, **10b** (240 mg, 0.96 mmol) was treated with NH₃/MeOH (24 h). FC (column 10 × 3 cm, *E*) and crystallization from MeOH gave **18** (190 mg, 84%). Colourless needles. Dec. > 170°. TLC (*E*): R_f 0.40. UV (MeOH): 278 (10900). ¹H-NMR ((D₆)DMSO): 1.91, 2.36 (2*m*, CH₂(3')); 2.7–2.5 (*m*, CH₂(2')); 3.46 (*m*, CH₂(5')); 4.33 (*m*, H-C(4')); 4.80 (*t*, *J* = 5.7, OH-C(5')); 6.61 (*dd*, *J* = 7.0, 3.5, H-C(1')); 8.12, 8.45 (2*s*, NH₂); 8.32 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 46.10, H 5.34, N 35.34.

From **16**: Deprotection of **16** (320 mg, 0.91 mmol) was carried out as described for **1**. Crystallization from MeOH yielded **18** (180 mg, 84%).

7-Amino-2-(2,3-dideoxy-α-D-glycero-pentofuranosyl)-2H-1,2,3-triazolo[4,5-d]pyrimidine (19). From **11b**: As described for **1**, **11b** (200 mg, 0.8 mmol) was treated with NH₃/MeOH (7 h). FC (column 10 × 3 cm, *E*) and crystallization from AcOEt yielded **19** (150 mg, 80%). Colourless crystals. M.p. 145° (dec.). TLC (*F*): R_f 0.50. UV (MeOH): 295 (8800), 264 (sh, 2400), 255 (sh, 1800). ¹H-NMR ((D₆)DMSO): 1.94, 2.35 (2*m*, CH₂(3')); 2.55 (*m*, CH₂(2')); 3.49 (*m*, CH₂(5')); 4.43 (*m*, H-C(4')); 4.85 (*t*, *J* = 5.3, OH-C(5')); 6.56 (*dd*, *J* = 7.1, 2.7, H-C(1')); 8.10 (*s*, NH₂); 8.32 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 46.00, H 5.22, N 35.62.

From **15/17**: From the slower migrating zone (see **2**), 95 mg (40.3%) of **19** were obtained.

7-Amino-1-(2,3-dideoxy-α-D-glycero-pentofuranosyl)-1H-1,2,3-triazolo[4,5-d]pyrimidine (20). As described for **1**, **12b** (200 mg, 0.8 mmol) was treated with NH₃/MeOH (36 h). FC (column 10 × 3 cm, *E*) and crystallization from AcOEt gave (135 mg, 72%). Colourless crystals. Dec. > 150°. TLC (*F*): R_f 0.55. UV (MeOH): 289 (7600). ¹H-NMR ((D₆)DMSO): 1.93, 2.16 (2*m*, CH₂(3')); 2.49 (*m*, CH₂(2')); 3.36 (*m*, CH₂(5')); 4.34 (*m*, H-C(4')); 4.76 (*t*, *J* = 5.2, OH-C(5')); 6.62 (*d*, *J* = 4.5, H-C(1')); 7.74 (*s*, NH₂); 8.32 (*s*, H-C(5)). Anal. calc. for C₉H₁₂N₆O₂ (236.23): C 45.76, H 5.12, N 35.58; found: C 46.06, H 5.21, N 35.80.

Glycosylation of 7-Amino-3H-1,2,3-triazolo[4,5-d]pyrimidine (13) [4] with 6. Powdered K₂CO₃ (1.0 g, 7.24 mmol) and TDA-1 (100 μl, 0.31 mmol) were stirred in DMF (100 ml) for 5 min. Then, **13** [4] (1.1 g, 8 mmol) was added and dissolved under warming (60°). The soln. was brought to r.t. and a freshly prepared, cold (–80°) THF soln. (40 ml) of **6** (16 mmol; calculated on the basis of 100% yield of **6**) was added in portions of 5 ml within 30 min. Stirring was continued for another 30 min, insoluble material filtered off, and the filtrate poured into 20% aq. NaHCO₃ soln. (150 ml). The aq. layer was twice extracted with CH₂Cl₂ and the combined org. layer dried (NaSO₄). After evaporation (40°), the residue was dissolved in *G* and applied to the top of a silica-gel column (40 × 3 cm). FC (*G*) gave 3 fractions.

7-Amino-3-{2,3-dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-glycero-pentofuranosyl}-3H-1,2,3-triazolo[4,5-d]pyrimidine (**14**). From *Fr. 2*, **14** (260 mg, 9.3%) was obtained. Colourless solid. TLC (*E*): R_f 0.6. UV (MeOH): 278 (11200), 255 (sh, 6100). $^1\text{H-NMR}$ ($(\text{D}_6$)DMSO): -0.17 , -0.13 , (2s, Me_2Si); 0.74 (s, *t*-Bu); 2.15 – 2.36 (m, $\text{CH}_2(3'')$); 2.72 , 2.6 (2m, $\text{CH}_2(2'')$); 3.60 (m, $\text{CH}_2(5'')$); 4.22 (m, H-C(4'')); 6.52 (dd, $J = 7.2$, 2.2 , H-C(1'')); 8.10 , 8.44 (2s, NH_2); 8.30 (s, H-C(5)). Anal. calc. for $\text{C}_{15}\text{H}_{26}\text{N}_6\text{O}_2\text{Si}$ (350.50): C 51.40, H 7.48, N 23.98; found: C 51.41, H 7.44, N 23.82.

7-Amino-3-{2,3-dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- α -D-glycero-pentofuranosyl}-3H-1,2,3-triazolo[4,5-d]pyrimidine (**16**). Evaporation of *Fr. 1* yielded **16** (240 mg, 8.5%). Colourless solid. TLC (*E*): R_f 0.6. UV (MeOH): 278 (11100), 255 (sh, 6000). $^1\text{H-NMR}$ ($(\text{D}_6$)DMSO): 0.03 , 0.48 (2s, Me_2Si); 0.86 (s, *t*-Bu); 1.93 (m, H-C(3'')); 2.6 (m, $\text{CH}_2(2'')$); 3.65 (m, $\text{CH}_2(5'')$); 4.37 (m, H-C(4'')); 6.59 (dd, $J = 6.7$, 3.1 , H-C(1'')); 8.13 , 8.46 (2s, NH_2); 8.31 (s, H-C(5)). Anal. calc. for $\text{C}_{15}\text{H}_{26}\text{N}_6\text{O}_2\text{Si}$ (350.50): C 51.40, H 7.48, N 23.98; found: C 51.74, H 7.62, N 23.69.

7-Amino-2-{2,3-dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-glycero-pentofuranosyl}-2H-1,2,3-triazolo[4,5-d]pyrimidine (**15**) and 7-Amino-2-{2,3-dideoxy-5-O-[(1,1-dimethylethyl)dimethylsilyl]- α -D-glycero-pentofuranosyl}-2H-1,2,3-triazolo[4,5-d]pyrimidine (**17**). From *Fr. 3*, **15/17** were obtained (390 mg, 13.9%), which could not be separated. For the desilylation and separation of the deprotected compounds, see **2** and **19**.

7-Amino-3-(2,3-dideoxy- β -D-glycero-pentofuranosyl)-3H-1,2,3-triazolo[4,5-d]pyrimidine 5'-[Tetrakis(triethylammonium) Triphosphate] (**21**·4 Et₃N). Compound **1** (47 mg, 0.2 mmol) and *N,N,N',N'*-Tetramethylnaphthalene-1,8-diamine (65 mg, 0.3 mmol) were dissolved in trimethyl phosphate (2 ml) under warming. The soln. was cooled to 0°, freshly distilled POCl₃ (50 μ l, 0.54 mmol) added, and the mixture allowed to stand at 4° for 2 h and then treated with a soln. of tributylammonium diphosphate (0.5N in DMF, 2 ml) and Bu₃N (200 μ l, 0.84 mmol). After stirring for 3 min at 0°, 20 ml of 1M aq. (Et₃NH)HCO₃ (TBK) was added. Evaporation resulted in a semi-solid, which was applied to the top of a DEAE-Sephadex column (30 \times 1.5 cm, HCO₃⁻ form). The elution was performed by a linear gradient of 0.8M TBK buffer (1 l) and H₂O (1 l). The main zone was eluted at 0.52M TBK and yielded **21** (0.11 mmol, 55%). Colourless solid. TLC (*H*): R_f 0.15. UV (H₂O): 280 (11000). $^{31}\text{P-NMR}$ (0.1M Tris-HCl, pH 7.5, 100 nM EDTA/D₂O): -8.32 (d, $J = 19.9$, P(γ)); -10.40 (d, $J = 19.4$, P(α)); -21.97 (t, $J = 19.4$, P(β)).

7-Amino-2-(2,3-dideoxy- β -D-glycero-pentofuranosyl)-2H-1,2,3-triazolo[4,5-d]pyrimidine 5'-[Tetrakis(triethylammonium) Triphosphate] (**22**·4 Et₃N). Compound **2** (23.5 mg, 0.1 mmol) was phosphorylated as described for **21**. After chromatography on DEAE-Sephadex (column 30 \times 1.5 cm, HCO₃⁻ form), **22** was eluted at 0.45M TBK. Evaporation yielded a colourless solid (0.058 mmol, 58%). TLC (*H*): R_f 0.2. UV (H₂O): 294 (8500). $^{31}\text{P-NMR}$ (0.1M Tris-HCl, pH 7.5, 100 nM EDTA/D₂O): -9.98 (d, $J = 19.2$, P(γ)); -10.49 (d, $J = 20.1$, P(α)); -22.34 (t, $J = 19.4$, P(β)).

REFERENCES

- [1] J. A. Montgomery, in 'Handbuch der experimentellen Pharmakologie', Eds. O. Eichler, A. Farha, H. Herken, and A. D. Welch, Springer, Heidelberg, 1974, Vol. 38/1, p. 76.
- [2] J. A. Montgomery, H. J. Thomas, in 'The Purines – Theory and Experiment', 'The Jerusalem Symposia on Quantum Chemistry and Biochemistry 1972', Eds. E. D. Bergmann and B. Pullmann, Academy of Sciences and Humanities, Jerusalem, Israel, Vol. IV, p. 446.
- [3] Y. Mizuno, T. Itoh, A. Nomura, *Heterocycles* **1982**, *17*, 61.
- [4] R. O. Roblin, J. O. Lampen, J. P. English, Q. P. Cole, J. R. Vaughan, *J. Am. Chem. Soc.* **1945**, *67*, 290.
- [5] W. Traube, *Ber. Dtsch. Chem. Ges.* **1900**, *33*, 3035.
- [6] A. Albert, in 'Advances in Heterocyclic Chemistry', Ed. A. R. Katritzky, Academic Press, New York, 1986, Vol. 39, p. 117.
- [7] J. Davoll, *J. Chem. Soc.* **1958**, 1593.
- [8] J. A. Montgomery, H. J. Thomas, *J. Org. Chem.* **1971**, *36*, 1962.
- [9] J. A. Montgomery, H. J. Thomas, S. J. Clayton, *J. Heterocycl. Chem.* **1970**, *7*, 215.
- [10] J. A. Montgomery, R. D. Elliot, in 'Nucleic Acid Chemistry', Eds. L. B. Townsend and R. S. Tipson, Wiley & Sons, New York, 1978, Vol. 68, p. 677.
- [11] J. A. Montgomery, A. T. Shortnacy, G. Arnett, W. M. Shannon, *J. Med. Chem.* **1977**, *20*, 401.
- [12] C. W. Smith, R. W. Sidwell, R. K. Robins, R. L. Tolman, *J. Med. Chem.* **1972**, *15*, 883.
- [13] J. Montgomery, H. J. Thomas, *J. Med. Chem.* **1972**, *15*, 305.
- [14] W. Hutzenlaub, R. L. Tolman, R. K. Robins, *J. Med. Chem.* **1972**, *15*, 879.

- [15] Z. Kazimierczuk, U. Bindig, F. Seela, *Helv. Chim. Acta* **1989**, *72*, 1527.
- [16] E. DeClercq, in 'Aids Research and Human Retroviruses', Mary Ann Liebert, Inc., Publishers, New York, 1992, Vol. 8, p. 119.
- [17] F. Seela, H. Rosemeyer, S. Fischer, *Helv. Chim. Acta* **1990**, *73*, 1602.
- [18] F. Seela, H.-P. Muth, A. Röling, *Helv. Chim. Acta* **1991**, *74*, 554.
- [19] F. Seela, R. Gumbiowski, *Helv. Chim. Acta* **1991**, *74*, 1048.
- [20] F. Seela, H. Rosemeyer, R. Gumbiowski, K. Mersmann, H.-P. Muth, A. Röling, *Nucleos. Nucleot.* **1991**, *10*, 409.
- [21] J. P. H. Verheyden, J. G. Moffat, to *Synthex (USA) Inc.* US 3.817.982, June 18, 1974 (CA: **1974**, *81*, 63942b).
- [22] M. Matthes, M. Janta-Lipinski, K. Reimer, W. Müller, H. Meisel, C. Lehmann, J. Schildt, to *Akademie der Wissenschaften der DDR*, EP 409.227, January 23, 1991 (CA: **1991**, *115*, 159680m).
- [23] E. Kojima, H. Yoshioka, H. Fukinbara, K. Murakami, to *Sanyo-Kokusaku Pulp Co., Ltd.*, Brit. UK Pat. Appl. GB 2.228.479, August 29, 1990 (CA: **1991**, *114*, 102707a).
- [24] W. Fischer, E. Kaun, U. Genz, to *IMA PCT*, Int. Appl. WO 9096.312, June 14, 1990 (CA: **1991**, *114*, 99958a).
- [25] M. Okabe, R.-C. Sun, S. Y.-K. Tam, L. B. Todaro, D. L. Coffen, *J. Org. Chem.* **1988**, *53*, 4780.
- [26] S. Sprang, R. Scheller, D. Rohrer, M. Sundaralingam, *J. Am. Chem. Soc.* **1978**, *100*, 2867.
- [27] H. Rosemeyer, G. Toth, B. Golankiewicz, Z. Kazimierczuk, W. Bourgeois, U. Kretschmer, H.-P. Muth, F. Seela, *J. Org. Chem.* **1990**, *55*, 5784.
- [28] D. Neuhaus, M. Williamson, in 'The Nuclear Overhauser Effect, Structural and Conformational Analysis', Verlag Chemie, Weinheim, 1989, p. 150.
- [29] Z. Kazimierczuk, F. Seela, *Liebigs Ann. Chem.* **1990**, 647.
- [30] T. R. Emerson, R. J. Swan, T. L. V. Ulbricht, *Biochemistry* **1967**, *6*, 843.
- [31] C. A. Bush, in 'Basic Principles in Nucleic Acid Chemistry', Ed. P. O. P. Ts'o, Academic Press, New York, 1974, Vol. 2, p. 129.
- [32] J. L. York, *J. Org. Chem.* **1981**, *46*, 2171.
- [33] R. Käppi, Z. Kazimierczuk, P. Järvinen, F. Seela, H. Lönnberg, *J. Chem. Soc., Perkin Trans. 2* **1991**, 595.
- [34] J. Ludwig, *Acta Biochem. Biophys. Acad. Sci. Hung.* **1981**, *16*, 131.
- [35] F. Seela, 1991, unpublished data.
- [36] H. Ballweg, *Liebigs Ann. Chem.* **1962**, *657*, 141.