

Reaction of γ -Hydroxy-*N*-[1-(dimethylcarbamoyl)ethyl]butanamides under the ‘Direct Amide Cyclization’ Conditions

by Boyan Iliev¹⁾, Anthony Linden, and Heinz Heimgartner*

Organisch-chemisches Institut der Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich


The preparation of the title compounds was achieved *via* the ‘azirine/oxazolone method’ starting from the corresponding γ -hydroxy acids. Upon subjecting the γ -hydroxy-*N*-[1-(dimethylcarbamoyl)ethyl]butanamides **4** to the so-called ‘direct amide cyclization’ (DAC) conditions, chlorinated acids **11** or imino lactones **12** were obtained as the sole products instead of the expected cyclodepsipeptides **A** or their cyclodimers (*Scheme 4*). Variation of the substituents in **4** did not affect the outcome of the reaction and a mechanism for the formation of both products from the intermediate oxazolone **13** has been proposed. Under the acidic conditions of the DAC, the imino lactones are formed as their HCl salts **12**, which, in polar solvents or on silica gel, reacted further to give the chlorinated acids **11**. Stabilization of the imino lactones was achieved by increasing the substitution in the five-membered ring, and their structure, in the form of the hydrochlorides, was established independently by X-ray crystallography (*Fig. 4*). A derivative **15** of the imino lactone **12a** was prepared by the reaction with the 2*H*-azirin-3-amine **10a**; its structure was also established by an X-ray crystal-structure determination (*Fig. 3*). Furthermore, the structures of the ω -chloro acids **11a** and **11b** were determined by X-ray crystallography (*Fig. 2*).

1. Introduction. – Cyclic depsipeptides, *i.e.*, heterodetic cyclopeptides which contain ester (depside) bonds as part of their backbone, have been found in many natural products, and show a wide spectrum of biological activity [1]. They are, therefore, sought after as promising lead compounds for drug design and discovery. Nature is a rich source of fascinating cyclodepsipeptides, and, although the significance of incorporating a depside bond is still not clear, it appears to be essential for biological activity, since all-amide analogues are often inactive [2]. The depside bond is recognized as being more difficult to incorporate into the backbone than the amide bond, although macrolactonizations have been studied extensively [2–6], and is, therefore, usually pre-formed in the linear precursor prior to the cyclization *via* amide-bond formation to give cyclic depsipeptides.

The most-well-known structures in this class of natural products belong to the ion-selective antibiotics, such as valinomycin [7], the closely related enniatin family [8], the actinomycins [9], and others. The reduction in the conformational freedom brought about by cyclization often results in higher-receptor binding affinity. Frequently, in these cyclic compounds, extra conformational restrictions are also built in, such as D-amino acids, *N*-alkylated amino acids or α,α -disubstituted amino acids.

¹⁾ Part of the Ph. D. thesis of *B.I.*, University of Zürich, 2005.

A very efficient method for the incorporation of the latter into depsipeptide rings, the so called ‘direct amide cyclization’ (DAC), has been developed in our research group in recent years. It has been used successfully for the synthesis of 6-, 9-, 12-, and 15-membered cyclodepsipeptides from α -hydroxy acids [10–12], as well as larger ring systems from α - and β -hydroxy acids [13–15]. Therefore, we were interested to investigate reactions of dipeptides, containing β -, γ -, and δ -hydroxy acids, and an α -amino acid, with the aim of synthesizing seven- to nine-membered analogues. As we reported earlier, the cyclization of β -hydroxy acid amides **1**, the linear precursors of the desired seven-membered rings, yielded only cyclodimers **3** by the dimerization process [16][17] (*Scheme 1, Table 1*).




Monosubstitution at C(α) of the amino acid moiety (R³=H) did not prevent dimerization, although the cyclization was carried out under different conditions [17]²). If, however, the hydroxy acid moiety in **1** was monosubstituted at C(α), no cyclic depsipeptides were obtained, although the formation of the intermediate 1,3-oxazol-5(4H)-one **2** has been monitored by IR spectroscopy. Instead, H₂O elimination occurred, to give derivatives of α,β -unsaturated acid amides.

In the present paper, we describe the results of reactions of the higher homologues of **1** containing γ -hydroxy acids, which were carried out with the aim of obtaining either 8- or 16-membered cyclodepsipeptides.

2. Results and Discussion. – The linear amides **4** were synthesized in five steps from the γ,δ -unsaturated aldehydes **5** (*Scheme 2 and Table 1*). Reduction gave the pentenols **6**, which were acetylated to give **7**³). The introduction of the COOH group was achieved by oxidative cleavage of the C=C bond of **7**, either with Ru^{IV}oxide/NaIO₄ [19] or with oxone/acetone [20], although purification proved to be easier in the first case, thus leading to **8** in higher yields. The protection of the OH group prevents the formation of the lactone under these oxidation conditions [21]. The coupling of **8** with the respective

²) The DAC conditions were not applicable in this case because of the sluggish formation of the 1,3-oxazol-5(4H)-one intermediate (*cf.* [18]).

³) The need for protection of the OH group arises from the fact that attempts to obtain the desired γ -hydroxy acids from the corresponding easily available lactones (*e.g.*, 4,4-dimethyltetrahydropyran-2-one) by alkaline hydrolysis failed. Upon acidification of the sodium salt, spontaneous lactonization occurred, and only the starting lactone could be isolated.



a) NaBH_4 , MeOH, 0° , 1 h. b) Ac_2O , Pyr, Et_2O , reflux, 1 h. c) NaIO_4 , Ru^{IV} oxide, MeCN, CCl_4 , H_2O , r.t., 14 h. d) **10**, THF, r.t., 24 h. e) LiOH, THF/ H_2O , r.t., 2 h.

amino acid to give **9** was performed by the reaction with 2,2,*N,N*-tetramethyl- or 2,2,*N*-trimethyl-*N*-phenyl-2*H*-azirine-3-amine (**10a** or **10b**, resp.), and *N,N*-dimethyl-1-azaspiro[2.4]hept-1-en-2-amine (**10c**), respectively, at room temperature ('azirine/oxazolone method' [22][23]). Deprotection of the hydroxy group by treatment with LiOH in THF/ H_2O led to the linear precursors **4**.


Table 1. Synthesized γ -Hydroxy Amides **4**

4	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
a	Me	Me	H	H	Me	Me	Me
b	H	H	H	H	Me	Me	Me
c	Me	Me	H	H	–(CH ₂)–		Me
d	Me	Me	H	H	Me	Me	Ph
e	H	Ph	H	H	Me	Me	Ph
f	Me	Ph	H	H	Me	Me	Ph
g	Me	Me	Me	H	Me	Me	Ph
h	Me	Me	Me	Me	Me	Me	Ph

Unexpectedly, the reactions of **4a–4c** (Table 1) under the conditions of the 'direct amide cyclization' yielded neither the eight-membered nor the 16-membered cyclodepeptides. Instead, after chromatographic workup, the chlorinated acids **11a–11c** were isolated as the sole products (Scheme 3 and Table 2). If the *N*-methyl-*N*-phenylamide **4d** was used instead of the *N,N*-dimethylamides **4a–4c**, the product formed could be isolated after evaporation of the solvent and washing the residue with CH_2Cl_2 . The isolated product was identified as the imino lactone hydrochloride **12a** (Scheme 3).

At first, it seems that R⁵ is determining the product of the reaction. Indeed, it has been shown that in some DAC reactions *N,N*-dimethyl- and *N*-methyl-*N*-phenyl amides behave quite differently [15]. Therefore, the cyclization of **4e** and **4f** was attempted under DAC conditions. Surprisingly, in both cases the product was not the imino lactone of type **12**, but the chlorinated acid of type **11**. Therefore, the presence of the

Scheme 3

Table 2. Synthesized Chlorinated Acids **11**

11	R ¹	R ²	R ⁵	R ⁶
a	Me	Me	Me	Me
b	H	H	Me	Me
c	Me	Me	–(CH ₂) ₄ –	
e	H	Ph	Me	Me
f	Me	Ph	Me	Me

PhN residue can not be the reason for the different behavior of **4d** under DAC conditions.

The explanation lies in the purification process. Since *N*-methylaniline hydrochloride is soluble in CH₂Cl₂ [16][17], and the products of the reaction are not, the crude residue in the case of **4d** was just washed with CH₂Cl₂, and thus the imino lactone was isolated as its hydrochloride **12a** in pure form. On the other hand, Me₂NH·HCl is insoluble in CH₂Cl₂, so that chromatographic workup was necessary in the case of **4a–4c**, which led to the formation of the corresponding chloro acids **11**. In the case of **4e**, and **4f**, due to the presence of the Ph substituent in the hydroxy acid moiety, the product of the reaction was soluble in CH₂Cl₂, and no precipitate was formed. For this reason, chromatographic purification was required leading to chlorinated acids **11** as the sole product (Scheme 3).

It is worth mentioning, that the ¹H-NMR spectrum of **12a** in (D₆)DMSO changes with time. After just 1/2 h at room temperature, the appearance of signals at 1.00, 1.31, 2.09, 3.59, and 8.05 ppm was observed, which grew stronger with time, while the signals at 1.15, 1.56, 3.13, and 4.56 ppm diminished (Fig. 1).

The new signals were sharp and well-defined, indicating the formation of a single compound, rather than decomposition. After ca. 12 h at room temperature, the ratio of both compounds was roughly 1:1, while, after 3 d, only traces of the starting compound **12a** could be observed. The spectrum of the newly formed substance coincides with the spectrum of **11a**. The same process was also observable in the ¹³C-NMR spectrum, the isomerization proceeding in a variety of polar solvents such as DMF and acetone. In the latter, the ratio **12a/11a** reached 1:1 in about a week and then remained constant. Flash chromatography of a mixture **11a/12a** allowed the isolation of analytical amounts of **12a**. Finally, the instability of **12a** in solution was established in a control





Fig. 1. Dynamic ¹H-NMR spectrum of **12a** in (D₆)DMSO: a) spectrum of **12a**, b) spectrum of **11a**

experiment in which a mixture **12a/11a** in CH_2Cl_2 containing 10% of MeOH was stirred at room temperature. After 14 h, only the chlorinated acid **11a** was isolated. Apparently, under the DAC conditions, the initially formed product is the imino lactone hydrochloride **12a**, which, in polar solvents, undergoes an isomerization to give **11a** (Scheme 4).



A reaction mechanism for the formation of **11a** and **12a** is shown in Scheme 4. Treatment of **4d** with HCl gas leads to the corresponding 1,3-oxazol-5(4*H*)-one **13**. When no external nucleophiles are present, the OH group acts as a nucleophile and attacks one of the two electrophilic centres in the ring. If the attack occurs at the C=O group (pathway *a*), the eight-membered cyclodepsipeptide **A** would be formed. On the other hand, the OH attack at the C=N group (pathway *b*) would lead to the opening of the oxazolone ring to yield the imino lactone hydrochloride **12a**. The latter can be isolated if no chromatographic workup is necessary. In a polar solvent, the Cl^- ion apparently attacks the CH_2 group adjacent to the O-atom, which leads to the open-chain ω -chloro acid **11a**⁴⁾.

Due to its instability in solution, no crystals of **12a** suitable for an X-ray crystal structure determination could be obtained, but the structures of the chlorinated acids **11a** and **11b** were confirmed by X-ray crystallography (Fig. 2)⁵⁾.

The OH group in **11a** forms an intermolecular H-bond with the amide O-atom of a neighboring molecule, thereby linking the molecules into extended chains which run parallel to the [0 1 0] direction and can be described by a graph set motif [25] of C(7). The amide group forms an intermolecular H-bond with the carboxylate carbonyl

4) We expected that, in the presence of a better nucleophile, a competition with the attack of Cl^- should be possible, which would lead to a mixture of products, but all reactions in the presence of CsI, Bu_4NI , and PhSn_3 , respectively, led to the formation of **11** exclusively.

5) The structure of **11c** has also been established by an X-ray crystal-structure determination. The quality of the crystals and, subsequently, the results of the structure determination were poor, although unambiguous. The results are not reported here, but have been deposited at the CCDC (see Footnote 6).





Fig. 2. ORTEP Plots [24] of the molecular structures of a) **11a** and b) one of the disordered conformations of one of the two symmetry-independent molecules of **11b** (arbitrary numbering of the atoms; 50% probability ellipsoids)

O-atom of a different neighboring molecule, thus also linking the molecules into extended chains. These chains run parallel to the $[1\ 0\ 0]$ direction and can be described by a graph set motif of C(5). The combination of both interactions generates a two-dimensional network, which lies parallel to the $(0\ 0\ 1)$ plane.

In the case of **11b**, there are two symmetry-independent molecules in the asymmetric unit. The most significant difference between the independent molecules is in the orientation of the Cl-atom. Furthermore, each of the two symmetry-independent molecules is disordered over the $\text{Cl}(\text{CH}_2)_3$ section of the molecule. Two positions were defined for these atoms in each molecule, except for the Cl-substituted C-atom, which is common to both conformations. The major conformation is present in *ca.* 65 and 80% of molecules A and B, respectively. Except for the orientations of the Cl-atoms, the major conformation of molecule A is almost identical to that of the minor conformation of molecule B and *vice versa*. The OH group of each molecule in **11b** forms an intermolecular H-bond with the amide O-atom of an adjacent molecule of the same type. These interactions link the molecules into extended chains composed entirely of molecules of type A or entirely of type B. These chains run in the $[1\ 0\ 1]$ direction and can be described by a graph set motif of C(7). The amide group of each molecule forms an intermolecular H-bond with the carbonyl O-atom of the carboxy group of an adjacent symmetry-independent molecule. These interactions link

the molecules into extended $\cdots A \cdots B \cdots A \cdots B \cdots$ chains which run in the $[0\ 0\ 1]$ direction and can be described by a binary graph set motif of $C_2^2(10)$. The combination of all H-bonding interactions links the molecules into extended two-dimensional networks which lie parallel to the $(0\ 1\ 0)$ plane.

To isolate the crucial intermediate **12a**, extraction with aq. Na_2CO_3 solution [26] and deprotection by treatment with polyvinylpyridine were attempted, but neither procedure gave satisfactory results. Although esterification of **12a** with CH_2N_2 yielded the corresponding crude methyl ester hydrochloride **14a** (Scheme 5), the product was also not stable in polar solvents and isomerized partially to the corresponding chloro ester.



Finally, the reaction of **12a** with *2H*-azirin-3-amine **10b** yielded the diamide **15**, which was stable in solution, and whose structure was confirmed by X-ray crystallography (Fig. 3). This is the first direct proof of the tetrahydrofuran-2-imine structure.





Fig. 3. ORTEP Plot [24] of the molecular structure of **15** (arbitrary numbering of the atoms; 50% probability ellipsoids)

The central amide group (N(4)H) of **15** has a weak intramolecular H-bonding interaction with the imine N-atom (N(7)), which results in a five-membered loop that can be described by a graph set motif [25] of $S(5)$.

When the hydroxy-protected amide **9d** was subjected to the DAC conditions, the corresponding protected 1,3-oxazol-5(4*H*)-one **16a** (Scheme 6) was isolated in good yield, which indicated that an oxazolone, *i.e.*, **13** in Scheme 4, is most probably the first intermediate from which the imino lactones **12** and the chloro acids **11** are formed. Therefore, we decided to prepare **9i–9l** with different protecting groups, which could potentially allow the deprotection and isolation of the oxazolone with a free OH group as a precursor for the cyclization to give **A**.



We planned to prepare **9i–9l** analogously to **9d** (Scheme 2). The benzyl (Bn), (benzyloxy)methyl (Bom), and (9*H*-fluoren-9-yl)methoxycarbonyl (Fmoc) protecting groups could be introduced before the oxidative cleavage of the double bond and yield the protected acids of type **8**, which could then be reacted with 2*H*-azirin-3-amine **10b** to give the protected diamides **9**. Because of the instability of the protecting group under the oxidation conditions, direct protection of the hydroxy amide **4d** was preferred (Scheme 6). Unfortunately, the deprotection of **16b–16d** failed in all cases.



Since the DAC method failed to give cyclic depsipeptides, we attempted to synthesize the hydroxy esters of type **17** from amides **4** and to subject them to the NaH cyclization procedure, as described earlier [16][17]. Unexpectedly, treatment of **4d** and **4f** with HCl gas in toluene containing 20% EtOH as an external nucleophile did not yield the desired esters **17**, but gave as the main products the imino lactone esters **14** in moderate yields together with the corresponding *ω*-chloro esters **18** (Scheme 7).


Further experiments with **4d** in the presence of MeOH showed that the ester could also be formed directly from the imino lactone **12a**, either under DAC conditions or by treatment with CH_2N_2 (Schemes 5 and 7).

If instead of the hydroxy amides **4**, the protected alcohols **9d** or **9f** were subjected to the DAC conditions in the presence of MeOH, the unprotected linear esters **17** were formed (Scheme 8). We believe that again oxazolones are the intermediates, and, since the OH group in **9** is protected, it is to be expected that protected oxazolones of type **16** (Scheme 6) are the intermediates. Nucleophilic addition of MeOH to the




oxazolone, followed by ring opening, leads to the ester group in **17**. The deacetylation could be explained by trans-esterification, *via* formation of AcOMe, which is a common reaction under acidic conditions.

The esters **17** were subjected to the conditions of the NaH-catalyzed cyclization, *i.e.*, they were treated with NaH in toluene at 80° [16][27], but no cyclic products were obtained. As in previous studies [16][17], the rigidity of the amide bond was considered to be the reason for the failure. Therefore, the amino ketone **19** was synthesized, as depicted in Scheme 9, and subjected to the same reaction conditions. Again, no cyclic products could be identified in the mixture.



a) HCHO, NaOH; 72%. b) Br₂; 88%. c) Ethyl 2-amino-2-methylpropanoate hydrochloride, Et₃N; 39%.

If indeed the formation of the cyclic products **12** and **14** occurs *via* the attack of the OH group at the iminium C-atom of the oxazolone **12** (see *Scheme 4*, path *b*), α -substitution in the hydroxy diamide **4** might present sufficient steric hindrance in order to direct the nucleophilic attack onto the ester group (*Scheme 4*, path *a*). Therefore, the amides **4g** and **4h** were prepared analogously to *Scheme 2*. Treatment of amides **4g** and **4h** with HCl gas in toluene at 100° (DAC conditions) yielded imino lactones as sole products (*Scheme 10*).



Unlike their analogues **12a**–**12c**, compounds **12g** and **12h** are stable in solution. The higher substitution of the ring apparently stabilizes it, *i.e.*, the gem-dimethyl effect, (*Thorpe–Ingold* effect [28]) makes the ring opening thermodynamically unfavorable. Thanks to their stability in solution, crystals suitable for an X-ray crystal structure determination could be grown for both hydrochlorides **12g** and **12h** (*Fig. 4*).

Since the space group of **12g** is centrosymmetric, the compound in the crystal is racemic. The OH group forms a H-bond with a neighboring Cl^- ion, while the iminium group forms a H-bond with a different Cl^- ion. Thus, each Cl^- ion accepts two H-bonds. These interactions link two cations and two anions in a cyclic manner into a tetrameric unit and can be described by a graph set motif [25] of $R_4^2(14)$.

Similar H-bonding interactions occur in **12h**, but this time the interactions link the ions into extended chains, which run parallel to the [1 0 0] direction and can be described by a graph set motif of $C_2^1(7)$.

Conclusions. – When γ -hydroxy diamides **4** were subjected to the DAC reaction conditions, neither the 8-membered nor the 16 membered depsipeptides were formed. Depending on the workup procedure, either the chlorinated acids **11** or the imino lactone hydrochlorides **12** with a COOH group were obtained as the sole products in good yields. Both products are formed *via* the intermediate oxazolones by an attack of the OH group at the iminium C-atom, instead of at the C=O group of the oxazolone (*Scheme 4*). This attack leads to the five-membered ring, instead of the desired eight-membered ring. The former are obviously more stable than the latter.

Compounds **12**, with α -H atoms next to the C=O group, are unstable in solution and isomerize to the corresponding ω -chloro acids **11**. Increased substitution of the imino lactone stabilizes the five membered ring and prevents isomerization. A few other cyclization methods were also tried, but they all failed to give cyclic depsipeptides.




Fig. 4. ORTEP Plots [24] of the molecular structures of a) **12g** and b) **12h** (arbitrary numbering of the atoms; 50% probability ellipsoids)

Experimental Part

1. *General.* See [16].

2. *Starting Materials.* 2,2,N,N-Tetramethyl-2H-azirin-3-amine (**10a**), 2,2,N-trimethyl-N-phenyl-2H-azirin-3-amine (**10b**), and N,N-dimethyl-1-azaspiro[2.4]hept-1-en-2-amine (**10c**) were prepared according to standard procedures (*cf.* [16] and refs. cit. therein). 2,2-Dimethylpent-4-enal (**5a**) was prepared from isobutyraldehyde according to *Noack* and *Goettlich* [29], 2-phenylpent-4-en-1-ol (**6e**) was obtained from 2-phenylpropanal by following the method of *Iqbal* and *Srivastava* [30], and 2-methyl-2-phenylpent-4-enal (**5f**) was synthesized from 2-phenylpropanal analogously to **5a**. All spectra were in accord-

ance with literature data [31]. 2,2,3-Trimethylpent-4-enal (**5g**) and 2,2,3,3-tetramethylpent-4-enal (**5h**) were prepared according to *Brannock* and *Burpiit* [32] from crotyl bromide and 1-bromo-3-methylbut-2-ene, resp. 1-Bromo-4-hydroxy-3,3-dimethylbutan-2-one was synthesized according to *Mihelcic* and *Moeller* [33]. All other products used were commercially available.

3. *Preparation of Pent-4-enyl Acetates. General Procedure 1 (GP 1)*. To a soln. of the corresponding pent-4-enal **5** (5 mmol) in MeOH (20 ml) at 0°, NaBH₄ (760 mg, 20 mmol) was added in small portions within 20 min, then the mixture was allowed to warm to r.t. and was stirred for a total of 1 h. The solvent was evaporated *i.v.*, and the residue was dissolved in H₂O. Extraction with Et₂O (5 × 30 ml), drying (MgSO₄), and evaporation *i.v.* yielded the pent-4-en-1-ols **6**, which were acetylated without further purification. For this purpose, **6** (5 mmol) was dissolved in Et₂O (50 ml) and pyridine (0.81 ml, 10 mmol) was added. The mixture was heated to reflux, and a soln. of Ac₂O (0.41 ml, 5.5 mmol) in Et₂O (10 ml) was added dropwise within 20 min. The mixture was stirred under reflux for another 1.5 h, cooled, the salt was filtered off, the org. layer was washed with 10% aq. CuSO₄ soln. and brine, dried (MgSO₄), evaporated *i.v.*, and purified by column chromatography (CC, SiO₂) to yield the desired pent-4-enyl acetates **7**.

3.1. *2,2-Dimethylpent-4-enyl Acetate (7a)*. According to *GP 1* from 2,2-dimethylpent-4-enal (**5a**, 560 mg, 5 mmol), CC (hexane/Et₂O 10 : 1). Yield: 593 mg, (79%) of **7a** as a colorless oil. All spectra were in accordance with literature data [34].

3.2. *2-Phenylpent-4-enyl Acetate (7e)*. To a soln. of 2-phenylpent-4-en-1-ol (**6e**, 5 mmol, 810 mg) in Et₂O (50 ml), pyridine (0.81 ml, 10 mmol) was added, the mixture was heated to reflux, and then a soln. of Ac₂O (0.71 ml, 5.5 mmol) in Et₂O (10 ml) was added dropwise over 20 min. The mixture was stirred under reflux for another 1.5 h, cooled, and the pyridinium acetate was filtered off, the org. layer was washed with 10% aq. CuSO₄ soln. and brine, dried (MgSO₄), evaporated *i.v.*, and purified by CC (hexane/Et₂O 15 : 1) to yield 860 mg (84%) of **7e**. Colorless oil. ¹H-NMR: 1.98 (*s*, MeCO); 2.24–2.41 (*m*, CH₂); 2.91–3.06 (*m*, PhCH); 4.11–4.26 (*m*, CH₂O); 4.92–5.09 (*m*, CH₂=CH); 5.62–5.83 (*m*, CH₂=CH); 7.12–7.39 (*m*, 5 arom. H). ¹³C-NMR: 20.8 (*q*, Me); 36.8 (*t*, CH₂); 44.4 (*d*, CH); 67.6 (*t*, CH₂O); 116.6 (*t*, CH₂=CH); 126.7, 127.4, 128.3 (*3d*, 5 arom. CH); 135.6 (*d*, CH₂=CH); 141.3 (*s*, arom. C); 170.8 (*s*, C=O). ESI-MS: 205 (100, [M+H]⁺).

3.3. *2-Methyl-2-phenylpent-4-enyl Acetate (7f)*. According to *GP 1* from 2-methyl-2-phenylpent-4-enal (**5f**; 870 mg, 5 mmol), CC (hexane/Et₂O 20 : 1). Yield: 785 mg (72%) of **7f**. Colorless oil. IR: 3075s, 2962vs, 2860vs, 1745vs, 1636s, 1482s, 1432s, 1386s, 1334s, 1188s, 1076s, 1029s, 912m. ¹H-NMR: 1.35 (*s*, Me); 2.05 (*s*, MeCO); 2.31–2.59 (*m*, CH₂); 4.13–4.28 (*m*, CH₂O); 4.87–5.11 (*m*, CH₂=CH); 5.45–5.59 (*m*, CH₂=CH); 7.14–7.44 (*m*, 5 arom. H). ¹³C-NMR: 20.7, 22.5 (*2q*, 2 Me); 41.0 (*s*, C(Me)(Ph)); 43.4 (*t*, CH₂); 71.4 (*t*, CH₂O); 117.8 (*t*, CH₂=CH); 126.1, 127.7, 128.1 (*3d*, 5 arom. CH); 133.3 (*d*, CH₂=CH); 144.3 (*s*, arom. C); 170.9 (*s*, C=O). ESI-MS: 219 (100, [M+H]⁺).

3.4. *2,2,3-Trimethylpent-4-enyl Acetate (7g)*. According to *GP 1*, from 2,2,3-trimethylpent-4-enal (**5g**, 5 mmol, 630 mg), CC (hexane/Et₂O 10 : 1). Yield: 604 mg (71%) of **7g**. Colorless oil. IR: 3077w, 2973vs, 2879s, 1744vs, 1637m, 1473m, 1377s, 1036s, 914m. ¹H-NMR: 0.86, 0.90 (2s, Me₂C); 0.98 (*d*, *J* = 4.5, Me); 1.98 (*s*, MeCO); 2.11–2.23 (*m*, MeCH); 3.36 (*s*, CH₂O); 4.90–5.04 (*m*, CH₂=CH); 5.74–5.92 (*m*, CH₂=CH). ¹³C-NMR: 14.6, 20.8, 21.6 (*3q*, Me, Me₂C, MeCO); 36.5 (*s*, Me₂C), 40.9 (*d*, MeCH); 70.4 (*t*, CH₂O); 114.1 (*t*, CH₂=CH); 141.5 (*d*, CH₂=CH); 171.8 (*s*, C=O). ESI-MS: 193 (100, [M+H]⁺).

3.5. *2,2,3,3-Tetramethylpent-4-enyl Acetate (7h)*. According to *GP 1*, from 2,2,3,3-tetramethylpent-4-enal (**5h**, 5 mmol, 700 mg), CC (hexane/Et₂O 10 : 1). Yield: 717 mg (78%) of **7h**. Colorless oil. IR: 3072m, 2964vs, 2909s, 1746vs, 1635m, 1473m, 1414m, 1380s, 1315s, 1040s, 913s. ¹H-NMR: 0.88, 0.99 (2s, 2 Me₂C); 2.02 (*s*, MeCO); 3.90 (*s*, CH₂O); 4.89–5.00 (*m*, CH₂=CH); 5.87–5.96 (*m*, CH₂=CH). ¹³C-NMR: 20.5 (*q*, Me₂C); 21.0 (*q*, MeCO); 22.6 (*q*, Me₂C); 38.3, 40.8 (2s, 2 Me₂C); 70.6 (*t*, CH₂O); 112.0 (*t*, CH₂=CH); 145.3 (*d*, CH₂=CH); 171.3 (*s*, C=O). ESI-MS: 207 (100, [M+H]⁺).

4. *4-Acetoxybutanoic Acids 8. General Procedure 2 (GP 2)*. To a soln. of pent-4-enyl acetates **7** (5 mmol) in MeCN/CCl₄/H₂O 2 : 2 : 3 (70 ml), NaIO₄ (2.28 g, 20 mmol) was added under stirring. After 10 min, a cat. amount of RuO₂·H₂O was added, and the mixture was stirred vigorously at r.t. for 4–12 h. Filtration of the white residue over *Celite*, washing with CH₂Cl₂, extraction of the collected mother liquor with CH₂Cl₂, drying (MgSO₄), evaporation *i.v.*, and purification by CC yielded **8** as colorless oils.

4.1. *4-Acetoxy-3,3-dimethylbutanoic Acid (8a)*. According to *GP 2* from **7a** (5 mmol, 780 mg), 6 h, CC (CH₂Cl₂/MeOH 20 : 1). Yield: 625 mg (73%) of **8a**. Colorless oil. IR: 3284s (br), 2969s, 1739vs,

1709vs, 1475w, 1379s, 1243s, 1041s, 926w. ¹H-NMR: 0.98 (s, Me₂C); 1.98 (s, MeCO); 2.28 (s, CH₂); 3.88 (s, CH₂O); 9.83 (br. s, COOH). ¹³C-NMR: 20.6 (q, Me); 24.9 (q, Me₂C); 33.5 (s, Me₂C); 42.8 (t, CH₂); 71.6 (t, CH₂O); 171.1 (s, C=O); 177.7 (s, COOH). ESI-MS: 197 (100, [M+Na]⁺).

4.2. 4-Acetoxy-3-phenylbutanoic Acid (**8e**). According to GP 2, **7e** (5 mmol, 1.020 g), 4 h, CC (CH₂Cl₂/acetone 20:1). Yield: 678 mg (61%) of **8e**. Colorless oil. ¹H-NMR: 1.99 (s, MeCO); 2.61–2.88 (m, CH₂); 3.42–3.58 (m, PhCH); 4.09–4.36 (m, CH₂O); 7.13–7.38 (m, 5 arom. H.); 10.63 (br. s, COOH). ¹³C-NMR: 20.6 (q, MeCO); 37.1 (t, CH₂); 40.7 (d, CH); 67.3 (t, CH₂O); 127.2, 127.5, 128.6 (3d, 5 arom. CH); 139.8 (s, arom. C); 170.7 (s, C=O); 177.3 (s, COOH). ESI-MS: 245 (100, [M+Na]⁺).

4.3. 4-Acetoxy-3-methyl-3-phenylbutanoic Acid (**8f**). According to GP 2 from **7f** (5 mmol, 1090 mg), 6 h, CC (CH₂Cl₂/MeOH 20:1). Yield: 861 mg (73%) of **8f**. Colorless oil. ¹H-NMR: 1.31 (s, Me); 2.01 (s, MeCO); 2.28–2.36 (m, CH₂); 4.18–4.45 (m, CH₂O); 7.08–7.38 (m, 5 arom. H.). ¹³C-NMR: 20.9 (q, MeCO); 26.6 (q, Me); 31.1 (s, Me₂C); 45.7 (d, CH); 69.3 (t, CH₂O); 127.3, 127.6, 128.9 (3d, 5 arom. CH); 141.2 (s, arom. C); 171.4 (s, C=O); 176.1 (s, COOH). ESI-MS: 259 (100, [M+Na]⁺).

4.4. 4-Acetoxy-2,3,3-trimethylbutanoic Acid (**8g**). According to GP 2 from **7g** (5 mmol, 850 mg), 6 h, CC (CH₂Cl₂/MeOH 20:1). Yield: 771 mg (82%) of **8g**. Colorless oil. IR: 3466s (br.), 2976s, 1738vs, 1710vs, 1467w, 1390s, 1245s, 1041s, 925w. ¹H-NMR: 0.95, 0.99 (2s, Me₂C); 1.18 (d, J=4.7, MeCH); 2.00 (s, MeCO); 2.48 (q, J=4.7, MeCH); 3.89 (s, CH₂O); 10.11 (br. s, COOH). ¹³C-NMR: 15.6, 20.6 (2q, 2 Me); 21.8, 22.1 (2q, Me₂C); 38.7 (s, Me₂C); 45.1 (d, CH); 70.8 (t, CH₂O); 171.0 (s, C=O); 181.3 (s, COOH). ESI-MS: 211 (100, [M+Na]⁺).

4.5. 4-Acetoxy-2,2,3,3-tetramethylbutanoic Acid (**8h**). According to GP 2 from **7h** (5 mmol, 920 mg), 4 h, CC (CH₂Cl₂/MeOH 20:1). Yield: 524 mg (52%) of **8h**. IR: 3290s (br.), 2972s, 1743vs, 1477m, 1377m, 1248s, 1214w, 1040s, 1021w, 926w. ¹H-NMR: 0.99, 1.19 (2s, 2 Me₂C); 2.03 (s, MeCO); 3.99 (s, CH₂O); 9.63 (br. s, COOH). ¹³C-NMR: 20.6 (q, MeCO); 21.1, 21.4 (2s, 2 Me₂C); 38.3, 46.9 (2s, 2 Me₂C); 70.1 (t, CH₂O); 171.1 (s, C=O); 183.3 (s, COOH). CI-MS: 203 (38, [M+H]⁺), 143 (100, [M-MeCO]⁺).

5. Coupling of **8** with 2H-Azirin-3-amines **10**. General Procedure 3 (GP 3). Acids **8** were taken up in dry THF (20 ml), and the corresponding aminoazirine **10** was added dropwise. The mixture was stirred overnight at r.t., the solvent was evaporated *i.v.*, and the residue was purified by CC to yield acetoxy diamides **9**.

5.1. 3-[[1-Methyl-1-(N,N-dimethylcarbamoyl)ethyl]carbamoyl]-2,2-dimethylpropyl Acetate (**9a**). According to GP 3 from **8a** (348 mg, 2 mmol) in dry THF (20 ml) and **10a** (235 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 30:1). Yield: 509 mg (89%) of **9a**. White powder. M.p. 99.1–99.2°. ¹H-NMR: 0.98, 1.52 (2s, 2 Me₂C); 2.05 (s, MeCO); 2.10 (s, CH₂); 3.08 (br. s, Me₂N); 3.98 (s, CH₂O); 7.12 (s, NH). ¹³C-NMR: 20.9 (q, MeCO); 24.5, 24.7 (2q, 2 Me₂C); 34.1 (s, Me₂C); 38.2 (q, Me₂N); 45.5 (t, CH₂); 56.8 (s, Me₂C); 71.8 (t, CH₂O); 169.4, 171.2, 173.2 (3s, 3 C=O). ESI-MS: 309 (100, [M+Na]⁺).

5.2. 3-[[1-(N,N-Dimethylcarbamoyl)cyclopropyl]carbamoyl]-2,2-dimethylpropyl Acetate (**9c**). According to GP 3 from **8a** (348 mg, 2 mmol) in dry THF (20 ml) and **10c** (290 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 20:1). Yield: 549 mg (88%) of **9c**. White powder. M.p. 139.0–140.2°. ¹H-NMR: 1.02 (s, Me₂C); 1.66–1.78, 1.83–1.93 (2m, 4 CH₂); 2.07 (s, MeCO); 2.18–2.26 (m, CH₂); 2.98 (br. s, Me₂N); 3.96 (s, CH₂O); 6.41 (br. s, NH). ¹³C-NMR: 21.0 (q, MeCO); 22.4 (t, CH₂); 24.6 (q, Me₂C); 34.3 (t, CH₂); 35.7 (s, Me₂C); 38.2 (q, Me₂N); 44.3 (t, CH₂); 66.8 (s, Me₂C); 71.9 (t, CH₂O); 169.8, 171.4, 172.5 (3s, 3 C=O). CI-MS: 313 (40, [M+H]⁺), 268 (100, [M-Me₂N]⁺).

5.3. 3-[[1-Methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]-2,2-dimethylpropyl Acetate (**9d**). According to GP 3 from **8a** (348 mg, 2 mmol) in dry THF (20 ml) and **10b** (365 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 40:1). Yield: 584 mg (84%) of **9d**. White powder. M.p. 94.1–95.8°. ¹H-NMR: 0.99, 1.48 (2s, 2 Me₂C); 2.03 (s, MeCO); 2.12 (s, CH₂); 3.26 (s, MeN); 3.91 (s, CH₂O); 6.38 (s, NH); 7.21–7.43 (m, 5 arom. H.). ¹³C-NMR: 20.9 (q, MeCO); 24.4, 26.0 (2q, 2 Me₂C); 34.0 (s, Me₂C); 41.5 (q, MeN); 46.0 (t, CH₂); 58.3 (s, Me₂C); 71.7 (t, CH₂O); 127.9, 128.2, 129.3 (3d, 5 arom. CH); 144.5 (s, arom. C); 169.6, 171.2, 173.4 (3s, 3 C=O). ESI-MS: 371 (100, [M+Na]⁺).

5.4. 3-[[1-Methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]-2-phenylpropyl Acetate (**9e**). According to GP 3 from **8e** (444 mg, 2 mmol) in dry THF (20 ml) and **10b** (365 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 50:1). Yield: 689 mg (87%) of **9e**. White powder. M.p. 133.3–134.2°. ¹H-NMR: 1.32, 1.38 (2s, Me₂C); 2.09 (s, MeCO); 2.18–2.25, 2.43–2.50 (2m, CH₂); 3.20–3.26 (m, MeN, CH); 3.72–3.77 (m, CH₂O); 5.97 (s, NH); 7.17–7.36 (m, 10 arom. H.). ¹³C-NMR: 21.1 (q, MeCO); 26.1, 26.2

(2*q*, Me₂C); 40.9 (*t*, CH₂); 41.3 (*q*, MeN); 44.5 (*d*, CH); 58.1 (*s*, Me₂C); 67.0 (*t*, CH₂O); 126.9, 127.5, 127.9, 128.6, 129.3 (5*d*, 10 arom. CH); 141.8, 144.4 (2*s*, 2 arom. C); 169.9, 171.0, 173.0 (3*s*, 3 C=O). ESI-MS: 419 (100, [M+Na]⁺).

5.5. 3-*[[1-Methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]-2-methyl-2-phenylpropyl Acetate (9f)*. According to GP 3 from **8f** (472 mg, 2 mmol) in dry THF (20 ml) and **10b** (365 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 50:1). Yield: 689 mg (84%) of **9f**. White powder. M.p. 135.1–136.6°. ¹H-NMR: 1.08, 1.16, 1.36 (3*s*, Me, Me₂C); 2.08 (*s*, MeCO); 2.11–2.17 (*m*, CH₂); 3.16 (*s*, MeN); 4.21 (*s*, CH₂O); 5.64 (*s*, NH); 7.09–7.41 (*m*, 10 arom. H). ¹³C-NMR: 20.8 (*q*, MeCO); 22.4, 26.1, 26.6 (3*q*, Me, Me₂C); 30.8 (*s*, Me₂C); 41.1 (*q*, MeN); 46.3 (*t*, CH₂); 57.4 (*s*, Me₂C); 71.1 (*t*, CH₂O); 126.1, 127.6, 127.9, 128.5, 128.9, 129.3 (6*d*, 10 arom. CH); 143.6, 144.8 (2*s*, 2 arom. C); 168.9, 170.8, 172.9 (3*s*, 3 C=O). CI-MS: 411 (42, [M+H]⁺), 304 (100, [M–Me(Ph)N]⁺), 175 (21).

5.6. 3-*[[1-Methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]-2,2-dimethylbutyl Acetate (9g)*. According to GP 3 from **8g** (372 mg, 2 mmol) in dry THF (20 ml) and **10b** (365 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 50:1). Yield: 637 mg (88%) of **9g**. White powder. M.p. 139.6–140.9°. ¹H-NMR: 0.99, 1.00 (2*s*, Me₂C); 1.05 (*d*, *J*=3.9, MeCH); 1.40, 1.41 (2*s*, Me₂C); 2.05–2.12 (*m*, MeCO, CH); 3.19 (*s*, MeN); 3.86 (*q*, CH₂O); 6.66 (*s*, NH); 7.14–7.46 (*m*, 5 arom. H). ¹³C-NMR: 12.2 (*q*, Me); 20.8 (*q*, MeCO); 21.6, 22.4, 24.8, 25.0 (4*q*, 2 Me₂C); 35.9 (*s*, Me₂C); 41.4 (*q*, MeN); 46.4 (*d*, CH); 58.5 (*s*, Me₂C); 71.3 (*t*, CH₂O); 128.0, 128.4, 129.3 (3*d*, 5 arom. CH); 144.1 (*s*, arom. C); 170.9, 172.7, 173.7 (3*s*, 3 C=O). ESI-MS: 385 (100, [M+Na]⁺).

5.7. 3-*[[1-Methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]-2,2,3-trimethylbutyl Acetate (9h)*. According to GP 3 from **8h** (404 mg, 2 mmol) in dry THF (20 ml) and **10b** (365 mg, 2.1 mmol), CC (CH₂Cl₂/MeOH 50:1). Yield: 684 mg (91%) of **9g**. White powder. M.p. 119.9–120.9°. ¹H-NMR: 0.93, 1.11, 1.47 (3*s*, 3 Me₂C); 2.04 (*s*, MeCO); 3.26 (*s*, MeN); 3.98 (*s*, CH₂O); 7.04 (*s*, NH); 7.18–7.41 (*m*, 5 arom. H). ¹³C-NMR: 21.0, 21.7, 24.9 (3*q*, 3 Me₂C); 38.6 (*s*, Me₂C); 41.7 (*q*, MeN); 47.2, 58.8 (2*s*, 2 Me₂C); 70.7 (*t*, CH₂O); 128.3, 128.6, 129.5 (3*d*, 5 arom. CH); 144.1 (*s*, arom. C); 171.1, 174.3, 176.9 (3*s*, 3 C=O). ESI-MS: 415 (20, [M+K]⁺), 399 (100, [M+Na]⁺), 377 (60, [M+H]⁺). Anal. calc. for C₂₁H₃₂N₂O₄ (376.50): C 66.99, H 8.57, N 7.44; found: C 66.35, H 8.28, N 7.15.

6. *Deprotection of 9 to Give Hydroxy-Diamides 4. General Procedure 4 (GP 4)*. The amides **9** were treated with 4 equiv. of LiOH in THF/H₂O 2:1 at r.t. for 4–12 h. Evaporation of the solvent *i.v.*, extraction of the residue with CH₂Cl₂, drying (MgSO₄), evaporation *i.v.*, and washing with Et₂O yielded the hydroxy diamides **4** as white powders, which were used without further purification.

6.1. 4-Hydroxy-3,3-dimethyl-N-*[1-methyl-1-(N,N-dimethylcarbamoyl)ethyl]butanamide (4a)*. According to GP 4 from **9a** (858 mg, 3 mmol), 4 h. Yield: 677 mg (93%) of **4a**. White powder. M.p. 138.6–139.9°. ¹H-NMR: 0.99, 1.64 (2*s*, Me₂C); 2.14 (*s*, CH₂); 3.12 (br. *s*, Me₂N); 3.91 (*s*, CH₂O); 7.02 (*s*, NH). ¹³C-NMR: 24.6, 24.8 (2*q*, 2 Me₂C); 34.6 (*s*, Me₂C); 38.8 (*q*, Me₂N); 45.4 (*t*, CH₂); 56.8 (*s*, Me₂C); 71.3 (*t*, CH₂O); 171.4, 173.0 (2*s*, 2 C=O). CI-MS: 244 (32, [M+H]⁺), 201 (100, [M–Me₂N]⁺).

6.2. 4-Hydroxy-N-*[1-methyl-1-(N,N-dimethylcarbamoyl)ethyl]butanamide (4b)*. A soln. of sodium 4-hydroxybutanoate (1.0 g, 7.9 mmol) in H₂O (10 ml) was acidified with 3*N* HCl. Extraction with AcOEt (5 × 30 ml), drying (MgSO₄), and evaporation of the solvent *i.v.* yielded 520 mg (63%) of 4-hydroxybutanoic acid as a viscous fluid, which was used without further purification. All spectra were in accordance with the data in [35]. 4-Hydroxybutanoic acid (600 mg, 5.77 mmol) was dissolved in dry THF (20 ml) and **10a** (711 mg, 6.35 mmol, 1.1 equiv.) was added dropwise. The mixture was stirred overnight at r.t., the solvent evaporated *i.v.*, and the residue was purified by CC (CH₂Cl₂/MeOH 10:1). Yield: 1134 mg (90%) of **4b**. White solid. M.p. 124.6–125.3°. ¹H-NMR: 1.48 (*s*, Me₂C); 1.73–1.82 (*m*, CH₂); 2.08 (*t*, *J*=6.2, CH₂); 3.26 (*s*, Me₂N); 3.65 (*t*, *J*=6.2, CH₂O); 6.38 (*s*, NH); 7.24–7.32 (*m*, 5 arom. H). ¹³C-NMR: 24.4 (*t*, CH₂); 26.0 (*q*, Me₂C); 41.5 (*q*, Me₂N); 46.0 (*t*, CH₂); 58.3 (*s*, Me₂C); 71.7 (*t*, CH₂O); 127.9, 128.2, 129.3 (3*d*, 5 arom. CH); 144.5 (*s*, arom. C); 171.2, 173.4 (2*s*, 2 C=O). CI-MS: 279 (85, [M+H]⁺), 172 (100, [M–Me₂N]⁺).

6.3. 4-Hydroxy-3,3-dimethyl-N-*[1-(N,N-dimethylcarbamoyl)cyclopentyl]butanamide (4c)*. According to GP 4 from **9c** (936 mg, 3 mmol), 6 h. Yield: 738 mg (91%) of **4c**. White powder. M.p. 138.4–139.9°. ¹H-NMR: 0.99 (*s*, Me₂C); 1.62–1.76, 1.84–2.02 (2*m*, 4 CH₂); 2.21–2.39 (*m*, CH₂); 3.03 (br. *s*, Me₂N); 3.38 (*s*, CH₂O); 6.61 (br. *s*, NH). ¹³C-NMR: 24.4 (*t*, CH₂); 25.2 (*q*, Me₂C); 35.7 (*s*, Me₂C); 37.3 (*t*, CH₂); 37.9 (*q*, Me₂N); 46.4 (*t*, CH₂); 66.6 (*s*, C); 71.9 (*t*, CH₂O); 172.2, 172.7 (2*s*, 2 C=O). ESI-MS: 309 (10, [M+K]⁺), 293 (100, [M+Na]⁺).

6.4. *4-Hydroxy-3,3-dimethyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]butanamide (4d)*. According to *GP 4*, from **9d** (1044 mg, 3 mmol), 6 h. Yield: 845 mg (92%) of **4d**. White powder. M.p. 138.4–139.9°. ¹H-NMR: 0.92, 1.44 (2s, 2 Me₂C); 2.13 (s, CH₂); 3.22 (s, MeN); 3.91 (s, CH₂O); 6.32 (s, NH); 7.16–7.41 (m, 5 arom. H). ¹³C-NMR: 25.2, 26.0 (2q, 2 Me₂C); 35.6 (s, Me₂C); 41.3 (q, MeN); 47.2 (t, CH₂); 58.2 (s, Me₂C); 71.4 (t, CH₂O); 127.9, 128.2, 129.2 (3d, 5 arom. CH); 144.3 (s, arom. C); 171.6, 173.1 (2s, 2 C=O). ESI-MS: 345 (20, [M+K]⁺), 329 (100, [M+Na]⁺). Anal. calc. for C₁₇H₂₆N₂O₃ (306.41): C 66.64, H 8.55, N 9.14; found: C 66.36, H 8.49, N 8.97.

6.5. *4-Hydroxy-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]-3-phenylbutanamide (4e)*. According to *GP 4*, from **9e** (1188 mg, 3 mmol), 8 h. Yield: 934 mg (88%) of **4e**. White powder. M.p. 129.0–129.9°. ¹H-NMR: 1.28, 1.35 (2s, Me₂C); 2.11–2.21, 2.34–2.46 (2m, CH₂); 3.17 (br. s, MeN, CH); 3.64–3.79 (m, CH₂O); 5.96 (s, NH); 7.10–7.41 (m, 10 arom. H). ¹³C-NMR: 26.1, 26.2 (2q, Me₂C); 40.9 (q, MeN); 41.2 (t, CH₂); 44.5 (d, CH); 58.2 (s, Me₂C); 67.1 (t, CH₂O); 126.8, 127.5, 127.9, 128.6, 129.3 (5d, 10 arom. CH); 141.7, 144.4 (2s, 2 arom. C); 171.1, 173.0 (2s, 2 C=O). CI-MS (i-butane): 355 (100, [M+H]⁺).

6.6. *4-Hydroxy-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]-3-methyl-3-phenylbutanamide (4f)*. According to *GP 4* from **9f** (820 mg, 2 mmol), 12 h. Yield: 587 mg (90%) of **4f**. White powder. M.p. 121.9–123.3°. ¹H-NMR: 1.22, 1.23, 1.24 (3s, Me, Me₂C); 2.11–2.19, 2.31–2.38 (2m, CH₂); 3.15 (s, MeN); 3.51–3.59, 3.83–3.89 (2m, CH₂O); 5.84 (s, NH); 7.11–7.43 (m, 10 arom. H). ¹³C-NMR: 23.6, 26.1, 26.2 (3q, Me, Me₂C); 41.4 (q, Me₂N); 42.7 (s, Me₂C); 47.4 (t, CH₂); 58.1 (s, Me₂C); 70.6 (t, CH₂O); 126.1, 126.5, 127.8, 128.5, 129.3 (5d, 10 arom. CH); 144.5, 145.5 (2s, 2 arom. C); 170.9, 173.0 (2s, 2 C=O). CI-MS (i-butane): 327 (80, [M+H]⁺), 221 (100, [M-Ph(Me)N]⁺).

6.7. *4-Hydroxy-2,3,3-trimethyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]butanamide (4g)*. According to *GP 4* from **9g** (724 mg, 2 mmol), 4 h. Yield: 576 mg (90%) of **4g**. White powder. M.p. 124.2–126.3°. ¹H-NMR: 1.00 (s, Me₂C); 1.09 (d, J=4.6, MeCH); 1.40, 1.42 (2s, Me₂C); 2.08 (q, J=4.6, CH); 3.18 (s, MeN); 3.51–3.59 (m, CH₂O); 6.96 (s, NH); 7.12–7.42 (m, 5 arom. H). ¹³C-NMR: 12.7 (q, Me); 23.4, 24.2, 25.0 (3q, 3 Me₂C); 37.6 (s, Me₂C); 41.7 (q, MeN); 49.9 (d, CH); 58.9 (s, Me₂C); 69.3 (t, CH₂O); 128.4, 128.6, 129.5 (3d, 5 arom. CH); 143.9 (s, arom. C); 173.8, 175.6 (2s, 2 C=O). ESI-MS: 343 (100, [M+Na]⁺).

6.8. *4-Hydroxy-2,2,3,3-tetramethyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]butanamide (4h)*. According to *GP 4*, from **9h** (752 mg, 2 mmol), 4 h. Yield: 608 mg (91%) of **4h**. White powder. M.p. 129.9–131.3°. ¹H-NMR: 0.88, 1.15, 1.46 (3s, 3 Me₂C); 3.29 (s, MeN); 3.44 (s, CH₂O); 6.48 (s, NH); 7.23–7.42 (m, 5 arom. H). ¹³C-NMR: 21.5, 21.9, 24.6 (3q, 3 Me₂C); 39.6 (s, Me₂C); 41.6 (q, MeN); 47.6, 58.9 (2s, 2 Me₂C); 70.7 (t, CH₂O); 128.2, 128.4, 129.4 (3d, 5 arom. CH); 143.9 (s, arom. C); 173.9, 177.8 (2s, 2 C=O). ESI-MS: 357 (100, [M+Na]⁺).

7. *Attempted Direct Amide Cyclizations of Amides 4. General Procedure 5 (GP 5)*. A suspension of **4** (0.5 mmol) in dry toluene (50 ml) was heated to 100°, and HCl gas was bubbled through the suspension for 4–6 min. Then, the mixture was allowed to cool to r.t. while bubbling N₂ through it (ca. 20 min). The solvent was evaporated, the white residue was washed with CH₂Cl₂ (3 × 15 ml) and dried in h.v. to yield imino lactone hydrochlorides **12** as white powders.

General Procedure 6 (GP 6). A suspension of **4** was treated as described in *GP 5*. The solvent was evaporated, and the oily residue was purified by CC to yield the ω-chloro acids **11**.

7.1. *2-[(4,4-Dimethyltetrahydrofuran-2-ylidene)amino]-2-methylpropanoic Acid Hydrochloride (12a)*. According to *GP 5* from **4d** (153 mg, 0.5 mmol). Yield: 101 mg (84%). M.p. 158.3–159.1°. IR: 3431w, 2938s, 2804s, 1735vs, 1678vs, 1529s, 1414m, 1319m, 1194s, 1166s, 1003m, 946m, 754m. ¹H-NMR ((D₆)DMSO): 1.15, 1.56 (2s, 2 Me₂C); 3.13 (s, CH₂); 4.56 (s, CH₂O); 13.2 (br. s, COOH). ¹³C-NMR ((D₆)DMSO): 23.5, 24.1 (2q, 2 Me₂C); 38.0, 60.1 (s, Me₂C); 88.0 (t, CH₂O); 162.0 (s, C=N); 180.2 (s, C=O). CI-MS: 200 (100, [M-Cl]⁺).

7.2. *2-[(3,4,4-Trimethyltetrahydrofuran-2-ylidene)amino]-2-methylpropanoic Acid Hydrochloride (12g)*. According to *GP 5* from **4g** (160 mg, 0.5 mmol). Yield: 78 mg (73%). M.p. 152.4–153.6°. IR: 3386w, 2986s, 2776s, 1739vs, 1671vs, 1514m, 1478s, 1240m, 1178s, 1165s, 997s, 773m. ¹H-NMR ((D₆)DMSO): 1.08 (d, J=4.9, Me); 1.21, 1.48 (2s, 2 Me₂C); 2.18 (q, J=4.9, CH); 4.42 (s, CH₂O); 10.0 (br. s, COOH). ¹³C-NMR ((D₆)DMSO): 12.0 (q, Me); 21.1, 22.2 (2q, 2 Me₂C); 36.5 (s, Me₂C); 46.1 (d, CH); 58.2 (s, Me₂C); 87.2 (t, CH₂O); 163.1 (C=N); 178.0 (s, C=O). CI-MS (i-butane): 214 (100, [M-

Cl]⁺), 108 (25). Anal. calc. for C₁₁H₂₀ClNO₃ (249.74): C 52.90, H 8.07, N 5.61; found C 50.55, H 7.87, N 5.25.

7.3. 2-[(3,3,4,4-Tetramethyltetrahydrofuran-2-ylidene)amino]-2-methylpropanoic Acid Hydrochloride (**12h**). According to GP 5 from **4h** (167 mg, 0.5 mmol). Yield: 79 mg (78%). M.p. 157.0–157.6°. IR: 3390w, 3024s, 2938s, 2778s, 1738vs, 1669vs, 1476s, 1400s, 1313m, 1178s, 998m, 880m, 772m. ¹H-NMR ((D₆)DMSO): 0.97, 1.26, 1.56 (3s, 3 Me₂C); 4.58 (s, CH₂O); 12.9 (br. s, COOH). ¹³C-NMR ((D₆)DMSO): 21.6, 22.5, 26.1 (3q, 3 Me₂C); 42.8, 46.6, 60.7 (3s, 3 Me₂C); 86.8 (t, CH₂O); 168.1 (C=N); 183.2 (s, C=O). ESI-MS: 228 (100, [M+Na]⁺).

7.4. 2-[(4-Chloro-3,3-dimethylbutanoyl)amino]-2-methylpropanoic Acid (**11a**). According to GP 6 from **4a** (122 mg, 0.5 mmol) or from **4d** (153 mg, 0.5 mmol), CC (CH₂Cl₂/acetone 10:1). Yield: 57 mg (48%) of **11a** from **4a** and 62 mg (52%) from **4d**. White powder. M.p. 116.1–117.3°. IR: 3356vs, 2981s, 2605m, 1717vs, 1620vs, 1548vs, 1473s, 1393s, 1294m, 1221s, 1158s, 1023m, 895m, 791s. ¹H-NMR ((D₆)DMSO): 1.00, 1.31 (2s, 2 Me₂C); 2.09 (s, CH₂); 3.59 (s, CH₂Cl); 7.12 (br. s, NH); 12.9 (br. s, COOH). ¹³C-NMR ((D₆)DMSO): 24.9, 25.8 (2q, 2 Me₂C); 36.6 (s, Me₂C); 44.9 (t, CH₂); 55.3 (t, CH₂Cl); 60.3 (s, Me₂C); 172.1, 181.1 (2s, 2 C=O). ESI-MS: 260 (32, [M(³⁷Cl)+Na]⁺), 258 (100, [M(³⁵Cl)+Na]⁺).

7.5. 2-[(4-Chlorobutanoyl)amino]-2-methylpropanoic Acid (**11b**). According to GP 6 from **4b** (140 mg, 0.5 mmol), CC (CH₂Cl₂/acetone 10:1). Yield: 62 mg (66%) of **11b**. White solid. M.p. 111.9–113.1°. IR: 3318vs, 2988s, 1721vs, 1623s, 156s, 1467s, 1399m, 1230m, 1166s, 1050w, 945w, 787m. ¹H-NMR: 1.57 (s, Me₂C); 2.07–2.13 (m, CH₂); 2.37 (t, J=6.1, CH₂); 3.61 (t, J=6.1, CH₂Cl); 7.29 (br. s, NH). ¹³C-NMR: 24.7 (q, Me₂C); 33.9 (t, CH₂); 44.4 (t, CH); 55.6 (t, CH₂Cl); 59.6 (s, Me₂C); 171.8, 176.6 (2s, 2 C=O). CI-MS: 210 (10, [M(³⁷Cl)+NH₄]⁺), 208 (25, [M(³⁵Cl)+NH₄]⁺), 172 (100, [M–Cl]⁺).

7.6. 1-[(4-Chloro-3,3-dimethylbutylamino)cyclopentanecarboxylic Acid (**11c**). According to GP 6 from **4c** (135 mg, 0.5 mmol), CC (CH₂Cl₂/acetone 10:1). Yield: 69 mg (53%) of **11c**. White powder. M.p. 113.2–114.7°. ¹H-NMR: 1.02, 1.09 (2s, Me₂C); 1.71–1.76, 1.94–2.02 (2m, 4 CH₂); 3.22 (m, CH₂); 3.63 (m, CH₂Cl); 7.47 (br. s, NH). ¹³C-NMR: 24.8 (t, CH₂); 25.3 (q, Me₂C); 36.3 (q, Me₂C); 37.5, 44.7 (2t, 2 CH₂); 55.0 (t, CH₂Cl); 66.1 (s, Me₂C); 171.7, 176.4 (2s, 2 C=O). CI-MS: 264 (28, [M(³⁷Cl)+H]⁺), 262 (72, [M(³⁵Cl)+H]⁺), 226 (100, [M–Cl]⁺).

7.7. [2-(4-Chloro-3-phenylbutanoyl)amino]-2-methylpropanoic Acid (**11e**). According to GP 6 from **4e** (184 mg, 0.5 mmol), CC (CH₂Cl₂/acetone 20:1). Yield: 51 mg (43%) of **11e**. White powder. M.p. 119.6–121.8° (dec.). IR: 3320vs, 2928s, 1722vs, 1626vs, 1606s, 1557s, 1496s, 1442s, 1315s, 1260m, 1170s, 1079m, 752s, 696s. ¹H-NMR: 1.36, 1.39 (2s, Me₂C); 2.53–2.60, 2.70–2.78 (m, CH₂); 3.48–3.52 (m, PhCH); 3.85–3.91 (m, CH₂Cl); 7.24–7.32 (m, 5 arom. H); 7.45 (br. s, NH); 12.3 (br. s, COOH). ¹³C-NMR: 24.8, 25.3 (2q, 2 Me₂C); 40.2 (t, CH₂); 45.3 (d, CH); 55.3 (t, CH₂Cl); 59.6 (s, Me₂C); 127.6, 128.7, 129.0 (3d, 5 arom. CH); 142.3 (s, arom. C); 171.1, 179.1 (2s, 2 C=O). CI-MS: 286 (15, [M(³⁷Cl)+H]⁺), 284 (39, [M(³⁵Cl)+H]⁺), 248 (100, [M–Cl]⁺).

8. Synthesis of Dipeptide Esters. General Procedure 7 (GP 7). A suspension of **4** (0.5 mmol) in a mixture of dry toluene (50 ml) and alcohol (5 ml) was heated to reflux, and HCl gas was bubbled through the suspension for 6 min. Then, the mixture was allowed to cool to r.t. while bubbling N₂ through it (ca. 20 min). The solvent was evaporated, and the oily residue was purified by CC to yield the products as colorless oils.

General Procedure 8 (GP 8). To solid **12a** (117 mg, 0.5 mmol), a 0.5M soln. of CH₂N₂ in Et₂O (1 ml, 0.5 mmol) was added dropwise at 0°. After the disappearance of the yellow color, another 1 ml of CH₂N₂ was added, the mixture allowed to warm to r.t., stirred for 15 min, filtered, and the solvent was evaporated *i.v.* The white solid **14a** was used without further purification.

8.1. Methyl 2-[(4,4-Dimethyltetrahydrofuran-2-ylidene)amino]-2-methylpropanoate (**14a**). According to GP 7 from **4d** (153 mg, 0.5 mmol), 5 ml of MeOH, CC (CH₂Cl₂/acetone 20:1). Yield: 50 mg (43%) of **14a**. White solid. M.p. 88.6–89.8°. IR: 2961s, 2876m, 1736vs, 1706vs, 1567s, 1360m, 1271s, 1143s, 101m, 919m, 731s. ¹H-NMR ((D₆)DMSO): 1.05, 1.28 (2s, 2 Me₂C); 2.24 (s, CH₂); 3.56 (s, MeO); 3.83 (s, CH₂O). ¹³C-NMR ((D₆)DMSO): 23.6, 25.8 (2q, Me₂C); 36.6 (s, Me₂C); 42.4 (t, CH₂); 50.1 (q, MeO); 58.9 (s, Me₂C); 80.3 (t, CH₂O); 162.2 (s, C=N); 175.3 (s, C=O). CI-MS: 231 (100, [M+NH₄]⁺).

An experiment according to GP 8 yielded 113 mg (98%) of **14a** as a white solid.

8.2. Ethyl 2-[(4,4-Dimethyltetrahydrofuran-2-yliden)amino]-2-methylpropanoate (**14b**). According to GP 7 from **4d** (153 mg, 0.5 mmol), 5 ml of EtOH, CC (CH₂Cl₂/acetone 20 : 1). Yield: 49 mg (43%) of **14b** and 51 mg of ethyl 2-[(4-chloro-3,3-dimethylbutanoyl)amino]-2-methylpropanoate **18a** (38%) as colorless oils.

Data of **14b**: IR: 2965vs, 2893s, 1714vs, 1697vs, 1468s, 1380s, 1273s, 1166s, 1035m, 911m, 731s. ¹H-NMR: 1.11 (s, Me₂C); 1.21 (t, *J* = 7.1, MeCH₂); 1.43 (s, Me₂C); 2.33 (s, CH₂); 3.82 (s, CH₂O); 4.18 (q, *J* = 7.1, MeCH₂). ¹³C-NMR: 14.3 (q, Me); 25.1, 26.6 (2q, Me₂C); 37.1 (s, Me₂C); 44.5 (t, CH₂); 60.4 (t, MeCH₂); 60.5 (s, Me₂C); 81.9 (t, CH₂O); 163.7 (s, C=N); 176.5 (s, C=O). ESI-MS: 477 (15, [2M + Na]⁺), 250 (100, [M + Na]⁺), 228 (12, [M + H]⁺).

Data of **18a**: ¹H-NMR: 1.05 (s, Me₂C); 1.16 (t, *J* = 7.2, MeCH₂); 1.27 (s, Me₂C); 2.24 (s, CH₂); 3.81 (s, CH₂Cl); 4.02 (q, *J* = 7.2, MeCH₂); 6.02 (br. s, NH). ¹³C-NMR: 14.2 (q, Me); 24.7, 25.8 (2q, 2 Me₂C); 35.8 (s, Me₂C); 38.4 (t, CH₂); 54.9 (t, CH₂O); 61.3 (t, MeCH₂); 65.6 (s, Me₂C); 170.4, 172.6 (2s, 2 C=O). ESI-MS: 288 (31, [M(³⁷Cl) + Na]⁺), 286 (100, [M(³⁵Cl) + Na]⁺).

8.3. Methyl 2-Methyl-2-[(4-methyl-4-phenyltetrahydrofuran-2-yliden)amino]propanoate (**14c**). According to GP 7 from **4f** (168 mg, 0.5 mmol), 5 ml of MeOH, CC (CH₂Cl₂/acetone 25 : 1). Yield: 77 mg (56%) of **14c**. Colorless oil. ¹H-NMR: 1.45, 1.46, 1.49 (3s, Me, Me₂C); 2.73–2.79, 2.97–3.02 (2m, CH₂); 3.68 (s, MeO); 4.31 (s, CH₂O); 7.16–7.36 (m, 5 arom. H). ¹³C-NMR: 26.1, 26.7, 27.1 (3q, Me, Me₂C); 43.0 (t, CH₂); 44.6 (s, Me₂C); 51.3 (q, MeO); 60.1 (s, Me₂C); 80.9 (t, CH₂O); 127.3, 126.8, 128.7 (3d, 5 arom. CH); 144.5 (s, arom. C); 163.0 (s, C=N); 176.7 (s, C=O). CI-MS: 276 (100, [M + H]⁺), 108 (22).

8.4. Ethyl 1-[(4,4-Dimethyltetrahydrofuran-2-yliden)amino]cyclopentanecarboxylate (**14d**). According to GP 7 from **4c** (135 mg, 0.5 mmol), 5 ml of EtOH, CC (CH₂Cl₂/acetone 20 : 1). Yield: 71 mg (28%) of **14d** and 45 mg (31%) of ethyl 1-[(4-chloro-3,3-dimethylbutanoyl)amino]cyclopentanecarboxylate (**18b**) as colorless oils.

Data of **14d**: ¹H-NMR: 1.12 (s, Me₂C); 1.21 (t, *J* = 7.0, MeCH₂); 1.81–1.94 (m, 4 CH₂); 2.41 (s, CH₂); 3.89 (s, CH₂O); 4.14 (q, *J* = 7.0, MeCH₂). ¹³C-NMR: 14.3 (q, Me); 25.1 (q, Me₂C); 37.3, 37.6, 38.4, 39.1 (4t, 4 CH₂); 44.2 (s, Me₂C); 44.9, 47.1 (2t, 2 CH₂); 60.6 (s, Me₂C); 82.2 (t, CH₂O); 162.4 (s, C=N); 174.9 (s, C=O). CI-MS: 254 (100, [M + H]⁺), 200 (18, [M – Et]⁺), 158 (36).

Data of **18b**: ¹H-NMR: 1.09 (s, Me₂C); 1.20 (t, *J* = 6.9, MeCH₂); 1.74–1.82, 1.84–1.94 (2m, 4 CH₂, Me₂C); 2.16 (s, CH₂); 3.56 (s, CH₂O); 4.16 (q, *J* = 6.9, MeCH₂); 6.02 (br. s, NH). ¹³C-NMR: 14.2 (q, Me); 24.7, 25.8 (2q, 2 Me₂C); 35.8 (s, Me₂C); 37.4, 37.7, 38.4, 39.1 (4t, 4 CH₂); 44.2 (t, CH₂); 54.9 (t, CH₂O); 61.3 (t, MeCH₂); 65.6 (s, Me₂C); 170.4, 172.6 (2s, 2 C=O). CI-MS: 292 (34, [M(³⁷Cl) + H]⁺), 290 (100, [M(³⁵Cl) + H]⁺), 254 (28).

8.5. Methyl 2-[(4-Hydroxy-3,3-dimethylbutanoyl)amino]-2-methylpropanoate (**17a**). According to GP 7 from **9d** (174 mg, 0.5 mmol), 5 ml of MeOH, CC (CH₂Cl₂/acetone 20 : 1). Yield: 56 mg (48%) of **17a**. Colorless oil. ¹H-NMR: 0.98, 1.43 (2s, 2 Me₂C); 2.11 (s, CH₂); 3.29 (s, CH₂O); 3.78 (s, MeO); 6.25 (s, NH). ¹³C-NMR: 24.6, 25.2 (2q, 2 Me₂C); 35.7 (s, Me₂C); 46.9 (t, CH₂); 52.5 (q, MeO); 56.4 (s, Me₂C); 71.2 (t, CH₂O); 171.9, 174.9 (2s, 2 C=O). ESI-MS: 254 (100, [M + Na]⁺).

8.6. Methyl 2-[(4-Hydroxy-3-methyl-3-phenylbutanoyl)amino]-2-methylpropanoate (**17b**). According to GP 7 from **9f** (205 mg, 0.5 mmol) with 5 ml MeOH, CC (CH₂Cl₂/acetone 20 : 1). Yield 57 mg (38%) of **17b**. Colorless oil. ¹H-NMR: 1.38 (s, Me₂C); 1.39 (s, Me); 2.47–2.51, 2.66–3.70 (2m, CH₂); 3.65–3.69, 3.89–3.92 (2m, CH₂O); 3.68 (s, MeO); 6.25 (s, NH); 7.19–7.42 (m, 5 arom. H). ¹³C-NMR: 23.3, 24.5, 24.6 (3q, 3 Me); 43.0 (t, CH₂); 46.5 (s, C(Me)(Ph)); 52.3 (q, MeO); 56.1 (s, Me₂C); 70.3 (t, CH₂O); 126.0, 126.4, 128.4 (3d, 5 arom. CH); 145.4 (s, arom. C); 171.2, 174.7 (2s, 2 C=O). CI-MS (i-butane): 294 (100, [M + H]⁺).

9. Protection of the Hydroxy Amide **4d**. 9.1. 4-(Benzyloxy)-3,3-dimethyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]butanamide (**9i**). To a soln. of **4d** (1 mmol, 306 mg) in dry THF (20 ml), NaH (44 mg, 1.1 mmol, 60% suspension in mineral oil) was added at 0°. After 1 h at r.t., BnBr (171 mg, 1 mmol) was added, and the mixture was heated to reflux for 3 h. Washing with brine, extraction with Et₂O, drying (MgSO₄), and evaporation *i.v.* yielded a colorless oil, which was purified by CC (CH₂Cl₂/acetone 40 : 1). Yield: 329 mg (73%) of **9i**. Colorless oil. ¹H-NMR: 0.99, 1.36 (2s, 2 Me₂C); 2.18 (s, CH₂); 3.12 (s, CH₂O); 3.22 (s, MeN); 4.43 (s, PhCH₂); 5.91 (br. s, NH); 7.12–7.43 (m, 10 arom. H). ¹³C-NMR: 25.4, 26.7 (2q, 2 Me₂C); 34.8 (s, Me₂C); 41.3 (q, MeN); 46.3 (t, CH₂); 57.7 (s, Me₂C); 73.1 (t, CH₂O); 78.5 (t, PhCH₂);

127.4, 127.6, 127.7, 127.8, 128.3, 129.2 (6d, 10 arom. CH); 138.4, 145.0 (2s, 2 arom. C); 170.6, 173.2 (2s, 2 C=O). CI-MS (i-butane): 397 (100, [M+H]⁺).

9.2. (2,2-Dimethyl-3-[[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]carbamoyl]propyl)[(9H-fluoren-9-yl)methyl] Carbonate (**9k**). To a soln. of **4d** (306 mg, 1 mmol) in CH₂Cl₂ (20 ml), pyridine (3 ml), and FmocCl (284 mg, 1.1 mmol) were added. The mixture was stirred at r.t. for 3 h, diluted with CH₂Cl₂ (30 ml), and washed with a 10% CuSO₄ soln. and brine, and the org. fractions were dried (MgSO₄), evaporated *i.v.*, and purified by CC (CH₂Cl₂/acetone 30:1). Yield: 450 mg (88%) of **9k**. White powder. M.p. 108.9–110.6°. IR: 3320vs, 2966s, 1745vs, 1678vs, 1650vs, 1599s, 1532s, 1441s, 1386s, 1316m, 1154m, 964m, 909s, 758m. ¹H-NMR: 0.99, 1.39 (2s, 2 Me₂C); 1.86 (s, CH₂); 3.19 (s, MeN); 3.92 (s, CH₂O); 1.15 (t, J=5.8, CH₂CH); 4.36 (d, J=5.8, CH₂CH); 6.12 (s, NH); 7.09–7.32, 7.50–7.80 (2m, 13 arom. H). ¹³C-NMR: 24.5, 25.5 (2q, 2 Me₂C); 34.2 (s, Me₂C); 41.3 (q, MeN); 45.3 (t, CH₂); 58.0 (s, Me₂C); 69.7 (t, CH₂O); 75.1 (t, CH₂CH); 119.9 (d, CH); 125.0, 127.0, 127.7, 127.8, 127.9, 129.1 (6d, 13 arom. CH); 141.2, 143.3, 144.5 (3s, 5 arom. C); 155.2 (s, NC(O)O); 169.4, 173.1 (2s, 2 C=O). ESI-MS: 551 (100, [M+Na]⁺).

9.3. 4-[(Benzyloxy)methoxy]-3,3-dimethyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]butanamide (**9l**). To a soln. of **4d** (306 mg, 1 mmol) in CH₂Cl₂ (20 ml) was added (i-Pr)₂NEt (142 mg, 1.1 mmol) and BnOCH₂Cl (BomCl, 156 mg, 1 mmol). The mixture was stirred at r.t. for 12 h, diluted with CH₂Cl₂ (30 ml), washed with NH₄Cl soln. and brine, dried (MgSO₄), evaporated *i.v.*, and purified by CC (CH₂Cl₂/acetone 40:1). Yield: 375 mg (88%) **9l**. Colorless oil. ¹H-NMR: 0.99, 1.42 (2s, 2 Me₂C); 1.84 (s, CH₂); 3.22 (s, MeN); 3.32 (s, CH₂); 4.51, 4.68 (2s, CH₂); 5.94 (br. s, NH); 7.08–7.42 (m, 10 arom. H). ¹³C-NMR: 25.0, 26.3 (2q, 2 Me₂C); 34.5 (s, Me₂C); 41.2 (q, MeN); 45.9 (t, CH₂); 57.8 (s, Me₂C); 69.3, 76.2, 94.8 (3t, 3 CH₂); 127.6, 127.7, 127.8, 128.3, 129.0, 129.2 (6d, 10 arom. CH); 137.6, 144.6 (2s, 2 arom. C); 170.2, 173.1 (2s, 2 C=O). CI-MS: 427 (16, [M+H]⁺), 320 (100, [M–Ph(Me)N]⁺), 108 (16).

10. Synthesis of 1,3-Oxazol-5(4H)-ones. 10.1. 3-(4,5-Dihydro-4,4-dimethyl-5-oxo-1,3-oxazol-2-yl)-2,2-dimethylpropyl Acetate (**16a**). According to GP 6 from **9d** (153 mg, 0.5 mmol), 4 min HCl gas, CC (CH₂Cl₂/acetone 40:1). Yield: 123 mg (51%) of **16a**. White solid. M.p. 99.5–101.7°. Recovered starting material: 33 mg (21%). IR: 3385m, 2976s, 2937s, 1820vs, 1739vs, 1672vs, 1525m, 1474s, 1379s, 1240vs, 1072s, 1041s, 966s, 897m. ¹H-NMR: 1.05, 1.36 (2s, 2 Me₂C); 2.04 (s, MeCO); 2.42 (s, CH₂); 3.92 (s, CH₂O); 7.21–7.34 (m, 5 arom. H). ¹³C-NMR: 20.8 (q, MeCO); 24.4, 24.6 (2q, 2 Me₂C); 34.3 (s, Me₂C); 38.0 (t, CH₂); 65.3 (s, Me₂C); 71.6 (t, CH₂O); 161.9 (s, C=N); 170.9, 181.4 (2s, 2 C=O). ESI-MS: 541 (100, [2(M+H₂O)+Na]⁺), 371 (25), 282 (70, [M+H₂O+Na]⁺).

10.2. 2-[3-(Benzyloxy)-2,2-dimethylpropyl]-4,4-dimethyl-1,3-oxazol-5(4H)-one (**16b**). According to GP 6, from **9i** (198 mg, 0.5 mmol), 4 min HCl gas, CC (CH₂Cl₂/acetone 50:1). Yield: 49 mg (34%) of **16b**. White powder. M.p. 104.2–106.0°. Recovered starting material: 124 mg (31%). IR: 2951s, 1894s, 2565w, 1822vs, 1723vs, 1612vs, 1534vs, 1458s, 1265s, 1101s, 1058m, 1040m, 911m. ¹H-NMR: 1.01, 1.38 (2s, 2 Me₂C); 2.48 (s, CH₂); 3.14 (s, CH₂O); 4.56 (s, PhCH₂); 7.21–7.34 (m, 5 arom. H). ¹³C-NMR: 24.5, 24.9 (2q, 2 Me₂C); 35.4 (s, Me₂C); 38.3 (t, CH₂); 65.2 (s, Me₂C); 73.3 (t, CH₂O); 78.6 (t, PhCH₂); 127.5, 128.1, 128.3 (3d, 5 arom. CH); 138.5 (s, arom. C); 162.5 (s, C=N); 182.1 (s, C=O). CI–MS (i-butane): 290 (18, [M+H]⁺), 200 (100, [M–Bn]⁺).

10.3. [3-(4,5-Dihydro-4,4-dimethyl-5-oxo-1,3-oxazol-2-yl)-2,2-dimethylpropyl] [(9H-fluoren-9-yl)methyl] Carbonate (**16c**). According to GP 6 from **9k** (169 mg, 0.33 mmol), 4 min HCl gas, CC (CH₂Cl₂/acetone 80:1). Yield: 64 mg (46%) of **16c**. White powder. M.p. 93.8–95.2°. Recovered starting material: 66 mg (39%). IR: 3320s, 2966s, 1821vs, 1745vs, 1678vs, 1650vs, 1599s, 1532s, 1441s, 1386m, 1316m, 1156m, 909s. ¹H-NMR: 1.01, 1.34 (2s, 2 Me₂C); 2.32 (s, CH₂); 3.87 (s, CH₂O); 4.12 (t, J=6.0, CH₂-CH); 4.38 (d, J=6.0, CH₂CH); 7.18–7.37, 7.53–7.81 (2m, 8 arom. H). ¹³C-NMR: 24.3, 24.8 (2q, 2 Me₂C); 34.5 (s, Me₂C); 37.9, 46.7 (2t, 2 CH₂); 65.1 (s, Me₂C); 65.9 (t, CH₂O); 69.8 (t, CH₂CH); 119.9 (d, CH); 125.0, 126.8, 127.1, 127.7 (4d, 8 arom. CH); 141.2, 143.3 (2s, 4 arom. C); 155.0 (s, NC(O)O); 161.6 (s, C=N); 181.1 (s, C=O). CI-MS: 422 (16, [M+H]⁺), 200 (100, [M–Fmoc]⁺), 179 (44).

10.4. 2-[3-[(Benzyloxy)methoxy]-2,2-dimethylpropyl]-4,5-dihydro-4,4-dimethyl-1,3-oxazol-5-one (**16d**). According to GP 6 from **9l** (143 mg, 0.33 mmol), 5 min HCl gas, CC (CH₂Cl₂/acetone 60:1). Yield: 42 mg (36%). White solid. M.p. 104.1–105.8°. Recovered starting material: 52 mg (35%). ¹H-NMR: 0.97, 1.36 (2s, 2 Me₂C); 2.42, 3.36, 4.59, 4.73 (4s, 4 CH₂); 7.14–7.43 (m, 5 arom. H). ¹³C-NMR:

24.4, 24.7 (2*q*, 2 Me₂C); 34.9 (s, Me₂C); 45.5 (t, CH₂); 56.8 (s, Me₂C); 69.3, 76.1, 94.8 (3*t*, 3 CH₂); 127.7, 128.3, 129.5 (3*d*, 5 arom. CH); 137.6 (s, arom. C); 162.4 (s, C=N); 172.6 (s, C=O). CI-MS (i-butane): 320 (100, [M+H]⁺).

10.5. 2,2-Dimethyl-3-(4-oxo-3-oxa-1-azaspiro[4.4]non-1-en-2-yl)propyl Acetate (**16e**). According to GP 6 from **9c** (156 mg, 0.5 mmol), 5 min HCl gas, CC (CH₂Cl₂/acetone 30:1). Yield: 45 mg (34%) of **16e**. White solid. M.p. 101.1–103.1°. Recovered starting material: 33 mg (21%). IR: 3385*w*, 2976*s*, 2937*s*, 1820*vs*, 1739*vs*, 1672*vs*, 1474*m*, 1379*s*, 1240*s*, 1072*s*, 1041*s*, 966*m*. ¹H-NMR: 1.01 (s, Me₂C); 1.73–2.04 (m, MeCO, 4 CH₂); 2.42 (s, CH₂); 3.90 (s, CH₂O). ¹³C-NMR: 20.8 (q, MeCO); 24.5 (q, Me₂C); 25.8 (t, 2 CH₂); 34.3 (s, Me₂C); 38.0, 38.2 (2*t*, 3 CH₂); 71.6 (t, CH₂O); 73.9 (s, spiro-C); 161.8 (s, C=N); 170.9, 182.1 (2*s*, 2 C=O). ESI-MS: 593 (100, [2M+H₂O+Na]⁺), 324 (58, [M+H₂O+K]⁺), 308 (64, [M+H₂O+Na]⁺), 286 (90, [M+H₂O+H]⁺).

11. 2-Methyl-N-[1-methyl-1-(N-methyl-N-phenylcarbamoyl)ethyl]-2-[(2,3,4,5-tetrahydro-4,4-dimethylfuran-2-ylidene)amino]propanamide (**15**). To solid **12a** (100 mg, 0.42 mmol), a soln. of **10b** (110 mg, 3 mmol, 1.5 equiv.) in CH₂Cl₂ (5 ml) was added under vigorous stirring. The solvent was immediately removed *i.v.*, and the suspension was left at r.t. overnight. CC (CH₂Cl₂/acetone 20:1) yielded 102 mg (64%) of **15**. White powder. M.p. 142.1–143.3°. IR: 3364*vs*, 2976*s*, 2937*s*, 1708*vs*, 1666*vs*, 1645*vs*, 1493*s*, 1380*s*, 1223*m*, 1111*m*, 1014*s*, 781*m*, 710*s*. ¹H-NMR: 1.08, 1.42, 1.50 (3*s*, 3 Me₂C); 2.26 (s, CH₂); 3.21 (s, MeN); 3.89 (s, CH₂O); 7.06–7.29 (m, 5 arom. H); 8.07 (br. s, NH). ¹³C-NMR: 23.8, 24.9, 26.1 (3*q*, 3 Me₂C); 36.5 (s, Me₂C); 41.0 (q, MeN); 45.2 (t, CH₂); 57.3, 60.6 (2*q*, 2 Me₂C); 81.5 (t, CH₂O); 127.3, 128.1, 129.4 (3*d*, 5 arom. CH); 144.8 (s, arom. C); 161.5 (s, C=N); 173.6, 176.3 (2*s*, 2 C=O). ESI-MS: 369 (100, [M+Na]⁺).

12. Methyl 2-[(4-Hydroxy-3,3-dimethyl-2-oxobutyl)amino]-2-methylpropanoate (**19**). To a soln. of 1-bromo-4-hydroxy-3,3-dimethylbutan-2-one (390 mg, 2 mmol) in Et₃N (5 ml), methyl 1-amino-2-methylpropanoate hydrochloride (456 mg, 3 mmol) was added. The mixture was heated under reflux overnight, cooled, diluted with CH₂Cl₂ (150 ml), and washed with a 10% CuSO₄ soln. and brine. Drying (MgSO₄), evaporation of the solvent, and CC (CH₂Cl₂/acetone 30:1) yielded 195 mg (42%) of **19**. Yellow oil. IR: 2973*m*, 2884*m*, 1725*vs*, 1660*vs*, 1528*m*, 1471*s*, 1364*s*, 1192*s*, 1046*m*. ¹H-NMR: 1.12, 1.46 (2*s*, 2 Me₂C); 3.50, 3.54 (2*s*, 2 CH₂); 3.71 (s, MeO); 6.74 (br. s, NH). ¹³C-NMR: 21.1, 25.1 (2*q*, 2 Me₂C); 48.4 (s, Me₂C); 49.9 (t, CH₂); 51.9 (q, MeO); 58.2 (s, Me₂C); 69.4 (t, CH₂O); 176.4, 212.6 (2*s*, 2 C=O). CI-MS (i-butane): 232 (70, [M+H]⁺), 146 (100), 128 (40).

13. X-Ray Crystal-Structure Determination of **11a**, **11b**, **12g**, **12h**, and **15** (Table 3 and Figs. 2–4)⁶⁾. All measurements were performed on a Nonius KappaCCD area-detector diffractometer [36] using graphite-monochromated MoK_α radiation (λ 0.71073 Å) and an Oxford Cryosystems Cryostream 700 cooler. The data collection and refinement parameters are given in Table 3, and views of the molecules are shown in Figs. 2–4. Data reduction was performed with HKL Denzo and Scalepack [37]. The intensities were corrected for Lorentz and polarization effects and, in the cases of **11a**, **12g**, and **12h**, an absorption correction based on the multi-scan method [38] was applied. Each structure was solved by direct methods using SIR92 [39], which revealed the positions of all non-H atoms. The non-H atoms were refined anisotropically.

In the case of **11b**, there are two independent molecules in the asymmetric unit. The atomic coordinates of the two molecules were tested carefully for a relationship from a higher symmetry space group using the program PLATON [40] but none could be found. Each molecule is disordered over two conformations. Two positions were defined for each atom of the Cl–(CH₂)₃– section of each molecule, except for the Cl-substituted C-atom, which is common to both conformations. Bond length restraints were applied to all bonds involving disordered atoms so as to maintain reasonable geometry. The best results were obtained with relative site occupation factors of 0.65:0.35 and 0.80:0.20 for the disordered components of molecules A and B, respectively.

⁶⁾ CCDC-286035–286040 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre, via http://www.ccdc.cam.ac.uk/data_request/cif

Table 3. Crystallographic Data of Compounds **11a**, **11b**, **12g**, **12h**, and **15**

	11a	11b	12g	12h	15
Crystallized from	toluene/CH ₂ Cl ₂ /MeCN/acetone	xylene/AcOEt/MeOH	CH ₂ Cl ₂ /acetone/i-PrOH	CH ₂ Cl ₂ /i-PrOH/acetone	CH ₂ Cl ₂ /Et ₂ O/hexane
Empirical formula	C ₁₀ H ₁₈ ClNO ₃	C ₈ H ₁₄ ClNO ₃	C ₁₁ H ₂₀ ClNO ₃	C ₁₂ H ₂₂ ClNO ₃	C ₂₁ H ₃₁ N ₃ O ₃
Formula weight [g mol ⁻¹]	235.71	207.66	249.74	263.76	373.49
Crystal color, habit	colorless, prism	colorless, plate	colorless, prism	colorless, prism	colorless, tablet
Crystal dimensions [mm]	0.07 × 0.17 × 0.25	0.02 × 0.15 × 0.20	0.13 × 0.20 × 0.25	0.18 × 0.20 × 0.25	0.12 × 0.15 × 0.30
Temp. [K]	160(1)	160(1)	160(1)	160(1)	160(1)
Crystal system	monoclinic	triclinic	triclinic	monoclinic	monoclinic
Space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> 2 ₁ / <i>n</i>
<i>Z</i>	4	4	2	4	4
Reflections for cell determination	18016	4764	13668	49886	4258
2 θ Range for cell determination [°]	4–60	4–55	4–55	4–60	4–52
Unit cell parameters					
<i>a</i> [Å]	5.8846(1)	9.5502(2)	8.1623(3)	7.0504(1)	10.1674(3)
<i>b</i> [Å]	10.5564(2)	10.3588(2)	8.8802(2)	15.0693(2)	19.4044(7)
<i>c</i> [Å]	19.0140(4)	11.3586(3)	9.0067(3)	13.0955(2)	11.5235(4)
α [°]	90	92.417(1)	86.145(2)	90	90
β [°]	91.887(1)	90.584(1)	88.338(2)	90.7171(8)	112.515(2)
γ [°]	90	109.313(1)	79.602(2)	90	90
<i>V</i> [Å ³]	1180.51(4)	1059.16(4)	640.56(4)	1391.22(3)	2100.2(1)
<i>D_x</i> [g cm ⁻³]	1.326	1.302	1.295	1.259	1.181
μ (MoK α) [mm ⁻¹]	0.312	0.338	0.291	0.272	0.0794
Scan type	ϕ and ω	ϕ and ω	ϕ and ω	ϕ and ω	ω
2 θ _(max) [°]	60	55	55	60	52
Transmission factors (min; max)	0.887; 0.979	–	0.855; 0.968	0.864; 0.957	–
Total reflections measured	32539	26265	15813	37833	37791
Symmetry independent reflections	3448	4824	2932	4054	4135
Reflections with <i>I</i> > 2 σ (<i>I</i>)	2459	3464	2535	3375	3006
Reflections used in refinement	3448	4822	2931	4052	4134
Parameters refined; restraints	148; 0	309; 10	159; 0	169; 0	256; 0
<i>R</i> (on <i>F</i> ; <i>I</i> > 2 σ (<i>I</i>) reflections)	0.0443	0.0667	0.0330	0.0370	0.0472
<i>wR</i> (on <i>F</i> ² ; all indept. reflections)	0.1158	0.2011	0.0833	0.0960	0.1132
Weighting parameters [<i>a</i> ; <i>b</i>] ^a)	0.0546; 0.3536	0.0996; 0.8346	0.0355; 0.2522	0.033; 0.7281	0.0406; 0.6032
Goodness-of-fit	1.046	1.069	1.055	1.055	1.048
Secondary extinction coefficient	–	–	0.019(5)	0.009(2)	0.015(2)
Final Δ_{\max}/σ	0.001	0.001	0.001	0.001	0.001
$\Delta\rho$ (max; min) [e Å ⁻³]	0.26; –0.32	0.92; –0.89	0.30; –0.25	0.28; –0.23	0.18; –0.19

^a) $w^{-1} = \sigma^2(F_o^2) + (aP)^2 + bP$ where $P = (F_o^2 + 2F_c^2)/3$

The OH and ammonium H-atoms in **12g** and **12h**, as well as the amide H-atom in **15**, and the amide and OH H-atoms in **11a** and **11b** were placed in the positions indicated by difference electron-density maps, and their positions were allowed to refine together with individual isotropic displacement parameters. All remaining H-atoms in all structures were placed in geometrically calculated positions and each was assigned a fixed isotropic displacement parameter with a value equal to $1.2U_{\text{eq}}$ of its parent atom ($1.5U_{\text{eq}}$ for the Me groups). The refinement of each structure F^2 was carried out on using full-matrix least-squares procedures, which minimized the function $\sum w(F_o^2 - F_c^2)^2$. Corrections for secondary extinction were applied, except in the cases of **11a** and **11b**. Neutral-atom-scattering factors for non-H atoms were taken from [41], and the scattering factors for H-atoms were taken from [42]. Anomalous dispersion effects were included in F_c [43]; the values for f' and f'' were those of [44]. The values of the mass attenuation coefficients are those of [45]. All calculations were performed using the SHELXL97 program [46].

We thank the analytical sections of our institute for the spectra and elemental analyses. Financial support of the Swiss National Science Foundation and F. Hoffmann-La Roche AG, Basel, is gratefully acknowledged.

REFERENCES

- [1] C. E. Ballard, H. Yu, B. Wang, *Curr. Med. Chem.* **2002**, *9*, 471.
- [2] J. S. Davies, *J. Peptide Sci.* **2003**, *9*, 471.
- [3] T. G. Back, *Tetrahedron* **1997**, *33*, 3041.
- [4] A. Yurek-George, F. Habens, M. Brimmell, G. Packham, A. Ganesan, *J. Am. Chem. Soc.* **2004**, *126*, 1030.
- [5] C. Palomo, M. Oiarbide, J. Garcia, A. Gonzalez, R. Razos, J. Ordiozola, P. Banuelos, M. Tello, A. Linden, *J. Org. Chem.* **2004**, *69*, 4126.
- [6] F. Sarabia, S. Chammaa, F. Lopez-Herrera, *Tetrahedron Lett.* **2002**, *43*, 2961.
- [7] W. L. Duax, J. F. Griffin, D. A. Langs, G. D. Smith, *Biopolymers*, **1996**, *40*, 141.
- [8] P. Vongvilai, M. Isaka, P. Kittakoop, P. Srikitkulchai, P. Kongsaree, S. Prabpai, Y. Thebtaranonth, *Helv. Chim. Acta* **2004**, *87*, 2066.
- [9] A. B. Mauger, *Top. Antibiot. Chem.* **1980**, *5*, 223.
- [10] D. Obrecht, H. Heimgartner, *Tetrahedron Lett.* **1983**, *24*, 1921.
- [11] D. Obrecht, H. Heimgartner, *Helv. Chim. Acta* **1987**, *70*, 329.
- [12] D. Obrecht, H. Heimgartner, *Helv. Chim. Acta* **1984**, *67*, 526.
- [13] J. E. F. Magirus, Ph.D. Thesis, University of Zurich, 1995.
- [14] K. N. Koch, A. Linden, H. Heimgartner, *Tetrahedron* **2001**, *57*, 2311.
- [15] P. Köttgen, planned Ph.D. Thesis, University of Zurich.
- [16] B. Iliev, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **2003**, *86*, 3215.
- [17] B. Iliev, A. Linden, H. Heimgartner, *Tetrahedron*, in press.
- [18] P. Wipf, H. Heimgartner, *Helv. Chim. Acta* **1990**, *73*, 13.
- [19] R. Miller, W. Theis, G. Heilig, S. Kirchmeyer, *J. Org. Chem.* **1991**, *56*, 1453.
- [20] B. Travis, R. Narayan, B. Borhan, *J. Am. Chem. Soc.* **2002**, *124*, 3824.
- [21] J. M. Schomaker, B. R. Travis, B. Borhan, *Org. Lett.* **2003**, *5*, 3089.
- [22] D. Obrecht, H. Heimgartner, *Helv. Chim. Acta* **1990**, *73*, 221.
- [23] H. Heimgartner, *Angew. Chem., Int. Ed.* **1991**, *30*, 238.
- [24] C. K. Johnson, 'ORTEP II', Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1976.
- [25] J. Bernstein, R. E. Davis, L. Shimoni, N.-L. Chang, *Angew. Chem., Int. Ed.* **1995**, *34*, 1555.
- [26] T. Joseph, J. Tam, M. Kitadani, Y. L. Chow, *Can. J. Chem.* **1976**, *54*, 3517.
- [27] S. Richard, G. Prie, A. Guignard, J. Thibonet, J. Luc Parrain, A. Duchene, M. Abarbri, *Helv. Chim. Acta* **2003**, *86*, 726.
- [28] R. M. Beesley, C. K. Ingold, J. F. Thorpe, *J. Chem. Soc.* **1915**, *105*, 1080.
- [29] M. Noack, R. Goettlich, *Eur. J. Org. Chem.* **2002**, 3171.

- [30] J. Iqbal, R. Srivastava, *Tetrahedron* **1991**, *47*, 3155.
- [31] S. Murahashi, Y. Makabe, K. Kunita, *J. Org. Chem.* **1988**, *53*, 4489.
- [32] K. Brannock, R. Burpitt, *J. Am. Chem. Soc.* **1961**, *83*, 3576.
- [33] J. Mihelcic, K. Moeller, *J. Am. Chem. Soc.* **2003**, *125*, 36.
- [34] E. Lee, C. Yoon, T. Lee, S. Kim, T. Ha, Y. Sung, S. Park, S. Lee, *J. Am. Chem. Soc.* **1998**, *120*, 7469.
- [35] J. P. Gouesnard, *Bull. Soc. Chim. Fr.* **1989**, 88.
- [36] R. Hooft, *KappaCCD Collect Software*, Nonius BV, Delft, The Netherlands, 1999.
- [37] Z. Otwinowski, W. Minor, in 'Methods in Enzymology', Vol. 276, 'Macromolecular Crystallography', Part A, Eds. C. W. Carter Jr., R. M. Sweet, Academic Press, New York, 1997, p. 307.
- [38] R. H. Blessing, *Acta Crystallogr., Sect. A* **1995**, *51*, 33.
- [39] A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori, M. Camalli, SIR92, *J. Appl. Crystallogr.* **1994**, *27*, 435.
- [40] A. L. Spek, PLATON, *J. Appl. Crystallogr.* **2003**, *36*, 7.
- [41] E. N. Maslen, A. G. Fox, M. A. O'Keefe, in 'International Tables for Crystallography', Ed. A. J. C. Wilson, Kluwer Academic Publishers, Dordrecht, 1992, Vol. C, Table 6.1.1.1, p. 477.
- [42] D. C. Creagh, W. J. McAuley, in 'International Tables for Crystallography', Ed. A. J. C. Wilson, Kluwer Academic Publishers, Dordrecht, 1992, Vol. C, Table 4.2.6.8, p. 219.
- [43] D. C. Creagh, J. H. Hubbell, in 'International Tables for Crystallography', Ed. A. J. C. Wilson, Kluwer Academic Publishers, Dordrecht, 1992, Vol. C, Table 4.2.4.3, p. 200.
- [44] R. F. Steward, E. R. Davidson, W. T. Simpson, *J. Chem. Phys.* **1965**, *42*, 3175.
- [45] J. A. Ibers, W. C. Hamilton, *Acta Crystallogr.* **1964**, *17*, 781.
- [46] G. M. Sheldrick, SHELXL97, Program for the Refinement of Crystal Structures, University of Göttingen, Germany, 1997.

Received October 12, 2005