


The First Stereoselective Total Synthesis of Naturally Occurring, Bioactive (3*R*,5*R*)-1-(4-Hydroxyphenyl)-7-phenylheptane-3,5-diol and the Synthesis of Its Enantiomer¹⁾

by Parigi Raghavendar Reddy, Chithaluri Sudhakar, Jayprakash Narayan Kumar, and Biswanath Das*

Organic Chemistry Division-I, CSIR-Indian Institute of Chemical Technology, Hyderabad-500 007, India
(phone: +91-40-7193434; fax: +91-40-7160512; e-mail: biswanathdas@yahoo.com)

The first stereoselective total synthesis of the naturally occurring anti-emetic diarylheptanoid (3*R*,5*R*)-1-(4-hydroxyphenyl)-7-phenylheptane-3,5-diol (**1**) was accomplished starting from 4-hydroxybenzaldehyde and involving a *Sharpless* kinetic resolution and an asymmetric epoxidation as the key steps (*Scheme 2*). The enantiomer **1a** of this compound was also simultaneously prepared.


Introduction. – Diarylheptanoids constitute an important class of natural products as they possess various interesting biological properties, such as anti-inflammatory, antioxidant, and anticancer activities [1]. A large number of these compounds have been isolated from nature [2]. Some of them contain a 1,3-diol system. The linear diarylheptanoid (3*R*,5*R*)-1-(4-hydroxyphenyl)-7-phenylheptane-3,5-diol (**1**) was isolated from *Alpinia officinarum* (Zingiberaceae) [3]. The compound exhibits anti-emetic activity. Thus, **1** is an important target for total synthesis. However, it is interesting that, though the synthesis of several stereoisomers and analogues of **1** have already been reported [4], the synthesis of **1** itself has not yet been disclosed. Here, we report the first stereoselective total synthesis of **1**. Its enantiomer **1a** was also simultaneously prepared.



Results and Discussion. – The retrosynthetic analysis (*Scheme 1*) revealed that compound **1** or **1a** can be synthesized from the acetonide derivative **2** or **2a**, respectively, and benzyltriphenylphosphonium ylide **3**. The compound **2** or **2a** can in turn be prepared from the achiral allyl alcohol **4** generated from 4-hydroxybenzaldehyde (**5**).

The present synthesis of **1** and **1a** was initiated (*Scheme 2*) by converting **5** [5] into its benzyl ether [5] followed by *Wittig* olefination with Ph₃PCHCOOEt and subsequent reduction with LiBH₄ to yield the known alcohol **6** [5]. The latter was oxidized under *Swern* conditions, and the corresponding aldehyde was treated with 4-methoxybenzyl (PMB)-protected propargyl alcohol (4-MeOC₆H₄CH₂OCH₂C≡CH) and BuLi to


¹⁾ Part 59 in the series 'Synthetic Studies on Natural Products'.

Scheme 1. Retrosynthetic Pathway of **1** and **1a**

furnish the acetylenic compound **7**. The reduction of **7** with LiAlH₄ in THF under reflux [6] afforded the (*E*)-configured allyl-alcohol **4** (major isomer) along with a small amount of its (*Z*)-isomer, which was separated by column chromatography. The *Sharpless* kinetic resolution [7] of **4** with ((+)-diisopropyl tartrate (+)-DIPT, titanium(IV) isopropoxide ((ⁱPrO)₄Ti), and *tert*-butyl hydroperoxide (^tBuOOH) resulted in the formation of the chiral epoxy alcohol **8a** and the allyl alcohol **9**. Next, compound **9** underwent *Sharpless* asymmetric epoxidation [5] with (–)-DIPT, (ⁱPrO)₄Ti, and ^tBuOOH to generate the enantiomeric epoxy alcohol **8**. Compounds **8** and **8a** were separately treated with *Red-Al*[®] (sodium bis(2-methoxyethoxy)aluminium hydride solution in toluene) in THF to give the diols **10** and **10a**, respectively. The two OH groups of these diols were protected by preparing the acetonide derivatives **2** and **2a**, and subsequently the protecting PMB group was removed with DDQ (=4,5-dichloro-3,6-dioxocyclohexol-1,4-diene-1,2-dicarbonitrile) to generate the alcohols **11** (from **10**) and **11a** (from **10a**). Each of the alcohols **11** and **11a** was again oxidized under *Swern* conditions, and the corresponding aldehydes were subjected to *Wittig* olefination with Ph₃P=CHPh to afford the olefins **12** and **12a**, respectively, along with minor amounts (10%) of their (*Z*)-isomers, which were separated by column chromatography. The protecting acetonide group of **12** and **12a** was removed by treatment with *p*-toluenesulfonic acid (TsOH) in MeOH followed by hydrogenation in the presence of Pd/C to yield the diols **1** and **1a**, respectively. The optical and spectroscopic (¹H-, and ¹³C-NMR and MS) properties of **1** were identical to those of the naturally occurring (3*R*,5*R*)-1-(4-hydroxyphenyl)-7-phenylheptane-3,5-diol [3]. The spectroscopic properties of **1a** were identical to those of **1**, but its optical-rotation value was reverse.

In conclusion, we developed the first stereoselective total synthesis of the anti-emetic compound (3*R*,5*R*)-1-(4-hydroxyphenyl)-7-phenylheptane-3,5-diol (**1**) starting from 4-hydroxybenzaldehyde. The synthesis of its enantiomer was also accomplished. The method can be utilized for the synthesis of related diarylheptanoids which can be employed for bioevaluation.

The authors thank UGC and CSIR, New Delhi, for financial assistance.

Scheme 2. Synthesis of **1** and **1a**


i) NaH, BnBr, THF, 0° to r.t., 4 h; 80%. ii) Ph₃P=CHCOOEt, CH₂Cl₂, r.t., 8 h; 82%. iii) LiBH₄, H₂O, Et₂O, 0° to r.t., 24 h; 91%. iv) 1. (COCl)₂, DMSO, Et₃N, -78°; 82%; 2. BuLi, dry THF, 4-MeOC₆H₄CH₂OCH₂C≡CH, -78°, 3.5 h; 90%. v) LiAlH₄, dry THF, 0° to reflux, 3 h; 78%. vi) (iPrO)₄Ti, (+)-DIPT, ^tBuOOH, dry CH₂Cl₂, -20°, 6 h; 46% (**8a**), 53% (**9**). vii) (iPrO)₄Ti, (-)-DIPT, ^tBuOOH, dry CH₂Cl₂, -20°, 7 h; 66%. viii) Red-Al[®], dry THF, 0° to r.t., 2 h; 72% (**10**), 74% (**10a**). ix) TsOH (cat.), 2,2-dimethoxypropane, CH₂Cl₂, r.t., 0.5 h; 95% (**2**), 94% (**2a**). x) DDQ, CH₂Cl₂/H₂O 19:1, r.t., 2 h; 63% (**11**), 64% (**11a**). xi) 1. (COCl)₂, DMSO, Et₃N, -78°, 3.5 h; 80%; 2. (Ph₃PCH₂Ph)Br, BuLi, dry THF, 4 h; 78% (**12**), 75% (**12a**). xii) 1. TsOH, MeOH, 3 h; 2. H₂, Pd/C, 2 h; 60% (**1**), 63% (**1a**).

Experimental Part

General. All commercially available reagents were used directly without further purification unless otherwise stated. The solvents used were all of AR grade and were distilled under dry N₂ where necessary. All reactions were performed in predried apparatus unless otherwise stated. Yields were those

of purified compounds unless otherwise stated. TLC: silica gel 60 F_{254} plates (SiO_2 ; Merck). Column chromatography (CC): silica gel (SiO_2 ; 60–120 mesh; Qingdao Marine Chemical, P. R. China). Optical rotations: Jasco-DIP-300 digital polarimeter. NMR Spectra: Gemini-200 spectrometer; in CDCl_3 ; δ in ppm rel. to Me_4Si as internal standard, J in Hz. ESI-MS: VG-Autospec-micromass instrument; in m/z . HR-MS: QSTAR-XL-Hybrid MS system (Applied Biosystems); in m/z .

6-[4-Methoxyphenyl)methoxy]-1-[4-(phenylmethoxy)phenyl]hex-4-yn-3-ol (**7**). To a stirred soln. of oxalyl chloride (8.07 ml, 132.20 mmol) in dry CH_2Cl_2 (160 ml), DMSO (0.55 ml, 18.2 mmol) was added at -78° , and the mixture was stirred at -78° for 0.5 h. A soln. of **6** (16 g, 66.11 mmol) in dry CH_2Cl_2 (60 ml) was added at -78° , and the mixture was stirred for 1.5 h at -78° . Et_3N (27.59 ml, 198.01 mmol) was added at 0° , and the mixture was stirred for an additional 30 min. After completion of the reaction, the mixture was diluted with H_2O (30 ml) and extracted with CH_2Cl_2 (2×100 ml). The combined org. layers were washed with brine (60 ml), dried (Na_2SO_4), and concentrated to give 3-[4-(phenylmethoxy)phenyl]propanal (13.01 g, 82%). Because of extensive oxidation of the aldehyde to the corresponding acid, we used it immediately for the next step.

Under N_2 , 1.6M BuLi in hexane (46.87 ml, 75 mmol) was added to a soln. of the PMB ether of propargyl alcohol (= 1-methoxy-4-[(prop-2-yn-1-yloxy)methyl]benzene; 13.2 g, 75 mmol) in dry THF (150 ml) at -78° , and the mixture was stirred for 30 min. Finally, a soln. of 3-[4-(phenylmethoxy)phenyl]propanal (12.0 g, 50.0 mmol) in dry THF (80 ml) was added and the mixture stirred for 3 h at -78° . After completion of the reaction, the mixture was quenched with sat. aq. NH_4Cl soln. (20 ml) and then extracted with AcOEt (2×100 ml), the extract dried (Na_2SO_4) and concentrated, and the residue purified by CC: pure **7** (1.74 g, 90%). $^1\text{H-NMR}$ (200 MHz): 7.37–7.12 (*m*, 7 H); 7.01 (*d*, $J = 8.0$, 2 H); 6.80–6.72 (*m*, 4 H); 4.90 (*s*, 2 H); 4.42 (*s*, 2 H); 4.29 (*t*, $J = 7.0$, 1 H); 4.06 (*s*, 2 H); 3.66 (*s*, 3 H); 3.29 (*br. s*, 1 H); 2.64 (*t*, $J = 7.0$, 2 H); 1.98–1.83 (*m*, 2 H). $^{13}\text{C-NMR}$ (50 MHz): 159.2; 156.9; 137.1; 133.4; 129.4; 129.2; 129.1; 128.3; 127.6; 127.2; 114.5; 113.3; 87.9; 80.6; 71.1; 69.6; 61.2; 56.9; 55.0; 39.6; 30.4. ESI-MS: 439 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{27}\text{H}_{28}\text{O}_4$ (416.51): C 77.89, H 6.73; found: C 77.94, H 6.68.

6-[4-(4-Methoxyphenyl)methoxy]-1-[4-(phenylmethoxy)phenyl]hex-4-en-3-ol (**4**). To the suspension of LiAlH_4 (6.57 g, 173.07 mmol) in dry THF (60 ml) under N_2 at 0° was added **7** (18.0 g, 43.26 mmol) in dry THF (75 ml), and the mixture was refluxed for 3 h. After the completion of the reaction, the mixture was cooled to 0° and diluted with Et_2O , and the reaction was quenched with sat. aq. Na_2SO_4 soln. (20 ml). When the effervescence subsided, the mixture was filtered through a pad of Celite and the pad washed with hot AcOEt (90 ml). The filtrate was washed with brine (2×30 ml), dried (Na_2SO_4), and concentrated, and the residue purified by CC: **4** (14.10 g, 78%). $[\alpha]_D^{25} = -0.42$ ($c = 1.0$, CHCl_3). $^1\text{H-NMR}$ (200 MHz): 7.42–7.23 (*m*, 5 H); 7.20 (*d*, $J = 8.0$, 2 H); 7.05 (*d*, $J = 8.0$, 2 H); 6.82 (*d*, $J = 8.0$, 2 H); 6.80 (*d*, $J = 8.0$, 2 H); 5.74–5.68 (*m*, 2 H); 4.99 (*s*, 2 H); 4.40 (*s*, 2 H); 4.17–4.11 (*m*, 1 H); 4.10–4.03 (*m*, 1 H); 3.98–3.85 (*m*, 1 H); 3.78 (*s*, 3 H); 2.68–2.55 (*m*, 2 H); 1.82–1.72 (*m*, 2 H). $^{13}\text{C-NMR}$ (50 MHz): 159.1; 157.2; 137.2; 135.7; 134.1; 130.2; 129.4; 129.3; 128.5; 127.8; 127.4; 114.5; 113.8; 72.0; 71.8; 69.9; 69.7; 55.2; 38.9; 31.0. ESI-MS: 441 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{27}\text{H}_{30}\text{O}_4$ (418.53): C 77.51, H 7.18; found: C 77.58, H 7.15.

(1S)-1-[(2S,3S)-3-[(4-Methoxyphenyl)methoxy]methyl]oxiran-2-yl]-3-[4-(phenylmethoxy)phenyl]propan-1-ol (**8a**). To a suspension of powdered molecular sieves (4 Å; 3 g) in dry CH_2Cl_2 (140 ml), $(i\text{-PrO})_4\text{Ti}$ (5.66 ml, 19.13 mmol), and (+)-DIPT (3.92 ml, 22.96 mmol) were added sequentially at -20° . After stirring for 30 min, **4** (16.0 g, 38.27 mmol) in dry CH_2Cl_2 (60 ml) was added, and stirring was continued for another 30 min at -20° . Then $t\text{-BuOOH}$ (3.50 ml, 21.05 mmol) was added and the stirring continued for another 5 h at -20° . After completion of the reaction, the mixture was quenched by the addition of H_2O (15 ml) and allowed to reach r.t. by stirring for 30 min. After recooling to 0° , 30% (*w/v*) aq. NaOH soln. (10 ml, sat. with brine) was added, and the mixture was stirred at 0° for 1 h. The mixture was extracted with Et_2O (2×40 ml), the combined org. extract washed with brine (2×20 ml), dried (Na_2SO_4), and concentrated, and the residue purified by CC: pure **9** (8.48 g, 53%) and **8a** (7.67 g, 46%). **8a**: $[\alpha]_D^{25} = -0.42$ ($c = 1.4$, CHCl_3).

Data of (3R)-6-[4-(4-Methoxyphenyl)methoxy]-1-[4-(phenylmethoxy)phenyl]hex-4-en-3-ol (**9**): $^1\text{H-NMR}$ (200 MHz): 7.42–7.28 (*m*, 5 H); 7.21 (*d*, $J = 8.0$, 2 H); 7.05 (*d*, $J = 8.0$, 2 H); 6.86 (*d*, $J = 8.0$, 2 H); 6.85 (*d*, $J = 8.0$, 2 H); 5.00 (*s*, 2 H); 4.50 (*d*, $J = 14.0$, 1 H); 4.41 (*d*, $J = 14.0$, 1 H); 3.79 (*s*, 3 H); 3.66–3.58 (*m*, 1 H); 3.48–3.39 (*m*, 1 H); 3.17–3.11 (*m*, 1 H); 3.10–3.02 (*m*, 1 H); 2.88–2.80 (*m*, 1 H);

2.77–2.58 (*m*, 2 H); 1.94 (*br. s*, 1 H); 1.90–1.69 (*m*, 2 H). $^{13}\text{C-NMR}$ (50 MHz): 159.7; 157.8; 137.9; 134.2; 130.0; 129.5; 129.3; 128.5; 127.9; 127.7; 115.0; 114.1; 73.1; 72.2; 70.0; 69.2; 58.3; 55.2; 35.5; 30.8. ESI-MS: 457 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{27}\text{H}_{30}\text{O}_5$ (434.52): C 74.65, H 6.91; found: C 74.78, H 6.88.

(1*R*)-1-[(2*R*,3*R*)-3-[(4-Methoxyphenyl)methoxymethyl]oxiran-2-yl]-3-[4-(phenylmethoxy)phenyl]propan-1-ol (**8**). As described for **8a**, with molecular sieves (4 Å, 2 g), CH_2Cl_2 (60 ml), $(i\text{PrO})_4\text{Ti}$ (5.66 ml, 19.11 mmol), (–)-DIPT (3.60 ml, 21.01 mmol), **9** (8 g, 19.13 mmol) instead of **4**, CH_2Cl_2 (50 ml), and $t\text{BuOOH}$ (7.97 ml, 47.84 mmol, stirring for 6 h). Extraction with Et_2O (2 × 80 ml). CC afforded pure **8** (5.50 g, 66%). $[\alpha]_{\text{D}}^{25} = +0.40$ ($c = 1.0$, CHCl_3). Spectroscopic data: identical to those of **8a**.

(2*S*,4*R*)-1-[(4-Methoxyphenyl)methoxy]-6-[4-(phenylmethoxy)phenyl]hexane-2,4-diol (**10**). To a stirred soln. of **8** (5.0 g, 11.52 mmol) in dry THF (50 ml) under N_2 at 0° was added *Red-Al*[®] soln. in toluene (6.84 ml, 34.55 mmol of 70% *w/w*) and the mixture was stirred at r.t. for 2 h. After completion of the reaction, the mixture was quenched with sat. aq. NH_4Cl soln. (20 ml) and then extracted with AcOEt (100 ml). The combined org. extract was washed with brine, dried (Na_2SO_4), and concentrated and the residue purified by CC: pure **10** (3.61 g, 72%). $[\alpha]_{\text{D}}^{25} = +3.2$ ($c = 2.3$, CHCl_3). $^1\text{H-NMR}$ (200 MHz): 7.40–7.23 (*m*, 5 H); 7.19 (*d*, $J = 8.0$, 2 H); 7.05 (*d*, $J = 8.0$, 2 H); 6.82 (*d*, $J = 8.0$, 4 H); 5.01 (*s*, 2 H); 4.48 (*d*, $J = 14$, 1 H); 4.42 (*d*, $J = 14.0$, 1 H); 4.08–4.02 (*m*, 1 H); 3.88–3.81 (*m*, 1 H); 3.78 (*s*, 3 H); 3.41 (*dd*, $J = 12.0$, 4.0, 1 H); 3.31 (*dd*, $J = 12.0$, 7.0, 1 H); 2.75–2.69 (*m*, 1 H); 2.66–2.59 (*m*, 1 H); 1.80–1.49 (*m*, 4 H). $^{13}\text{C-NMR}$ (50 MHz): 159.9; 157.2; 138.2; 134.1; 130.0; 129.9; 129.8; 129.0; 128.1; 127.8; 113.9; 113.1; 77.8; 73.4; 71.9; 70.5; 55.8; 39.9; 39.3; 30.5. ESI-MS: 459 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{27}\text{H}_{32}\text{O}_5$ (436.54): C 74.31, H 7.34; found: C 74.22, H 7.39.

(2*R*,4*S*)-1-[(4-Methoxyphenyl)methoxy]-6-[4-(phenylmethoxy)phenyl]hexane-2,4-diol (**10a**). As described for **10**: **10a** (74%). $[\alpha]_{\text{D}}^{25} = -2.9$ ($c = 2.0$, CHCl_3). Spectroscopic data: identical to those of **10**.

(4*S*,6*R*)-4-[(4-Methoxyphenyl)methoxymethyl]-2,2-dimethyl-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxane (**2**). To **10** (3.5 g, 8.02 mmol) in dry CH_2Cl_2 (30 ml), 2,2-dimethoxypropane (20 ml) was added followed by the addition of a cat. amount of TsOH. The mixture was stirred for 30 min at r.t. After completion of the reaction, sat. aq. NaHCO_3 soln. (10 ml) was added, the mixture extracted with CH_2Cl_2 (3 × 30 ml), the combined org. layer dried (Na_2SO_4) and concentrated, and the residue purified by CC: pure **2** (3.63 g, 95%). $[\alpha]_{\text{D}}^{25} = +10.67$ ($c = 2.9$, CHCl_3). $^1\text{H-NMR}$ (200 MHz): 7.41–7.22 (*m*, 5 H); 7.15 (*d*, $J = 8.0$, 2 H); 7.02 (*d*, $J = 8.0$, 2 H); 6.79 (*d*, $J = 8.0$, 2 H); 6.77 (*d*, $J = 8.0$, 2 H); 5.00 (*s*, 2 H); 4.50 (*d*, $J = 14.0$, 1 H); 4.41 (*d*, $J = 14.0$, 1 H); 4.05–3.97 (*m*, 1 H); 3.78 (*s*, 3 H); 3.66–3.60 (*m*, 1 H); 3.42–3.30 (*m*, 2 H); 2.66–2.59 (*m*, 1 H); 2.57–2.51 (*m*, 1 H); 1.75–1.49 (*m*, 4 H); 1.42–1.34 (*m*, 3 H); 1.33 (*s*, 3 H). $^{13}\text{C-NMR}$ (50 MHz): 159.2; 157.2; 137.3; 134.1; 129.4; 129.3; 129.2; 128.5; 127.7; 127.3; 114.9; 113.2; 100.3; 73.1; 72.8; 70.0; 66.9; 65.8; 55.0; 38.2; 35.2; 31.0; 25.1; 24.9. ESI-MS: 499 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{30}\text{H}_{36}\text{O}_5$ (476.60): C 75.63, H 7.56; found: C 75.74, H 7.53.

(4*R*,6*S*)-4-[(4-Methoxyphenyl)methoxymethyl]-2,2-dimethyl-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxane (**2a**). As described for **2**: **2a** (94%). $[\alpha]_{\text{D}}^{25} = -10.4$ ($c = 2.2$, CHCl_3). Spectroscopic data: identical to those of **2**.

(4*S*,6*R*)-2,2-Dimethyl-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxan-4-methanol (**11**). To a soln. of **2** (3.5 g, 7.35 mmol) in $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ 19:1 was added DDQ (5.0 g, 22.02 mmol) at r.t. in portions. The mixture was stirred for 2 h at r.t. and filtered. The filtrate was concentrated and the residue purified by CC: **11** (1.64 g, 63.0%). $[\alpha]_{\text{D}}^{25} = +2.7$ ($c = 0.9$, CHCl_3). $^1\text{H-NMR}$ (200 MHz): 7.96 (*d*, $J = 8.0$, 2 H); 7.42–7.31 (*m*, 5 H); 6.99 (*d*, $J = 8.0$, 2 H); 5.12 (*s*, 2 H); 4.38–4.32 (*m*, 1 H); 4.16–4.10 (*m*, 1 H); 3.71–3.65 (*m*, 1 H); 3.58–3.51 (*m*, 1 H); 2.76–2.69 (*m*, 1 H); 2.37–2.29 (*m*, 1 H); 1.90–1.52 (*m*, 4 H); 1.26 (*s*, 6 H). $^{13}\text{C-NMR}$ (50 MHz): 156.4; 141.0; 129.7; 129.6; 128.8; 128.1; 127.8; 114.9; 96.5; 74.0; 70.2; 69.9; 40.2; 40.0; 31.7; 27.1; 26.8. ESI-MS: 379 ($[M + \text{Na}]^+$). Anal. calc. for $\text{C}_{22}\text{H}_{28}\text{O}_4$ (356.46): C 74.16, H 7.87; found: C 74.24, H 7.81.

(4*R*,6*S*)-2,2-Dimethyl-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxan-4-methanol (**11a**). As described for **11**: **11a** (64%). $[\alpha]_{\text{D}}^{25} = -2.9$ ($c = 0.9$, CHCl_3). Spectroscopic data: identical to those of **11**.

(4*S*,6*R*)-2,2-Dimethyl-4-[(1*E*)-2-phenylethenyl]-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxane (**12**). To a stirred soln. of oxalyl chloride (0.5 ml, 8.42 mmol) in dry CH_2Cl_2 (20 ml), DMSO (0.38 ml, 12.62 mmol) was added at -78° , and the mixture was stirred at -78° for 0.5 h. A soln. of **11** (1.5 g, 4.21 mmol) in CH_2Cl_2 (10 ml) was added at -78° , and the mixture was stirred for 1.5 h at -78° Et_3N

(0.74 ml, 20.99 mmol) was added at 0°, and the mixture was stirred for an additional 30 min. After completion of the reaction, the mixture was diluted with H₂O (5 ml) and extracted with CH₂Cl₂ (2 × 20 ml) and the extract washed with brine (20 ml), dried (Na₂SO₄), and concentrated to give (4*S*,6*R*)-2,2-dimethyl-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxan-4-carboxaldehyde (1.3 g, 80%). Because of extensive oxidation of the aldehyde to the corresponding acid, we used it immediately for the next step.

To a soln. containing benzyltriphenylphosphonium bromide (1.5 g, 5.48 mmol) in dry THF (5 ml) was added dropwise 1.6*M* BuLi in hexane (2.52 ml, 4.03 mmol) *via* a syringe under N₂ at r.t. After stirring for 10 min, the preceding aldehyde (1.3 g, 3.67 mmol), diluted in dry THF, was dropwise added to the red-orange *Wittig* reagent. The mixture was stirred for 4 h at r.t. to afford a yellow soln. with a white precipitate. Then CH₂Cl₂ and sat. NH₄Cl soln. were added until pH 6. The aq. layer was extracted with CH₂Cl₂, the combined org. layer dried (MgSO₄) and concentrated, and the residue purified by CC: **12** (1.2 g, 78%). $[\alpha]_D^{25} = +5.6$ (*c* = 0.9, CHCl₃). ¹H-NMR (200 MHz): 7.99 (*d*, *J* = 8.0, 2 H); 7.53–7.48 (*m*, 4 H); 7.41–7.29 (*m*, 6 H); 7.10 (*d*, *J* = 8.0, 2 H); 6.88 (*dd*, *J* = 14.0, 7.0, 1 H); 6.28 (*d*, *J* = 14.0, 1 H); 5.29 (*s*, 2 H); 4.65–4.58 (*m*, 1 H); 3.72–3.65 (*m*, 1 H); 2.63–2.56 (*m*, 2 H); 1.73–1.49 (*m*, 4 H); 1.42 (*s*, 6 H). ¹³C-NMR (50 MHz): 157.0; 141.8; 137.4; 134.7; 130.2; 129.8; 129.7; 128.9; 128.2; 128.0; 114.9; 96.5; 70.2; 68.8; 68.5; 39.8; 39.4; 31.6; 31.0. ESI-MS: 451 ($[M + Na]^+$). Anal. calc. for C₂₉H₃₂O₃ (428.56): C 81.31, H 7.48; found: C 81.25, H 7.54.

(4*R*,6*S*)-2,2-Dimethyl-4-[(1*E*)-2-phenylethenyl]-6-[2-[4-(phenylmethoxy)phenyl]ethyl]-1,3-dioxane (**12a**). As described for **12**: **12a** (75%). $[\alpha]_D^{25} = -5.2$ (*c* = 1.5, CHCl₃). Spectroscopic data: identical to those of **12**.

(3*R*,5*R*)-1-(4-Hydroxyphenyl)-7-phenylheptane-3,5-diol (**1**). A soln. of **12** (200 mg, 2.80 mmol) in MeOH (5 ml) in the presence of TsOH (10 mg) was stirred for 3 h. After the completion of the reaction, concentration afforded the crude diol. This residue of unsaturated diol in MeOH (15 ml) in the presence of a cat. amount of Pd/C was stirred for 2 h under H₂ (1 atm). After filtration and concentration, the crude residue was purified by CC: **1** (82 mg, 60%). $[\alpha]_D^{25} = +9.2$ (*c* = 1.1, CHCl₃). ¹H-NMR (200 MHz): 7.26–7.12 (*m*, 5 H); 6.93 (*d*, *J* = 8.0, 2 H); 6.71 (*d*, *J* = 8.0, 2 H); 4.02–3.89 (*m*, 2 H); 2.81–2.50 (*m*, 4 H); 1.89–1.69 (*m*, 4 H); 1.60 (*t*, *J* = 6.0, 2 H). ¹³C-NMR (50 MHz): 154.2; 142.0; 133.4; 129.4; 128.4; 128.3; 126.0; 115.5; 69.0; 68.9; 42.4; 39.1; 39.0; 32.1; 31.2. ESI-MS: 301 ($[M + H]^+$). Anal. calc. for C₁₉H₂₄O₃ (300.39): C 76.00, H 8.00; found: C 76.18, H 8.06.

(3*S*,5*S*)-1-(4-Hydroxyphenyl)-7-phenylheptane-3,5-diol (**1a**). As described for **1**: **1a** (63%). $[\alpha]_D^{25} = -10.2$ (*c* = 1.3, CHCl₃). Spectroscopic data: identical to those of **1**.

REFERENCES

- [1] K.-S. Chun, K.-K. Park, J. Lee, M. Kang, Y.-J. Surh, *Oncol. Res.* **2002**, *13*, 37; J. Ishida, M. Kozuka, H. Tokuda, H. Nishino, S. Nagumo, K.-H. Lee, M. Nagai, *Bioorg. Med. Chem.* **2002**, *10*, 3361; P. N. Yadav, Z. Liu, M. M. Rafi, *J. Pharmacol. Exp. Ther.* **2003**, *305*, 925; H.-J. Kim, S.-H. Yeom, M.-K. Kim, J.-G. Shim, I.-N. Paek, M.-W. Lee, *Arch. Pharmacol. Res.* **2005**, *28*, 177; H. Mohamad, N. H. Lajis, F. Abas, A. M. Ali, M. A. Sukari, H. Kikuzaki, N. Nakatani, *J. Nat. Prod.* **2005**, *68*, 285; K. Akiyama, H. Kikuzaki, T. Aoki, A. Okuda, N. H. Lajis, N. Nakatani, *J. Nat. Prod.* **2006**, *69*, 1637; H. Matsuda, S. Ando, T. Kato, T. Morikawa, M. Yoshikawa, *Bioorg. Med. Chem.* **2006**, *14*, 138.
- [2] K. K. Lee, B. D. Bahler, G. A. Hofmann, M. R. Mattern, R. K. Johnson, D. G. I. Kingston, *J. Nat. Prod.* **1998**, *61*, 1407.
- [3] S. Uehara, I. Yasuda, K. Akiyama, H. Morita, K. Takeya, H. Itokawa, *Chem. Pharm. Bull.* **1987**, *35*, 3298; D. Shin, K. Kinoshita, K. Koyama, K. Takahashi, *J. Nat. Prod.* **2002**, *65*, 1315.
- [4] M. Narasimhulu, T. S. Reddy, K. C. Mahesh, A. S. Krishna, J. V. Rao, Y. Venkateswarlu, *Bioorg. Med. Chem. Lett.* **2009**, *19*, 3125; V. U. Pawar, V. S. Shinde, *Tetrahedron: Asymmetry* **2011**, *22*, 8; S. P. Reddy, K. Ashalatha, D. K. Reddy, B. C. Babu, Y. Venkateswarlu, *Synthesis* **2011**, 3180; S. Bhosale, V. P. Vyavahare, U. V. Prasad, V. P. Palle, D. Bhuniya, *Tetrahedron Lett.* **2011**, *52*, 3397.
- [5] Y. Gao, J. M. Klunder, R. M. Hanson, H. Masamune, S. Y. Ko, K. B. Sharpless, *J. Am. Chem. Soc.* **1987**, *109*, 5765; B. Das, K. Suneel, G. Satyalakshmi, D. N. Kumar, *Tetrahedron: Asymmetry* **2009**, *20*, 1536.

- [6] J. Cossy, C. Meyer, M. Defosseux, N. Blanchard, *Pure Appl. Chem.* **2005**, 77, 1131.
[7] T. Katsuki, K. B. Sharpless, *J. Am. Chem. Soc.* **1980**, 102, 5974; B. Das, B. Veeranjanyulu, P. Balasubramanyam, M. Srilatha, *Tetrahedron: Asymmetry* **2010**, 21, 2762.

Received March 29, 2012