

Substituted 1,3,2,4-Benzodithiadiazines: Novel Derivatives, By-Products, and Intermediates*

Alexander Yu. Makarov, Irina Yu. Bagryanskaya, Yuri V. Gatilov, Tatiana V. Mikhailina, Makhmut M. Shakirov, Lyudmila N. Shchegoleva, and Andrey V. Zibarev

Institute of Organic Chemistry, Russian Academy of Sciences, 630090 Novosibirsk, Russia

Received 6 September 2000; revised 19 January 2001

ABSTRACT: The synthesis of the title compounds **1** by 1:1 condensation of $Ar-N=S=N-SiMe_3$ **2** with SCl_2 followed by intramolecular ortho-cyclization of each $[Ar-N=S=N-S-Cl]$ intermediate is complicated by further reaction of **1** with SCl_2 to give Herz salts **3**. With the **2**: SCl_2 ratio of 2:1, the formation of by-products **3** is reduced and novel compounds **1** are accessible. With ortho-I containing starting material **2j**, the parent compound **1s** is obtained as the result of an unexpected I, not H, substitution. The rate of the **1** + SCl_2 reaction depends upon a substituent's position, and the minor 8-R isomers **1l,p** (R = Br, I) are isolated for the first time from mixtures with the major 6-R isomers due to reduced reactivity toward SCl_2 . The synthesized compounds **1–3** are characterized by multinuclear (including nitrogen) NMR and X-ray crystallography. According to the X-ray diffraction data, **1j** (6-Br) and **1k** (7-Br) derivatives are planar, whereas **1i** (5-Br) and **1l** (8-Br) are bent along the $S1 \cdots N4$ line by $\sim 5^\circ$ and $\sim 4^\circ$, respectively, and the **1r** (7- OCH_3) derivative is planar in contrast to the known 5- OCH_3 isomer, which possesses a significantly folded hetero-

cycle. The distortion of the planar geometry of some compounds **1** is interpreted in terms of a pseudo-Jahn-Teller effect as the result of π -highest occupied molecular orbital (HOMO) – σ^* -(LUMO) lowest unoccupied molecular orbital + 1 mixing in a planar conformation. The **2p** compound is the first structurally defined $Ar-N=S=N-SiMe_3$ azathiene. The compound $Ar-N=S=N-S-NH-Ar$ **6** modeling the aforementioned intermediate has been isolated and structurally characterized. We describe the attempts to synthesize compounds **1** from 2-aminobenzenethiols and $(SN)_4$ and from salts **3** and Me_3SiN_3 , and we discuss the reaction pathways. © 2001 John Wiley & Sons, Inc. Heteroatom Chem 12:563–576, 2001

INTRODUCTION

Recently, various 5-R, 6-R and 7-R substituted 1,3,2,4-benzodithiadiazines (**1**) (Scheme 1) were synthesized by 1:1 condensation of $RC_6H_4-N=S=N-SiMe_3$ (**2**) with SCl_2 followed by intramolecular electrophilic ortho-cyclization [3]. These heterocycles reveal some formal features of antiaromaticity [4] along with high and varied, essentially unpredictable, heteroatom reactivity [1,3,5,6] thus provoking keen interest. However, their chemistry has been only briefly investigated. In particular, it was found that mild thermolysis of **1** in dilute hydrocarbon solutions resulted in practically quantitative yields of stable 1,2,3-benzodithiazolyl π -radicals [1], interesting as potential building blocks in the design and

*Part XV of the series "Cyclic Aryleneazachalcogenenes"; part XIV: Ref. [1], part XIII: Ref. [2].

Correspondence to: Andrey V. Zibarev. E-mail: zibarev@nioch.nsc.ru

Contract Grant Sponsor: Russian Foundation for Basic Research.

Contract Grant Number: 99-03-33115.

Contract Grant Number: 96-03-40001.

Contract Grant Number: 96-07-89187.

© 2001 John Wiley & Sons, Inc.

synthesis of organic molecules with high-spin ground states (molecular magnets) [7, and references there-in]. On the whole, it is believed that many new reactions will be observed and many new structural types will be found among the reaction products of **1**.

In some cases the preparations of compounds **1** from **2** [3] were unsuccessful, especially with *para*-substituted starting materials (precursors of 7-R isomers of **1**), independent of the character of each substituent R. The 8-R substituted compounds **1**, minor products of regioselective cyclizations of *meta*-substituted starting materials leading predominantly to 6-R isomers, were characterized only spectroscopically without isolation [3].

The present article deals with the improved synthesis of 7-R and some other derivatives of **1** including the first-ever isolation of minor 8-R isomers, as well as with the by-products and intermediates of the cyclization. Attempts to synthesize **1** by different approaches to the ring closure are also described.


RESULTS AND DISCUSSION

Precursors

The starting materials **2** bearing a strong donor (Me_2N) or a strong acceptor (NO_2) substituent R, as well as an I substituent, were of special interest [8].

The novel compounds **2** were synthesized from $\text{LiN}(\text{SiMe}_3)_2$ and corresponding $\text{RC}_6\text{H}_4\text{-N}=\text{S}=\text{O}$ (cf. [3,9]) or $\text{RC}_6\text{H}_4\text{-N}=\text{SCl}_2$ ($\text{R}=\text{NO}_2$) (Scheme 2). Surprisingly, both approaches were not effectual with $\text{R} = 3\text{-NO}_2$ and the first one with $\text{R} = 2\text{-Me}_2\text{N}$ or $4\text{-Me}_2\text{N}$.


The **2p** compound (Figure 1) is the first structurally defined $\text{Ar-N}=\text{S}=\text{N-SiMe}_3$ azathiene [**2p**: $\text{Ar} = 4\text{-(4'-IC}_6\text{H}_4\text{)C}_6\text{H}_4$]. In the crystal, **2p** adopts the *Z,E* configuration with a dihedral angle between the NSN plane and the adjacent ring of $14.3(2)^\circ$, and between the rings of $29.9(1)^\circ$. The S^1N^2 bond distance (Figure 1) is rather short but corresponds to those found in the related (*Z,E*) $\text{Ph}_3\text{Si-N}=\text{S}=\text{N-SiPh}_3$ compound (1.506 and 1.508 Å) [10]. Other bond lengths and bond angles are typical [11].



SCHEME 1

Improved Cyclization

The synthesis of **1** by a 1:1 reaction of **2** with SCl_2 is complicated by formation of Herz salts (**3**) (1,2,3-arenodithiazolium chlorides [12]). In some cases they are even the major products. Control experiments (see Cyclic By-Products) indicate that the salts **3** come from further reaction of **1** with SCl_2 . It follows that the 2: SCl_2 ratio should be changed in favor of **2** to suppress this side-process while preparing **1**. Indeed, with the 2: SCl_2 ratio of 2:1 it is possible to



SCHEME 2 Compounds 2:

I/R	Me_2N	NO_2	Br	I	F	OCH_3
2	(a)	d	g	j		
3	b	(e)	h	k		
4	(c)	f	i	l	m	n

3,5- Br_2 : **2o**, 4-(4'- IC_6H_4): **2p**

Parentheses indicate that corresponding compound was not synthesized by this way.





FIGURE 1 Structure of molecule **2p** in the crystal. Selected bond lengths (Å) and bond angles ($^\circ$): N^1S^1 1.513(3); S^1N^2 1.508(3); N^1C^1 1.417(5); N^2Si^1 1.740(3); I^1C^{10} 2.096(4); $\text{C}^1\text{N}^1\text{S}^1$ 130.3(2); $\text{N}^1\text{S}^1\text{N}^2$ 119.0(2); $\text{S}^1\text{N}^2\text{Si}^1$ 127.7(2).

obtain 7-R derivatives of **1** from the cyclizations previously unsuccessful [3] with the 1:1 ratio (i.e., **1q,r**; R = F, OCH₃) as well as to prepare several novel derivatives (Scheme 3).

With 4-RC₆H₄-N=S=N-SiMe₃, the target heterocycle formation proceeds also with R = Br, I, 4-IC₆H₄. On the other hand, with R = NO₂ or CF₃ the expected compounds are not obtained under the improved reaction conditions (cf. Ref. [3]).

In the case of 2-RC₆H₄-N=S=N-SiMe₃ the cyclization readily occurs with R = Br. With starting material **2j** (R = I) the parent 1,3,2,4-benzodithiadiazine **1s** was isolated from the reaction mixture instead of the expected 5-I derivative **1m**. One can think that in this case the ring closure is mediated by the bipolar ion (Scheme 3) [13] originated from initial electrophilic attack on iodine, rather than on carbon [14]. The postulated (Scheme 3) ICl by-product should react further with both **2j** and **1s**. At any rate, control experiments have shown that Cl₂ is highly reactive toward **1** to give a complex mixture of unidentified products. With R = NO₂, the cyclization failed.

Earlier [3], it was discovered that for 3-RC₆H₄-N=S=N-SiMe₃ (R = CH₃, OCH₃, F, Cl) the cycliza-



SCHEME 3 Heterocycles 1:

n/R	Me ₂ N	NO ₂	Br	I
5	(a)	(e)	i	(m)
6	(b)	(f)	j	n
7	(c)	(g)	k	o
8	(d)	(h)	l	p

R = 7-F, **1q**; 7-OCH₃, **1r**; H, **1s**; 6,8-Br₂, **1t**; 7-(4-IC₆H₄), **1u**
 Parentheses indicate that corresponding heterocycle was not synthesized by this way.

tion is highly regioselective, leading predominantly to 6-R substituted compounds **1**, with the preferred direction of ring closure being consistent with the thermodynamics of the corresponding σ -complexes as well as the factors of kinetic control for orbital-controlled EI-Nu interaction. In this work, for R = Br, the ratio of the major 6-Br (**1j**) isomer to the minor 8-Br (**1l**) one was found to be 80:20. For R = I, however, the ratio of 6-I (**1n**) and 8-I (**1p**) isomers was only 60:40.

The observed ratio of isomers would be affected by their further reaction with SCl₂ at different rates. Control experiments on available mixtures of 6-R and 8-R isomers (R = Br, I) have shown that the 6-R isomer reacts with SCl₂ much more rapidly than the 8-R isomer, which makes possible the eventual isolation of 8-Br and 8-I derivatives. Thus, the cyclization itself seems to be more regioselective than it appears from the observed ratios of isomers in the reaction mixtures presented in [3] and this work.

With R = Me₂N (**2b**) or CF₃ the cyclization failed.

Molecular Structures

According to the previous data of X-ray crystallography, the parent compound **1s** [6] and its 5-CF₃ and 6-F derivatives [3] are planar, whereas 5-OCH₃, 6-CH₃ [3], and 5,6,7,8-F₄ [15] derivatives are bent along the S1...N4 line by $\sim 11^\circ$, $\sim 7^\circ$, and 5.5° , respectively. The MP2/6-31G** calculations do not reproduce this intriguing structural dichotomy with **1s**, its 5-OCH₃ and 5,6,7,8-F₄ derivatives, and the model 1,3,2,4-dithiadiazine (**11**) being found to be bent along the S1...N4 line by $\sim 38^\circ$, $\sim 39^\circ$, $\sim 34^\circ$, and $\sim 45^\circ$ respectively. On the other hand, B3LYP/6-31G* calculations [2] predict a bent (26°) conformation for a free molecule of **1s**, but a perfectly planar conformation for its 5,6,7,8-F₄ derivative.

In the present work, both types of molecular geometry were observed again in the crystal (Figure 2, Table 1). For example, among **1i-l** isomers, **1j** (6-Br) and **1k** (7-Br) are planar, whereas **1i** (5-Br) and **1l** (8-Br) are bent along the S1...N4 line by $5.3(3)^\circ$ and $3.8(3)^\circ$, respectively (throughout this work the term *planar* conventionally means that the angle above is smaller than 3°). Furthermore, for two crystallographically independent molecules of **1j**, one is perfectly planar, whereas another is bent by $3.1(2)^\circ$, which directly indicates the importance of packing effects. The **1r** (7-OCH₃) derivative (Figure 2, Table 1) is planar, in contrast to the significantly folded 5-OCH₃ isomer (see previous paragraph). Thus, the type of the molecular geometry of the title compound in the crystal, planar or bent, depends




FIGURE 2 Structure of molecules **1i-l,r** in the crystal. For selected bond lengths and bond angles, see Table 1.

not only on the substituent character, but also on its position.

The bond lengths and bond angles of **1i-l,r** are typical [3,11]. Within experimental accuracy they are the same for all molecules (Table 1). In the N=S=N fragment the bond lengths differ except for **1l** (Table 1).

It can be assumed that the distortion of the planar C_s geometry in the calculations reflects the tendency of the molecules to minimize thermodynamic destabilization associated with antiaromaticity via a pseudo-Jahn-Teller effect [16]. With a simplifying

molecular orbital (MO) model, one can identify orbital interactions capable of reducing molecular symmetry of the compounds under consideration from C_s to C_1 . With C_s symmetry the molecular ground state is $(\dots\pi^2)$, namely, ${}^1A'$. It follows from the calculations (Table 2) that for both **1s** and **1l** the C_s structure is the TS of the molecular bending. The transition to C_1 symmetry is possible by vibronic interaction with excited ${}^1A''$ states achievable by $\pi \rightarrow \sigma^*$ and $\sigma \rightarrow \pi^*$ excitations. For **1s** the lowest-energy excitation of these types is $6a'' \rightarrow 20a'$. The mixing of $6a''$ (π -highest occupied molecular

TABLE 1 Selected Bond Lengths (Å) and Bond Angles (°) in Molecules **1i-1r**^a

	1i	1j (two independent molecules)	1k	1l	1r
S ¹ N ²	1.674(8)	1.684(5), 1.679(5)	1.664(4)	1.684(5)	1.678(3)
N ² S ³	1.547(8)	1.543(4), 1.548(5)	1.546(4)	1.538(5)	1.538(3)
S ³ N ⁴	1.526(4)	1.527(4), 1.533(4)	1.524(4)	1.533(3)	1.528(3)
N ⁴ C ¹⁰	1.413(7)	1.425(5), 1.417(5)	1.406(5)	1.426(7)	1.421(4)
S ¹ C ⁹	1.779(6)	1.785(4), 1.787(4)	1.787(3)	1.788(4)	1.785(3)
C ⁹ C ¹⁰	1.413(8)	1.401(6), 1.400(6)	1.408(5)	1.405(6)	1.402(4)
C ⁹ S ¹ N ²	106.5(3)	105.6(2), 106.0(2)	105.8(2)	105.7(2)	106.1(2)
S ¹ N ² S ³	122.9(3)	123.5(2), 123.3(3)	123.8(2)	123.8(3)	123.1(2)
N ² S ³ N ⁴	119.2(3)	119.3(2), 119.2(2)	118.8(2)	118.9(2)	119.3(2)
S ³ N ⁴ C ¹⁰	123.3(4)	123.0(3), 122.9(3)	123.7(3)	123.3(3)	123.5(2)
N ⁴ C ¹⁰ C ⁹	124.4(5)	124.2(4), 124.9(4)	124.2(3)	124.1(4)	123.8(3)
C ¹⁰ C ⁹ S ¹	123.2(4)	124.2(3), 123.7(3)	123.7(3)	123.9(4)	124.0(2)

^aFor atom numbering, see Figure 1.**TABLE 2** Quantum Chemical Calculations Data for Molecules **1s** and **11**

Compound (symmetry)	RHF/6-31G* - E _{tot} (PES point) ^a (a.u.)	E _{rel} (kcal mol ⁻¹)	MP2/6-31G* - E _{tot} (PES point) ^a (a.u.)	E _{rel} (kcal mol ⁻¹)
11 (C _s)	980.664074 (TS)	0.86		
11 (C ₁)	980.665451 (min)	0		
1s (C _s)	1133.337707 (TS)	0.28	1134.702832 (TS)	2.43
1s (C ₁)	1133.338154 (min)	0	1134.706710 (min)	0

^aThe types of stationary PES points were determined by the Hessian calculations.

orbital (HOMO)) with 20a' (the lowest virtual σ*-MO, which is next to the 7a'' π-lowest unoccupied molecular orbital (LUMO), i.e., LUMO + 1) (Figure 3) causes the molecule to fold along the S1...N4 line in accordance with the results of MP2/6-31G** calculations and the data of X-ray crystallography for 5-OCH₃, 5-Br, 6-CH₃, 8-Br, and 5,6,7,8-F₄ derivatives. According to the atomic orbital (AO) contributions to the 6a'' and 20a' MOs (Figure 3), the density of the corresponding one-electron transition is localized mainly on S1. The energy gain on going from C_s to C₁ symmetry is smaller by 1 kcal mol⁻¹ at the RHF level of theory becoming about nine-fold higher (for **1s**) at the MP2 level (Table 2).

Spectral Properties

In accordance with previous data [3], in ¹⁵N NMR spectra of the title compounds, δ¹⁵N values lie inside a narrow range, ~280–250 ppm, demonstrating weak dependence upon position and character of a carbocyclic substituent (for details, see Table 6 in Experimental). In the ¹⁵N{¹H} spectrum of the parent compound **1s**, the signal at 263.1 ppm (Table 3) is a


**FIGURE 3** The 6a'' π-HOMO and 20a' σ*-LUMO + 1 (see text) of C_s-**1s** and the corresponding MOs of C_s-**11** (RHF/6-31G*).

TABLE 3 Assignment of ^1H , ^{13}C , and ^{15}N NMR Spectra of **1s**^a

δ (J, Hz)	δ (J, Hz)
N^2 269.2 (s)	C^{8a} 115.3 (tdd, 8.5, 1.7, 0.8)
N^4 263.1 (d, 2.3)	C^{4a} 138.5 (tt, 9.3, 2.1)
H^5 5.90 (dd, 7.8, 1.3)	C^5 123.0 (ddt, 163.2, 8.5, 1.3)
H^6 6.63 (tm, 7.8)	C^6 133.2 (ddm, 164.8, 8.5)
H^7 6.79 (tm, 7.6)	C^7 130.5 (ddm, 164.5, 7.2)
H^8 5.79 (dd, 7.5, 1.2)	C^8 124.0 (dddd, 161.5, 8.9, 2.1, 1.3)

^aFor atom numbering, see Scheme 1.

doublet under selective decoupling with H^8 , but a singlet under selective decoupling with H^5 , thus proving the empirical assignment given in [3]. Table 3 provides also the complete assignment of the ^1H and ^{13}C NMR spectra of **1s** based on double resonance, COLOC, and off-resonance techniques.




The long-wave absorption maxima in the UV–vis spectra of the title compounds are found in the range of ~ 610 – 635 nm (cf. [3]), again with weak dependence upon the position and character of the substituent R (for details, see Table 6).

Cyclic By-Products

The Herz salts **3**, are cyclic by-products. Special preparative experiments as well as ^{14}N NMR identification of thiazyl chloride NSCl in the reaction mixtures suggest that it is possible to explain the formation of **3** by further reaction of **1** with SCl_2 (Scheme 4). One can postulate singlet 1,2,3-benzodithiazol-2-yl-nitrene (**5**, an isomer of **1**) to be the key intermediate of the reaction of **1** with SCl_2 . Then, the reaction seems to be an oxidative imination of SCl_2 with **5** followed by an elimination of NSCl from the product to give the final salt **3** (Scheme 4) [17]. Previously, similar elimination of NSCl has been observed during thermal decomposition of $\text{PhSO}_2\text{N}=\text{SCl}_2$ [18] (Scheme 4). It follows from control experiments that NSCl (taken as its trimer) also reacts with **1** to give Herz salts.

A thermodynamic gain of transformation of an antiaromatic system into an aromatic system seems to be a driving force of the reaction of **1** with SCl_2 . With planar geometry [19] and 10π -electrons, the cations of **3** possess aromaticity, although reduced as compared with the symmetric isomer **4** [20,21a].

Previously described 1:1 addition of **1** to PPh_3 that afforded a derivative of (1,2,3-benzodithiazol-2-yl)iminophosphorane (Scheme 4) [5] can also be interpreted as an oxidative imination of the phosphine with nitrene **5**. Thus, the reactions of **1** with formal both an electrophile (SCl_2) and a nucleophile (PPh_3)



SCHEME 4 Salts **3**: R = H (**a**), 4-OCH₃ (**b**), 6-OCH₃ (**c**), 5-Br (**d**), 5-CH₃ (**e**), 5-I (**f**)

probably proceed in a similar way. Formation of 1,2,3-benzodithiazolyls by thermolysis of **1** [1] is very likely mediated by **5** as well [17a].

Me_3SiCl is inert toward **1** under conditions of the cyclization, in contrast to its reaction with the related 1,3,5,2,4-benzotrithiadiazepine, leading to 1,3,2-benzodithiazolium chloride (**4**) [23], isomeric with **3a**.

A Model for the Cyclization Intermediate

Compound 4-FC₆H₄-N=S=N-S-NH-C₆H₄F-4 (**6**), actually isolated from a reaction of 4-FC₆H₄-N=S=N-SiMe₃ (**2m**) with salt **3a**, can formally be considered as a [4-FC₆H₄-N=S=N-S-Cl] intermediate in the synthesis of **1q** (Scheme 3), trapped by the parent amine 4-FC₆H₄NH₂. According to the data of X-ray crystallography (Figure 4), in the crystal this model compound adopts the *E,Z* configuration (cf. *Z,E* configuration of Ar-N=S=N-SiMe₃ **2p**, Figure 1). As the cyclization requires a *Z,Z* configuration, one can conclude that the model belongs to the early stages of the process. The compound **6** features considerable nonplanarity. The molecule consists of three planar fragments: ring C1-C6 (A plane, standard deviation 0.005 Å), chain NSNSN (B plane, standard deviation 0.74 Å), and ring C7-C12 (C plane, standard deviation 0.003 Å) with dihedral angles A-B, B-C, and A-C of 101.4°, 30.7° and 71.6°, respectively. The bond lengths and bond angles of




FIGURE 4 Structure of molecule **6** in the crystal. Selected bond lengths (Å) and bond bond angles (°): N¹S¹ 1.639(4); S¹N² 1.690(3); N²S² 1.549(3); S²N³ 1.551(3); C¹N¹ 1.402(5); C⁷N³ 1.415(4); C¹N¹S¹ 125.2(3); N¹S¹N² 101.2(2); S¹N²S² 118.5(2); N²S²N³ 110.0(2); S²N³C⁷ 119.4(3).

the sulfur-nitrogen fragment of **6** are typical (cf. the nearest analog Ph–N=S=N–S–Ph [24]).


Attempted Approaches to the Ring Closure

An attempt was made to synthesize **1** by reaction of 2-aminobenzenethiols with (SN)₄ as a source of the sulfur-nitrogen fragment necessary to the ring closure. However, the reaction proceeded only as oxidation of the thiol to the corresponding disulfide (Scheme 5).

Another approach tried was based on the reaction of a Herz salt **3** with Me₃SiN₃. It was believed that the reaction will lead via the corresponding covalent azide to a singlet nitrene **5** capable of isomerization into a compound **1** (cf. singlet nitrenes Ar–N: [25] and Ar–S–N: [26,27]). However, only persistent 1,2,3-benzodithiazolyl radicals **8** [1] were identified as reaction products (Scheme 5; cf. [27]). It should be noted that formation of a Herz-type ionic azide is also possible in this system followed by a redox process directly leading to an **8** radical [21b].

EXPERIMENTAL

The ¹H, ¹³C, ¹⁴N, and ¹⁵N NMR spectra were measured on a Bruker DRX-500 spectrometer at frequencies of 500.13, 125.76, 36.13, and 50.68 MHz,



SCHEME 5

respectively, with standards tetramethylsilane (TMS) and NH₃ (liq.); the ¹⁹F NMR spectra were taken on a Bruker AC-200 machine at a frequency of 188.28 MHz with C₆F₆ as the standard. The mass spectra (EI, 70 eV) were collected on a Finnigan MAT MS-8200 instrument, and the UV–vis spectra were recorded on Specord M40 and HP 8453 spectrophotometers.

The X-ray structure determinations (Table 4) were carried out on a Syntex P2₁ diffractometer using Cu Kα radiation (compounds **1k,r, 6**) and on a Bruker P4 diffractometer using Mo Kα radiation (compounds **1i,j,l, 2p**), with a graphite monochromator. Corrections for absorption by crystal habit were used for **1i-l** and **6**. The structures were solved by direct methods using the SHELX-86 (**1j,k,r, 6**), SHELXS-97 (**2p**), and XS-SHELXTL (**1i,l**) programs and refined by the least-squares method in the full-matrix anisotropic (isotropic for H atoms) approximation using the SHELXL-93 and SHELXL-97 (**1j,k,l,r,2p, 6**) and XL-SHELXTL (**1i**) programs. Hydrogen atoms were located on difference Fourier maps (**1j,l,r, 2p**) or geometrically (**1i,k, 6**).

The ESR spectra were recorded on a Bruker EMX spectrometer (MW power, 0.64 mW; modulation frequency, 100 KHz; modulation amplitude, 0.1 G).

The RHF/6-31G*, MP2/6-31G*, MP2/6-31G**, and the PM3 calculations were performed using the GAMESS program [28] and the MNDO-92 program [29], respectively.

Compounds **1s** [6], **2f,m,n** [3,9], **3a–c** [12], and **7** [30] were known previously.

The syntheses described subsequently were performed in an argon atmosphere in absolute solvents with stirring. The reagents were added dropwise, and the solvents were distilled off at reduced pressure. Tables 5 and 6 list the physical and analytical data for the compounds synthesized.

TABLE 4 Crystal and Refinement Data of Compounds **1i–1r**, **2p**, and **6**^a

	1i	1j	1k	
Formula	C ₆ H ₃ BrN ₂ S ₂	C ₆ H ₃ BrN ₂ S ₂	C ₆ H ₃ BrN ₂ S ₂	
M	247.13	247.13	247.13	
Crystal system	Orthorhombic	Triclinic	Triclinic	
Space group	<i>Fdd2</i>	P-1	P-1	
<i>a</i> (Å)	13.094(2)	3.9933(5)	6.067(2)	
<i>b</i> (Å)	31.286(5)	10.903(1)	7.444(2)	
<i>c</i> (Å)	7.741(1)	18.428(2)	8.784(2)	
α (°)		97.532(7)	85.00(2)	
β (°)		91.743(9)	84.42(2)	
γ (°)		95.351(8)	86.84(2)	
<i>V</i> (Å ³)	3171.1(8)	791.21(15)	392.9(2)	
<i>Z</i>	16	4	2	
<i>D_c</i> (g cm ⁻³)	2.071	2.075	2.089	
μ (mm ⁻¹)	5.638	5.649	11.518	
<i>F</i> (000)	1920	480	240	
Crystal size (mm)	0.09 × 0.50 × 0.90	0.01 × 0.14 × 1.7	0.09 × 0.40 × 1.20	
Scan mode	$\theta-2\theta$	$\theta-2\theta$	$\theta-2\theta$	
2θ range (°)	<55	<50	<140	
Reflections measured	1103	2587	1490	
Observed $F_0 > 4\sigma_F$	1016	2075	1381	
Transmission	0.13–0.60	0.47–0.84	0.04–0.38	
<i>R</i> (observed)	0.0346	0.0340	0.0405	
<i>wR</i> ₂ (all data)	0.0957	0.0884	0.1134	
Goodness of fit	1.064	1.038	1.093	
	1l	1r	2p	6
Formula	C ₆ H ₃ BrN ₂ S ₂	C ₇ H ₆ N ₂ OS ₂	C ₁₅ H ₁₇ IN ₂ SSi	C ₁₂ H ₉ F ₂ N ₃ S ₂
M	247.13	198.26	412.36	297.34
Crystal system	Monoclinic	Monoclinic	Triclinic	Monoclinic
Space group	<i>P2₁/c</i>	<i>P2₁/n</i>	P-1	<i>P2₁/n</i>
<i>a</i> (Å)	13.6916(17)	6.052(1)	6.2520(4)	12.463(3)
<i>b</i> (Å)	3.9952(4)	10.554(2)	11.8804(8)	4.652(1)
<i>c</i> (Å)	14.393(2)	12.938(2)	12.2049(7)	23.276(6)
α (°)			96.821(5)	
β (°)	95.79(1)	97.49(2)	101.336(5)	102.86(2)
γ (°)			93.620(5)	
<i>V</i> (Å ³)	783.28(16)	819.3(2)	879.1(1)	1315.6(5)
<i>Z</i>	4	4	2	4
<i>D_c</i> (g cm ⁻³)	2.096	1.607	1.558	1.501
μ (mm ⁻¹)	5.709	5.479	2.001	3.812
<i>F</i> (000)	480	408	402	608
Crystal size (mm)	0.025 × 0.4 × 0.84	0.12 × 0.17 × 1.40	0.06 × 0.47 × 1.30	0.12 × 0.16 × 1.60
Scan mode	$\theta-2\theta$	$\theta-2\theta$	$\theta-2\theta$	
2θ range (°)	<50	<130	<50	<140
Reflections measured	1340	1390	3079	2328
Observed $F_0 > 4\sigma_F$	1101	1086	2595	1436
Transmission	0.19–0.89		0.39–0.85	0.38–0.70
<i>R</i> (observed)	0.0391	0.0476	0.0334	0.0542
<i>wR</i> ₂ (all data)	0.1045	0.1277	0.0957	0.1395
Goodness of fit	1.084	1.110	1.057	1.020

^aAtomic coordinates, thermal parameters, bond lengths, and bond angles have been deposited at the Cambridge Crystallographic Data Centre.

TABLE 5 Characterization of the Compounds

Compound	m.p. (°C), b.p. (°C/mm)	Yield (%)	Formula	MS M ⁺ (m/z) Measured (calculated, ³⁵ Cl, ⁷⁹ Br)
1i	103–105	18	C ₆ H ₃ BrN ₂ S ₂	245.8925 (245.8922)
1j	92–93	15	C ₆ H ₃ BrN ₂ S ₂	245.8925 (245.8922)
1k	118–120	3	C ₆ H ₃ BrN ₂ S ₂	245.8932 (245.8922)
1l	95–97	3	C ₆ H ₃ BrN ₂ S ₂	245.8922 (245.8922)
1n	130–130.5	1	C ₆ H ₃ IN ₂ S ₂	293.8780 (293.8784)
1o	137–138	1	C ₆ H ₃ IN ₂ S ₂	293.8786 (293.8784)
1p	124–126	8	C ₆ H ₃ IN ₂ S ₂	293.8771 (293.8784)
1q	88–89	3	C ₆ H ₃ FN ₂ S ₂	185.9720 (185.9722)
1r	109–110	18	C ₇ H ₆ N ₂ OS ₂	197.9920 (197.9922)
1t	152–153	10	C ₆ H ₂ Br ₂ N ₂ S ₂ ^a	323.8032 (323.8027)
1u	108–110	2	C ₁₂ H ₇ IN ₂ S ₂	369.9096 (369.9097)
2b	105–107/1	37	C ₁₁ H ₁₉ N ₃ SSi	253.1077 (253.1069)
2d	105–107/1	67	C ₉ H ₁₃ N ₃ O ₂ SSi	255.0494 (255.0498)
2f	144–145/1, 52–53	54	C ₉ H ₁₃ N ₃ O ₂ SSi	255.0507 (255.0498)
2g	126–128/1	62	C ₉ H ₁₃ BrN ₂ SSi	287.9723 (287.9753)
2h	124–126/1	60	C ₉ H ₁₃ BrN ₂ SSi	287.9761 (287.9753)
2i	28–29, 125–126/3	48	C ₉ H ₁₃ BrN ₂ SSi	287.9731 (287.9753)
2j	141–143/1	30	C ₉ H ₁₃ IN ₂ SSi	335.9618 (335.9615)
2k	134–135/1	65	C ₉ H ₁₃ IN ₂ SSi	335.9618 (335.9615)
2l	121–123/0.4	20	C ₉ H ₁₃ IN ₂ SSi	335.9621 (335.9615)
2o	125–126/1	60	C ₉ H ₁₂ Br ₂ N ₂ SSi	366 (366)
2p	75–76	17	C ₁₅ H ₁₇ IN ₂ SSi	411.9926 (411.9928)
3a	194–196 (dec.)	54	C ₆ H ₄ CINS ₂	153.9775 (153.9785) ^b
3b	154–158 (dec.)	74	C ₇ H ₆ CINOS ₂	183.9887 (183.9891) ^b
3c	170–174 (dec.)	79	C ₇ H ₆ CINOS ₂	183.9896 (183.9891) ^b
3d	193–197 (dec.)	60	C ₆ H ₃ BrCINS ₂	231.8874 (231.8891) ^b
3e	185–190 (dec.)	58	C ₇ H ₃ Cl ₂ NS ₂	201.9512 (201.9552) ^b
3f	180–185 (dec.)	54	C ₆ H ₃ ClIN ₂ S	279.8750 (279.8754) ^b
6	126–127		C ₁₂ H ₉ F ₂ N ₃ S ₂	297.0221 (297.0221)

TABLE 5 (Continued) Characterization of the Compounds

Compound	m.p. (°C), b.p. (°C/mm)	Yield (%)	Formula	MS M ⁺ (m/z) Measured (calculated, ³⁵ Cl, ⁷⁹ Br)
7	109–110 ^c	82	C ₁₂ H ₁₀ Cl ₂ N ₂ S ₂	315.9665 (315.9662)
10a	107–108/1	47	C ₈ H ₁₀ N ₂ OS	182.0510 (182.0514)
10b	55–56	95	C ₆ H ₃ Br ₂ NOS	294.8303 (294.8308)
10c	131–132	90	C ₁₂ H ₈ INSO	340.9372 (340.9373)
10d	120–122/4	92	C ₈ H ₁₀ N ₂ OS	182.0514 (182.0514)

^aElemental analysis, found (calculated), %: 1t: C, 22.21 (22.09); H, 0.51 (0.61); Br, 48.90 (49.08); N, 8.47 (8.59); S, 19.86 (19.63).

^b(M⁺ – Cl) peak.

^cCorresponds to Ref. [30].

1-Aryl-3-trimethylsilyl-1,3-diaza-2-thiaallenes (2b,d,f,g–k,o,p)

A solution of 0.06 mol of the corresponding ArNSO in 25 mL of hexane was added to a suspension of 10.0 g (0.06 mol) of LiN(SiMe₃)₂ in 50 mL of hexane at –30°C. During 2 hours, the temperature was raised to 20°C, and 6.6 g (0.06 mol) of Me₃SiCl in 10 mL of hexane was added. The precipitate was filtered off, the solvent was distilled off, and the residue (except for 2p) was distilled in vacuum. Compounds 2b,g–k,o were obtained as orange oils (2i solidified to orange-yellow crystals upon standing), and 2p was obtained as yellow crystals (from hexane). With Ar = 2-Me₂NC₆H₄, 4-Me₂NC₆H₄, only ArNHSiMe₃ (45 and 60%, respectively) and (ArN=)₂S were isolated. With Ar = 3-O₂NC₆H₄, in both hexane and Et₂O (see [3,9]) only the corresponding (ArN=)₂S [31] was obtained in high yield. For previously described 3-CF₃C₆H₄NSNSiMe₃, the yield in hexane was 82% compared to 30% in Et₂O [3].

To a suspension of 4.01 g (0.024 mol) of LiN(SiMe₃)₂ in 50 mL of Et₂O at –60°C, 5.73 g (0.024 mol) of *n*-O₂NC₆H₄N=SCl₂ (*n* = 2–4) [32] was added in small portions during 15 minutes. During 2 hours, the temperature was raised to 20°C, the precipitate was filtered off, and the residue distilled in vacuum. With *n* = 2, 4, compounds 2d,f were isolated as yellow oils (2f solidified upon standing). With *n* = 3, a mixture of ArNSNSiMe₃ (2e) and ArNHSiMe₃ along with (ArN=)₂S was obtained.

TABLE 6 Spectral Data of the Compounds

Compound	NMR ^a δ			UV/Vis ^b λ_{max} (nm) (log ϵ)
	¹ H	¹³ C	¹⁵ N (J, Hz)	
1i	6.85, 6.59, 5.69	137.1, 134.8, 133.3, 122.9, 119.5, 117.6	276.9 (s), 256.4 (s)	635 (2.67), 391 (3.18), 303 (4.18), 296 (4.18)
1j	6.92, 6.06, 5.66	139.4, 135.3, 125.9, 125.0, 123.4, 114.4	273.0 (s), 259.4 (d, 2.3)	633 (2.57), 377 (3.02), 290 (4.33)
1k	6.73, 5.92, 5.75	137.5, 133.0, 126.7, 125.7, 124.1, 117.2	266.2 (s), 260.3 (d, 2.4)	629 (2.52), 376 (2.92), 365 (2.93), 298 (4.06), 290 (4.66)
1l	6.78, 6.39, 5.69	139.6, 136.1, 130.9, 122.0, 119.6, 116.9	264.7 (s), 252.3 (d, 2.8)	624 (2.65), 386 (3.16), 298 (4.20)
1n	7.14, 6.24, 5.53	141.5, 139.0, 131.2, 125.2, 115.4, 94.2	272.3 (s), 259.2 (d)	629 (2.37), 379 (3.01), 289 (4.25)
1o	6.98, 6.10, 5.61	139.5, 138.3, 132.2, 124.3, 117.0, 96.7	264 ^d	626 (2.66), 402 (3.09, sh), 379 (3.23), 291 (4.20)
1p	7.02, 6.27, 5.72	142.1, 140.6, 131.6, 122.9, 120.0, 95.2	269.2 (s), 254.7 (d)	624 (2.61), 393 (3.11), 288 (3.99)
1q^c	6.25, 5.90, 5.59	165.2, 134.5, 124.6, 117.5, 115.4, 112.3	260.7 (s), 256.5 (s)	608 (2.63), 296 (4.24), 288 (4.23)
1r	6.06, 5.89, 5.41, 3.63	163.4, 131.8, 124.6, 116.5, 112.0, 111.5, 55.3	263.4 (d, 2.5), 249.1 (s)	626 (2.84), 378 (3.39), 357 (3.09), 303 (4.23)
1t	6.90, 5.77	140.7, 137.6, 125.2, 123.4, 119.7, 116.7	270 ^d , 251 ^d	632 (2.38), 390 (2.82), 303 (4.01), 294 (4.00)
1u	7.60, 7.00, 6.68, 5.86, 5.83			626 (2.77), ~380 (3.38, sh), 311 (4.35)
2b	7.07, 6.91, 6.48, 2.90, 0.21			401 (3.40, sh), 342 (3.97), 281 (3.97), 248 (4.12)
2d	7.91, 7.55, 7.36, 7.23, 0.16			344 (3.75), 266 (4.05)
2g	7.69, 7.63, 7.34, 6.88, 0.27			351 (3.47), 270 (3.53), 267 (3.56)
2h	7.80, 7.48, 7.23, 7.12, 0.22			342 (3.67), 245 (3.59)
2i	7.85, 7.34, 0.37			352 (3.94)
2j	7.84, 7.34, 7.23, 6.79, 0.16			350 (3.15), 263 (3.54), 257 (3.56)
2k	7.98, 7.51, 7.44, 6.99, 0.22			342 (3.74), 255 (3.51), 225 (3.82)
2l	7.58, 7.42, 0.24			358 (3.57), 245 (3.44)
2o	7.78, 7.51, 0.30			340 (3.78)
2p	7.84, 7.77, 7.54, 7.35, 0.34			367 (4.28), 270 (4.22)
3a^c	9.09, 9.00, 8.65, 8.46 ^f	164.0, 156.2, 139.1, 133.9, 128.1, 123.4	406 ^d	426 (3.25), 347 (4.38) ^g
3b	8.60, 8.39, 7.62, 4.42	159.8, 158.1, 157.8, 145.6, 115.7, 111.8, 58.1	378 ^d	551 (3.54), 344 (4.05)
3c	8.79, 8.29, 7.76, 4.39 ^f	173.5, 163.9, 161.8, 131.9, 130.8, 105.2, 59.7	389 ^d	439 (3.87), 368 (3.77)
3d	9.22, 8.84, 8.61	166.3, 156.7, 143.6, 131.7, 131.3, 125.0	406 ^d	
3e	8.99, 8.90, 2.95	165.5, 157.2, 154.9, 148.6, 129.8, 125.3, 21.6	400 ^d	440 (3.43, sh), 378 (4.21)
3f	9.49, 8.77, 8.72	167.2, 158.3, 149.7, 139.2, 125.9, 103.1	401.7 (d, 1.4)	
6^c	7.04–6.98, 5.79	160.6, 158.7, 142.0, 140.6, 116.0, 115.8	298 ^{d,h}	414 (4.00)

TABLE 6 (Continued) Spectral Data of the Compounds

Compound	NMR ^{a,δ}			UV/Vis ^b λ _{max} (nm) (log ε)
	¹ H	¹³ C	¹⁵ N (J, Hz)	
7	7.12–7.04, 6.55, 4.21			
10a	7.14–7.09, 6.62, 2.90			436 (3.20)
10b	7.90, 7.68			313 (3.70)
10c	7.93, 7.78, 7.59, 7.33			354 (4.32)
10d	8.35, 7.10, 6.75, 6.70, 2.87			392 (2.00)

^aIn CDCl₃; **7**, **10a**, **b**, CCl₄; **3a–d**, CF₃COOH; **2d**, neat liquid.

^bIn CHCl₃; **2b**, **d**, **g–i**, **o**, **6**, **10c**, heptane; **3a–f**, CF₃COOH.

^cδ¹⁹F: **1q**, 55.5; **6**, 45.1, 39.9.

^dδ¹⁴N

^ecf. 1,3,2-benzodithiazolium chloride (**4**) [23]: δ¹H: 9.09, 8.22; δ¹³C: 163.3, 132.6, 122.8; δ¹⁴N: 378; λ_{max} (log ε) 398 (3.38).

^fcf. Ref. [45]

^gcf. Ref. [46]

^hUnresolved asymmetric broad line.

Substituted 1,3,2,4-Benzodithiadiazines (**1i–l**, **n–r**, **t**, **u**)

Solutions of 0.01 mol of Ar–N=S=N–SiMe₃ (**2**, this work and [3]) and 0.51 g (0.5 mol) of SCl₂ (0.0075 mol in the case of **2g**, **h**, **k**), each in 30 mL of CH₂Cl₂, were mixed by adding them to 300 mL of CH₂Cl₂ at 20°C, over a period of 1 hour. After an additional 1 hour, the reaction mixture was filtered, the solvent was distilled off, and the residue was sublimed in vacuum and recrystallized from hexane. Compounds **1j**, **l**, **o**, **q**, **r**, **u** were obtained as black crystals, **1i**, **k** were obtained as shiny black-green prisms, and **1t** was obtained as lustrous dark-green crystals.

A mother liquor from recrystallization of **1j** was evaporated to dryness. The residue (which consisted of a mixture of ~60% of **1l** and ~40% of **1j** according to the ¹H NMR data) was treated with an amount of SCl₂ equimolar to **1j** under conditions described below (Reactions of compounds **1** with SCl₂). Compound **1l** was isolated as black crystals.

The compounds **1n**, **p** (38%) were obtained by sublimation of reaction residue as a 3:2 mixture (¹H NMR). Fractional crystallization from hexane afforded pure **1n** as black needles. A sample of pure **1p** was prepared by reaction of the **1n**, **p** mixture with SCl₂ (see below Reactions of compounds **1** with SCl₂).

1,3,2,4-Benzodithiadiazine (**1s**)

Under the conditions described previously, compound **1s** [6] was obtained from **2j** in 9% yield.

Reactions of Compounds **1** with SCl₂. 1,2,3-Benzodithiazolium Chloride (**3a**) and Its 4-Methoxy- (**3b**), 6-Methoxy- (**3c**), 5-Bromo- (**3d**), 5-Methyl-6-chloro- (**3e**), and 5-Iodo (**3f**) Derivatives, and ¹⁴N NMR Detection of Thiacyl Chloride NSCl

At 20°C, a solution of 0.003 mol of the corresponding **1** (this work and [3]) in 10 mL of CH₂Cl₂ was added during 45 minutes to a solution of 0.31 g (0.003 mol) of SCl₂ in 10 mL of the same solvent. After 15 minutes, the precipitate was filtered off, dried in vacuum, and, in the case of **3a–c**, recrystallized from SOCl₂:CCl₄ (3:1). Salts **3a–e** [12] were obtained as crystalline solids (in the case of **3e**, **d** chlorination of a carbocycle occurred during reaction or recrystallization, respectively).

To detect NSCl, the interaction was performed in CCl₄ with more concentrated (~0.6 M) solutions, and the ¹⁴N NMR spectrum of the reaction mixture was measured periodically during the reaction. The ¹⁴N NMR identification of NSCl (δ¹⁴N: 730) was based on previously reported data [33] and on control measurements with an authentic sample prepared [33] by heating a solution of (NSCl)₃ [34] in CCl₄. At the same time, (NSCl)₃ [33,35] was not found in the reaction mixtures in accordance with the conclusion that trimerization of NSCl is very kinetically hindered [33].

At 20°C, to a solution of 0.247 g (0.001 mol) of a 1:1 mixture of 6-Br (**1j**) and 8-Br (**1l**) derivatives of **1** in 5 mL of CH₂Cl₂ was added during 15 minutes to a solution of 0.051 g (0.0005 mol) of SCl₂ in 5 mL of

the same solvent. The precipitate was filtered off. Salt **3d** (0.081 g; 60%) was obtained as a yellow powder. Its purity was checked by ^1H NMR spectroscopy. No traces of the 7-Br isomer have been found. The filtered solution was evaporated to dryness, and the residue was sublimed in vacuum. A 10:1 (^1H NMR) mixture (0.121 g) of **1i** and **1j** was obtained. Recrystallization from hexane gave pure **1i** (0.060 g).

In a similar way, **3f** (light brown powder) and **1p** (black crystals) were prepared from the 3:2 mixture of **1n** and **1p**.

Reaction of 5,6,7,8-tetrafluoro-1,3,2,4-benzodithiadiazine [15] with SCl_2 provided only unidentified products without any evidence of the formation of the polyfluorinated analog of **3a**.

Reaction of 1,3,2,4-Benzodithiadiazine (**1s**) with $(\text{NSCl})_3$ and Cl_2

At 20°C , to a solution of 0.17 g (0.0007 mol) of $(\text{NSCl})_3$ [34] in 2 mL of CCl_4 was added a solution of 0.34 g (0.002 mol) of **1s** [6] in 2 mL of the same solvent. The crude salt **3a** (0.38 g; 80%) was obtained as a brown precipitate.

At 20°C , an excess of Cl_2 was passed through a solution of 0.34 g (0.002 mol) of **1s** in 10 mL of CCl_4 . A slightly exothermic fast reaction afforded a complex mixture (^1H NMR, MS) of unidentified products.

Attempted Preparation of **1** from 2-Aminobenzenethiols

*Reaction of 5-Chloro-2-aminobenzenethiol (**9**) with $(\text{SN})_4$* A mixture of 0.80 g (0.005 mol) of **9** [30], 0.46 g (0.0025 mol) of $(\text{SN})_4$ (see below), and 50 mL of MeCN was boiled until NH_3 evolution ceased (~ 1 hour). The mixture was filtered, the solvent was distilled off, and the residue was recrystallized from toluene. Compound **7** was obtained as yellow crystals.

Attempted Preparation of **1** from Herz Salts

*Reaction of 1,2,3-Benzodithiazolium Chloride (**3a**) with Trimethylsilyl Azide, and ESR Detection of 1,2,3-benzodithiazolyl (**8**)* A mixture of 0.115 mg (10^{-6} mol) of Me_3SiN_3 [36], 0.190 mg (10^{-6} mol) of **3a**, and 1 mL of hexane, outgassed by three freeze-pump-thaw cycles, was heated in an ESR valve-equipped quartz capillary tube to 50°C . The ESR spectrum of the solution was identical to that of **8** [1]. Experiments on a preparative scale (in both hexane and MeCN) were accompanied by gas evolution and, after usual work-up, gave unidentified red tar. Similar results were obtained with salts **3b,c**.

Cyclotetra(azathiene) ($(\text{SN})_4$)

At 20°C , solutions of 4.12 g (0.02 mol) of $(\text{Me}_3\text{SiN}=\text{S})_2$ [37] and 2.06 g (0.02 mol) of SCl_2 , each in 30 mL of CH_2Cl_2 , were mixed by adding them to 100 mL of CH_2Cl_2 over a period of 1 hour. After an additional 2 hours, the mixture was filtered, the solvent was distilled off, and the residue was recrystallized from toluene. $(\text{SN})_4$ [38] (1.57 g; 85%) was obtained as orange-yellow needles, m.p. (decomposition) $185\text{--}186^\circ\text{C}$ (cf. [38]).

*Reaction of 1-(4-fluorophenyl)-3-trimethylsilyl-1,3-diaza-2-thiaallene (**2m**) with 1,2,3-benzodithiazolium chloride (**3a**)*

At 20°C , to a suspension of 0.40 g (0.0021 mol) of **3a** (this work and Ref. [12]) in 5 mL of CH_2Cl_2 was added, during 20 minutes, a solution of 0.48 g (0.0021 mol) of **2m** [3] in 5 mL of the same solvent. The mixture was stirred overnight, the solution was filtered off, the solvent was distilled off, and the residue was recrystallized twice from hexane. Compound **6** was obtained in 0.01 g yield as small orange needles.

N-Sulfinylanilines. $\text{RC}_6\text{H}_4\text{NSO}$ (**10**) was prepared by the Michaelis reaction [39] from the corresponding amines and SOCl_2 [40] except for the Me_2N -substituted derivatives synthesized from ArNH_2 and PhSO_2NSO [40] or ArNH_2 and dipyrid-2-yl sulfite [47]. Most compounds were known earlier [40], except for those in which $\text{R} = 3\text{-Me}_2\text{N}$ (**10a**, dark red oil), $3,5\text{-Br}_2$ (**10b**, pale yellow crystals), $4\text{-}(4'\text{-IC}_6\text{H}_4)$ (**10c**, bright yellow crystals) and $2\text{-Me}_2\text{N}$ (**10d**, dark red oil) derivatives.

4-Iodo-4'-aminobiphenyl [41]. 70%, Colorless plates, m.p. $165\text{--}165.5^\circ\text{C}$ (hexane/toluene) for preparation of **10c** was obtained by SnCl_2/HCl reduction of 4-iodo-4'-nitrobiphenyl synthesized in one step from biphenyl under the Tronov–Novikov conditions [42] by a known procedure [43] [37%, colorless crystals, m.p. $217\text{--}218^\circ\text{C}$ (toluene) (212°C [43])]. Both compounds were purified by column chromatography on silica followed by recrystallization. Warning: the compounds needed to be treated with caution because they are potential carcinogens [44].

ACKNOWLEDGMENTS

The authors are grateful to Mr. Ivan V. Vlasyuk for ESR measurements and Mr. Yuri V. Pavlov for assistance in some preparations.

REFERENCES

- [1] Bagryansky, V. A.; Vlasyuk, I. V.; Gatilov, Yu. V.; Makarov, A. Yu.; Molin, Yu. N.; Shcherbukhin, V. V.; Zibarev, A. V. *Mendeleev Commun* 2000, 5–7.
- [2] Manaev, A. V.; Makarov, A. Yu.; Gatilov, Yu. V.; Latosinska, J. N.; Shcherbukhin, V. V.; Traven, V. F.; Zibarev, A. V. *J Electron Spectrosc Relat Phenom* 2000, 107, 33–38.
- [3] Bagryanskaya, I. Yu.; Gatilov, Yu. V.; Makarov, A. Yu.; Maksimov, A. M.; Miller, A. O.; Shakirov, M. M.; Zibarev, A. V. *Heteroat Chem* 1999, 10, 113–124.
- [4] Planar or nearly planar molecular geometry in the crystal, united molecular 12ρ -electron system with markedly low first ionization energy (to 7.75 eV), and low-lying excited states ($\lambda_{\max} \sim 625$ nm) [2,3]; for introduction into the problem, see: Minkin, V. I.; Simkin, B. Ya.; Glukhovzev, M. N. *Aromaticity and Anti-aromaticity: Electronic and Structural Aspects*; Wiley Interscience: New York, 1994.
- [5] Zibarev, A. V.; Gatilov, Yu. V.; Bagryanskaya, I. Yu.; Maksimov, A. M.; Miller, A. O. *J Chem Soc Chem Commun* 1993, 298–299.
- [6] Cordes, A. W.; Hojo, M.; Koenig, H.; Noble, M. C.; Oakley, R. T.; Pennington, W. T. *Inorg Chem* 1986, 25, 1137–1145.
- [7] (a) Barclay, T. M.; Beer, L.; Cordes, A. W.; Oakley, R. T.; Preuss, K. E.; Taylor, N. J.; Reed, R. W. *J Chem Soc Chem Commun* 1999, 531–532; (b) Torroba, T. J. *Prakt Chem* 1999, 341, 99–113.
- [8] One intention is to adopt substitution reactions of iodoarenes to further modification of the title compounds under mild conditions including their coupling (on an attempt to prepare benzo(bis)dithiadiazine), see: Codding, P. W.; Koenig, H.; Oakley, R. T. *Can J Chem* 1983, 61, 1561–1566.
- [9] Zibarev, A. V.; Miller, A. O.; Gatilov, Yu. V.; Furin, G. G. *Heteroat Chem* 1990, 1, 443–453.
- [10] Herberhold, M.; Gerstmann, S.; Milius, W.; Wrackmeyer, B.; Borrmann, H. *Phosphorus Sulfur Silicon* 1996, 112, 261–279.
- [11] Allen, F. H.; Kennard, O.; Watson, D. G.; Brammer, L.; Orpen, A. G.; Taylor, R. *J Chem Soc Perkin Trans 2* 1987, S1–S19.
- [12] (a) Kirsch, G. In *Methoden der Organischen Chemie* (Houben-Weyl), Bd. 8Ed.; Schaumann, E., Ed.; Thieme: Stuttgart, 1994, 3–12; (b) Sammes, M. P. In *Comprehensive Heterocyclic Chemistry*; Katritzky, A. R., Rees, C. W., Eds.; Pergamon Press: Oxford, 1984, Vol. 6, pp 897–946.
- [13] In contrast to extremely labile neutral sulfur iodides, sulfur iodine cations are relatively stable and structurally characterized: Klapoetke, T.; Passmore, J. *Acc Chem Res* 1989, 22, 234–240.
- [14] It follows from the data of Penning ionization electron spectroscopy that HOMO of C_6H_5I featured high reactivity upon electrophilic attack due to a great contribution from the halogen; it is not the case for other C_6H_5Hal (Hal = F, Cl, Br): Fujisawa, S.; Ohno, K.; Masuda, S.; Harada, Y. *J Am Chem Soc* 1986, 108, 6505–6511.
- [15] Zibarev, A. V.; Gatilov, Yu. V.; Miller, A. O. *Polyhedron* 1992, 11, 1137–1141.
- [16] (a) Burdett, J. K. *Molecular Shapes*; Wiley-Interscience: New York, 1990; (b) Schastnev, P. V.; Shchegoleva, L. N. *Molecular Distortions in Ionic and Excited States*; CRC Press: Boca Raton, FL, 1995.
- [17] (a) The detailed data of stationary and laser flash photolysis, matrix isolation study, and nonempirical quantum chemical calculations to support the proposed participation of **5** in the reactions of the title compounds, including formation of 1,2,3-benzodithiazolyls, will be presented elsewhere. For preliminary communication, see: Gritsan, N. P.; Bagryansky, V. A.; Vlasyuk, I. V.; Molin, Yu. N.; Makarov, Yu. A.; Zibarev, A. V. *Abstracts of Papers, Conference on Reactive Intermediates and Unusual Molecules*, Vienna, Austria, 2000; p 64; (b) The formation of 1,2,3-benzodithiazolium salt is suppressed on going from **1s** to its tetrafluoro derivative, probably owing to the impossibility of positively charged SSN moiety stabilization at the electron-withdrawing tetrafluorobenzene ring.
- [18] Markovsky, L. N.; Fedyuk, G. S.; Levchenko, E. S.; Kirsanov, A. V. *Zh Org Khim* 1973, 9, 2502–2506 (in Russian); *Chem Abstr* 1974, 80, 82816.
- [19] Bats, J. W.; Fuess, H.; Weber, K. L.; Roesky, H. W. *Chem Ber* 1983, 116, 1751–1755.
- [20] Zibarev, A. V.; Beregovaya, I. V. *Rev Heteroat Chem* 1992, 7, 171–190.
- [21] (a) At the PM3 level of theory, the cation of **3a** (ΔH_f° 249.1 kcal mol⁻¹) is less stable than the isomeric cation of **4** (Δ_f° 233.5 kcal mol⁻¹). The S–S bond length is very long in the Herz salts (2.023 Å in the experimental geometry of 4,6-di(*tert*-butyl) derivative of **3a** [19]), probably due to Coulombic repulsion of positively charged S atoms ($3a, q: S^1, 0.36; S^2, 0.45$), being comparable with the average S–S ordinary bond length in organic disulfides (2.048 Å [11]). The S–S bond order in the cation of **3a** is only 1.04. The absolute hardness, η , which can serve as a unifying measure of aromaticity [22], is equal to 3.205 and 3.471 eV for the cations of **3a** and **4**, respectively. One can speculate that in the case of salts **3** the 10- ρ -electron system of a cation is very close to the situation with a topological cleavage at the S–S bond; (b) It follows from E_{tot} values (PM3) that N_3^- is more stable than N_3^\bullet by 3.1 eV (IE), whereas cation of **3a** is less stable than the corresponding **8** radical by 8.0 eV (EA).
- [22] (a) Zhou, Z.; Navangul, H. V. *J Phys Org Chem* 1990, 3, 784–788; (b) Pearson, R. G. *Chemical Hardness*; Wiley-VCH: New York, 1997.
- [23] (a) Zibarev, A. V.; Bagryanskaya, I. Yu.; Gatilov, Yu. V.; Shchegoleva, L. N.; Dolenko, G. N.; Furin, G. G. *Khim Geterotsikl Soedin* 1990, 1683–1688 (in Russian); *Chem Abstr* 1991, 115, 7769; (b) Zibarev, A. V.; Rogoza, A. V.; Konchenko, S. N.; Fedotov, M. A.; Furin, G. G. *Zh Obshch Khim* 1991, 61, 441–445 (in Russian); *Chem Abstr* 1991, 115, 92167; (c) Awere, E. G.; Burford, N.; Haddon, R. C.; Parsons, S.; Passmore, J.; Waszczak, J. V.; White, P. S. *Inorg Chem* 1990, 29, 4821–4830.
- [24] (a) Bagryanskaya, I. Yu.; Gatilov, Yu. V.; Shakirov, M. M.; Zibarev, A. V. *Zh Strukt Khim* 1996, 37, 363–367 (in Russian); *Chem Abstr* 1997, 126, 18934; (b) Bagryanskaya, I. Yu.; Gatilov, Yu. V.; Shakirov, M. M.; Zibarev, A. V. *J Struct Chem* 1996, 37, 318–322.
- [25] (a) Schuster, G. B.; Platz, M. S. *Adv Photochem* 1992, 17, 69–143; (b) Gritsan, N. P.; Pritchina, E. A. *Usp*

- Khim 1992, 61, 910–939 (in Russian); Chem Abstr 1992, 117, 211713.
- [26] Atkinson, R. S.; Judkins, B. D.; Khan, N. J Chem Soc Perkin Trans 1 1982, 2491–2491.
- [27] Mayer, R.; Decker, D.; Bleisch, S.; Domschke, G. J Prakt Chem 1987, 329, 81–86.
- [28] Schmidt, M. V.; Baldrige, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. J.; Koseki, S.; Matsunaga, N.; Nguen, K. A.; Su, S.; Windus, T. L.; Dupuis, M.; Montgomery, J. A. J Comput Chem 1993, 14, 1347–1363.
- [29] Specialized fund of quantum chemical programs of the Siberian Division of the Russian Academy of Sciences, Institute of Chemical Kinetics and Combustion, Novosibirsk, Russia.
- [30] Palmer, P. J.; Trigg, R. B.; Warrington, J. K. J Med Chem 1971, 14, 248–251.
- [31] Hoerhold, H.H.; Beck, J. J Prakt Chem 1969, 311, 621–629.
- [32] (a) Levchenko, E. S.; Sheinkman, I. E. Zh Obshch Khim 1966, 36, 428–432 (in Russian) Chem Abstr 1996, 65, 634; (b) Levchenko, E. S.; Sheinkman, I. E. J Gen Chem USSR 1966, 36, 446–450.
- [33] Passmore, J.; Schriver, M. Inorg Chem 1988, 27, 2749–2751.
- [34] Alange, G. G.; Banister, A. J.; Bell, B. J Chem Soc Dalton Trans 1972, 2399–2340.
- [35] Chivers, T.; Oakley, R. T.; Scherer, O. J.; Wolmershaeuser, G. Inorg Chem 1981, 20, 914–917.
- [36] Birkofer, L.; Ritter, A.; Richter, P. Chem Ber 1963, 96, 2750–2757.
- [37] Bagryanskaya, I. Yu.; Gatilov, Yu. V.; Miller, A. O.; Shakirov, M. M.; Zibarev, A. V. Heteroat Chem 1994, 5, 561–565.
- [38] (a) Becke-Goehring, M. Inorg Synth 1960, 6, 123–128; (b) Heal, H. G. Adv Inorg Chem Radiochem 1972, 15, 375–412.
- [39] Obviously, this reaction of general significance (see, for example: Bussas, R.; Kresze, G.; Muensterer, H.; Schwoebel, A., Sulfur Reports 1986, 2, 215–378) is worthy of naming. Surprisingly, it is not made as yet (see, for example Schneller, S. Int J Sulfur Chem B 1972, 7, 155–171; Schneller, S. Int J Sulfur Chem 1976, 8, 579–597; For scientific biography of Prof. August Michaelis (1847–1916), one of the pioneers of heteroatom chemistry, see Teller, J.; Teller, M. Beitrage zur Geschichte der Universitaet Rostock, 1984, 27–41.
- [40] Gmelin Handbook of Inorganic Chemistry; Sulfur-Nitrogen Compounds, Part 6; Springer: Berlin, 1990; pp 113–171.
- [41] Belcher, R.; Nutten, A. J.; Stephen, W. I. J Chem Soc 1953, 1334–1337.
- [42] Merkushev, E. B. Synthesis 1988, 923–937.
- [43] Sedov, A. M.; Sergeeva, A. A.; Novikov, A. N. Izv Vyssh Ucheb Zaved Khim Khim Technol 1970, 13, 591–592 (in Russian); Chem Abstr 1970, 73, 87559.
- [44] You, Z.; Brezzell, M. D.; Das, S. K.; Hooberman, B. H.; Sinsheimer, J. E. Mutat Res 1994, 320, 45–58.
- [45] Akulin, Yu. I.; Gel'mont, M. M.; Strelets, B. Kh.; Efros, L. S. Khim Geterotsikl Soedin 1978, 912–916 (in Russian); Chem Abstr 1979, 90, 54275.
- [46] Strelets, B. Kh.; Efros, L. S. Zh Org Khim 1969, 5, 153–158 (in Russian); Chem Abstr 1969, 70, 86889.
- [47] Kim, S.; Yi, K. Y. Tetrahedron Lett 1986, 27, 1925–1928.