

TOTAL SYNTHESIS OF (+)-ANGUSTOLINE

Ichiya Ninomiya\* and Takeaki Naito


Kobe Women's College of Pharmacy,  
Motoyamakita, Higashinada, Kobe, Japan.

(+)-Angustoline, an alkaloid of Strychnos angustiflora, was synthesized via the route including enamide photocyclization.

As an extension of synthetic work on angustidine,<sup>1</sup> we report the first total synthesis of (+)-angustoline, a major alkaloid of Strychnos angustiflora,<sup>2</sup> thus completing formal synthesis of all three alkaloids so far isolated.

Treatment of sodium 5-acetylnicotinate<sup>3</sup> with oxalyl chloride produced the acid achloride, which without purification was reacted with harmalane to afford the enamide (Ia) as a yellow solid,  $\nu_{\max}^{\text{cm}^{-1}}$  (CHCl<sub>3</sub>) 3520 (NH), 1695 (Ac), and 1640 (NCO). A 0.02 M methanolic solution of (Ia) was irradiated with a low pressure mercury lamp at room temperature for 5 h. Chromatography of the reaction mixture on alumina separated two products (IIa and IIIa) in 30 and 8 % yields respectively; (IIa), m.p. > 300°, M<sup>+</sup> 329; (IIIa), m.p. > 300°, M<sup>+</sup> 329.

Sodium borohydride reduction of (IIa) afforded the alcohol (IIb), m.p. 292-294° (lit. 310-314°)<sup>2</sup>, M<sup>+</sup> 331, which was shown to be (+)-angustoline by comparisons of i.r., n.m.r., and mass spectra. Since angustoline had been converted into angustine and dihydroangustine,<sup>4</sup> this synthesis formally completed the total synthesis of all three alkaloids so far isolated.<sup>2</sup>



In relation to the above synthesis, photocyclization of the enamide (Ib), which was obtained from harmalane and 5-(1-chloroethyl)nicotinoyl chloride, was carried out and afforded three photocyclized products (IIc, IIIb, and IIIc), which were separated by chromatography on alumina; (IIc), m.p. 220-223°,  $M^+$  345; (IIIb), m.p.  $>300^\circ$ ,  $M^+$  315; (IIIc), m.p. 266-269°,  $M^+$  345. The compound (IIc) should be designated as (+)-angustoline methyl ether. The formation of these three photoproducts would seem to be formed either by hydrogenolysis or substitution of the chloride by solvent during the course of irradiation.

ACKNOWLEDGEMENT We thank Dr. H.T. Cheung of the University of Sydney for i.r., n.m.r., and mass spectra of natural angustoline.

## REFERENCES

- 1 I. Ninomiya, H. Takasugi, and T. Naito, J. Chem. Soc. Chem. Comm., 1973, 732.
- 2 a) T. Y. Au, H. T. Cheung, and S. Sternhell, J. Chem. Soc. Perkin I, 1973, 13; b) J. D. Phillipson, S. R. Hemingway, N. G. Bisset, P. J. Houghton, and E. J. Shellard, Phytochemistry, 1974, 13, 973.
- 3 R. Lukes and P. Vaculik, Chem. Listy, 1957, 51, 1510.
- 4 Synthesized independently by two groups; R. T. Brown, A. A. Charalambides, and H. T. Cheung, Tetrahedron Letters, 1973, 4837; A. Shafiee and E. Winterfeldt, Chem. Ber., 1974, 107, 966; Synthesis, 1974, 185.

Received, 13th August, 1974