

THE FRIEDEL-CRAFTS REACTION OF 1-METHYL-2(1H)-PYRIDONE AND ITS DERIVATIVES

Hiroshi Tomisawa, Reiko Fujita, Hiroshi Hongo, Hideki Kato, and Masao Watanabe
Tohoku College of Pharmacy
Komatsushima, Sendai 983

The Friedel-Crafts reaction of 1-methyl-2(1H)-pyridone (I), 1-methyl-2(1H)-quinolone (II), and 2-methyl-1(2H)-isoquinolone (III) was carried out, and the main results are shown in the following table. The structures of the products were confirmed by spectral or chemical method.

Compound No.	Reagent	Catalyst	Temp. (°C)	Period (days)	Product (Yield %)
I ¹⁾	(MeCO) ₂ O	H ₂ SO ₄	140	4	IV(34.3)
I ¹⁾	(EtCO) ₂ O	H ₂ SO ₄	180	2	IV(66.4), V(2.5)
I ¹⁾	(C ₃ H ₇ CO) ₂ O	H ₂ SO ₄	180	2	IV(75.9), V(7.4)
I ²⁾	(C ₆ H ₅ CO) ₂ O	H ₂ SO ₄	180	5	IV(38.2), V(36.7), VI(2.7)
I ³⁾	C ₆ H ₅ COCl	AlCl ₃	180	2	IV(30.9), V(3.8), VI(12.4)
II ⁴⁾	MeCOCl	AlCl ₃	50	14	VII(37.5)
II ⁴⁾	EtCOCl	AlCl ₃	80	14	VII(6.5), VIII(25.8)
II ²⁾	(EtCO) ₂ O	H ₂ SO ₄	180	3	VII(16.1)
II ³⁾	C ₆ H ₅ COCl	AlCl ₃	120	2	VII(28.3), VIII(2.3)
III ²⁾	(MeCO) ₂ O	H ₂ SO ₄	140	1/24	IX(75.6), (72) ⁶⁾
III ⁵⁾	MeCOCl	AlCl ₃	50	2	IX(40.4), X(6.9)
III ²⁾	(EtCO) ₂ O	AlCl ₃	180	3	IX(58)
III ²⁾	(C ₆ H ₅ CO) ₂ O	H ₂ SO ₄	120	5	IX(96.4)
III ³⁾	C ₆ H ₅ COCl	AlCl ₃	120	7	IX(21.6), XI(19.5)

IV: 5-acylated I, V: 3,5-diacylated I, VI: 3-acylated I, VII: 6-acylated II, VIII: 3-acylated II, IX: 4-acylated III, X: 7-acylated III, XI: 5-acylated III

References

- 1) R. Fujita, Chem. Pharm. Bull. (Tokyo), submitted. 2) H. Tomisawa, R. Fujita, H. Hongo, and H. Kato, ibid., submitted. 3) H. Tomisawa, R. Fujita, H. Hongo, and H. Kato, ibid., in the press. 4) H. Tomisawa, M. Watanabe, R. Fujita, and H. Hongo, ibid., 18, 917 (1970). 5) H. Tomisawa and R. Fujita, ibid., 21, 2585 (1973).
- 6) D. E. Horning, G. Lacasse, and J. M. Muchowski, Can. J. Chem. 49, 2785 (1971).