

SYNTHESES OF DIBENZO(b,f)AZONINE AND DIBENZO(b,g)AZECINES
THROUGH RING TRANSFORMATION

Shinzo Kano, Toshihisa Ogawa, Eiji Komiyama, Tsutomu Yokomatsu,

Yoko Takahagi, and Shiroshi Shibuya

Tokyo Collge of Pharmacy

3-20-1, Kitashinjuku, Shinjuku-ku, Tokyo 160

The benzyne reaction of 1-(2-bromo-4,5-dimethoxyphenethyl)-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline(21) using sodium dimsyl afforded 5,6,7,12,13,14-hexahydro-10-hydroxy-7-methansulfinylmethyl-2,3,9-trimethoxy-dibenzo[b,g]azecine(22). This product was previously assigned to 5,6,7,8,9,10-hexahydro-3-hydroxy-8-methansulfinylmethyl-2,12,13-trimethoxy-7-methyldibenzo[e,g]azecine(25a), which was revised to (22). Reductive deoxygenation of (22) with Zn-Hg gave 7-(methylthio)methyl derivative(23) whose desulfurization yielded 7-methyldibenzo[b,g]azecine(24). The similar reaction using 1-(2-bromo-4,5-methylenedioxyphenethyl)- and 1-(3-bromo-4-methoxyphenethyl)-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline also examined to give the corresponding dibenzo[b,g]azecines(28) and (29), respectively. These 7-methansulfinylmethyl-dibenzo[b,g]azecines were converted to 7-methyldibenzo[b,g]azecines through deoxygenation of (28) and (29), followed by desulfurization of 7-(methylthio)methyl derivatives (30) and (31). The structure of (30) was confirmed by comparison of the spectroscopic data with those of the authentic specimen alternatively synthesized.

This ring transformation reaction was applied to the formation of 6-substituted dibenzo[b,g]azonines from the 1-halogenobenzyl-6-hydroxyisoquinolines.