


INDOLE-SINGLET OXYGEN REACTIONS. A NOVEL REARRANGEMENT OF
THE PEROXIDIC INTERMEDIATES TO 2,3-DIHYDRO-1,4-BENZOXAZINES

Isao Saito, Mitsuru Imuta, Seichi Matsugo, and Teruo Matsuura

Department of Synthetic Chemistry, Faculty of Engineering, Kyoto University

Yoshida, Kyoto 606

Dye-sensitized photooxygenation of various N-methylindoles has been investigated. In most cases, N-methylindoles gave the corresponding C₂-C₃ ring cleavage products which are considered to be formed through a dioxetane or an ionic peroxide such as a zwitterionic peroxide or a perepoxyde. In an attempt to intercept the peroxidic intermediates by the functional group of the side chain, we have carried out the photooxygenation of N-methyltryptophol(1). Photooxygenation of 1 at room temperature produced the C₂-C₃ ring cleavage product, whereas at low temperature 1 gave the 3-hydroperoxyindoline 2 in 95% yield. In addition, the hydroperoxidic product 2 readily underwent a new type of rearrangement to give the corresponding 2,3-dihydro-1,4-benzoxazine 3 in the presence of catalytic amounts of acid.



Under similar conditions, photooxygenation of N-methylindole-3-propionic acid, tryptamine hydrochloride, and tryptophan methyl ester hydrochloride in methanol gave the corresponding 2,3-dihydro-1,4-benzoxazine in excellent yield. Thus the results provide a simple and general method for the oxidative transformation of indoles into 1,4-benzoxazine systems. The results also offer a suggestive information that such transformations may potentially be involved in the oxidation of indoles in biological systems. The participation of singlet oxygen in these reactions has also been demonstrated.