

SYNTHESIS AND PROPERTIES OF STABLE QUINONE METHIDES IMINES,  
4-(1,3-DITHIOL-2-YLIDENE)-2,5-CYCLOHEXADIEN-1-IMINES

Jyuzo Nakayama, Naohiko Matsumaru, and Masamatsu Hoshino  
Department of Chemistry, Saitama University, Urawa, Saitama 338

Toshiro Takemasa

College of Liberal Arts, Saitama University, Urawa, Saitama 338

Synthesis and spectroscopic properties of stable quinone methide imines were reported.

The reaction of 2,6-diethylaniline with 2 molar amounts of 1,3-benzodithiolylium tetrafluoroborate in acetonitrile at room temperature gave 2-(4-amino-3,5-diethylphenyl)-1,3-benzodithiolylium tetrafluoroborate (1a) as deep green crystals in 75% yield. Treatment of 1a with 1,8-diazabicyclo[5.4.0]-7-undecene (DBU) in acetonitrile afforded the stable quinone methide imine, 2,6-diethyl-4-(1,3-benzodithiol-2-ylidene)-2,5-cyclohexadien-1-imine (2a), as orange crystals in 61% yield.

2-Methylthio-1,3-dithiolylium iodide reacted with 2,6-diethylaniline in boiling acetonitrile, giving rise to 2-(4-amino-3,5-diethylphenyl)-1,3-dithiolylium iodide (1b) in 51% yield. Treating 1b with DBU in acetonitrile gave the quinone methide imine, 2,6-diethyl-4-(1,3-dithiol-2-ylidene)-2,5-cyclohexadien-1-imine (2b) in 72% yield.

A variety of stable quinone methide imines were prepared in similar ways.

