


PO 34

TAUTOMERIC FORMS OF THE BENZOTHIAZOLYL DERIVATIVES OF β -DICARBONYL COMPOUNDS


Halgaš J., Regec V., Zahradník P.

Department of Organic Chemistry, Faculty of Science, Comenius University, 816 00 Bratislava

2-Benzothiazolyl derivatives of β -dicarbonyl compounds may exist in several tautomeric forms. Recently, on the basis of nmr spectra, the diketo form I was discarded and the thiazoline form II was suggested for ethyl malonate, ethyl acetoacetate, ethyl cyanoacetate and malononitrile derivatives.



We have prepared the compounds quoted (extended by acetylacetone derivative) by the reaction of 2-fluorobenzothiazole with sodium salts of appropriate C-acids in ether or THF. For ethyl acetoacetate and acetylacetone derivatives we have also considered the enol form III ($Z = \text{CH}_3, \text{OC}_2\text{H}_5$). As comparative samples we have synthesised N-methyl derivatives IV and complexes V:




Spectroscopic and quantum chemical investigations has been performed to confirm or deny the thiazoline and enol forms.

PO 35


YLIDES OF MALONYL HETEROCYCLIC COMPOUNDS

Thomas Kappe*, Gertraud Korbuly and Wolfgang Stadlbauer
Institute of Organic Chemistry, University of Graz, Austria

During the course of work on sixmembered mesoionic compounds of type 1, containing the negatively charged malonyl residue, we became interested in ylides of malonyl heterocyclic compounds (2) in which the positive charge is located outside of the heterocyclic ring system. For a comparative study a number of synthetic routes for the preparation of pyridinium-, iodonium-, sulfonium-, and phosphonium-ylides (2) have been developed.



We have already described the synthesis of pyridinium-ylides from dichloro-1) and monochloro-malonyl derivatives²⁾. We have now found, that iodonium-ylides, which are readily obtained from malonylheterocycles and aryl-iodonium diacetates, are converted to pyridinium ylides with pyridine in the presence of acids. Without acidic catalysis rearrangement to arylethers takes place. These reactions are exemplified with coumarin and carbostyryl derivatives according to the following reaction scheme:



The iodonium-ylides 3 undergo a thermal rearrangement yielding the 3-iodo-4-arylethers 4, which can be converted by photocyclization to benzofuran derivatives (f. i. 4, $X = \text{O}$, — „coumeston“). However, with pyridine in acetic acid the pyridinium-ylides 5 are obtained. The reaction of isoatoic anhydride (6) with the pyridinium betaine 7 yields the ylide 5 ($X = \text{NH}$), too.

The corresponding phosphonium- and sulfonium-ylides can be prepared by exchange reactions starting with iodonium-ylides, however, the latter are also obtained from the unsubstituted malonyl heterocycles and sulfoxides with acetic anhydride as dehydrating agent.

REFERENCES

- 1) T. Kappe, R. Khorchid-Zadeh, H. Steininger, Z. Naturforsch. 30b, 773 (1975).
- 2) G. Schindler, T. Kappe, Synthesis 1977, 243.

PO 36

CONFORMATIONAL ANALYSIS OF MORPHINE-TYPE OPIATE AGONISTS AND ANTAGONISTS. RELATIONSHIP BETWEEN THE CONFORMATION OF THE PIPERIDINE RING IN THESE OPIATES AND THE TYPE OF THEIR ACTIVITY (AGONIST OR ANTAGONIST)

Vera M. Kolb

Department of Chemistry and Biochemistry, and Medical School, Southern Illinois University, Carbondale, Illinois, 62901, USA

Qualitative conformational analysis of naloxone (I, an opiate antagonist) shows that the overall stability of naloxone is greater when the piperidine ring is in the skew boat conformation (N-allyl pseudo-equatorial) rather than the chair conformation (N-allyl equatorial) since in the latter conformation there are unfavorable interactions between the N-allyl group and the ring B. The most stable piperidine conformation of oxymorphone (II, an opiate agonist) is, however, a chair form (N-methyl equatorial). The results of this conformational analysis revealed a new aspect of the structure-activity relationship of morphine-type agonists and antagonists and led to a proposal of a new opiate-receptor model (V. M. Kolb, J. Pharm. Sci., in press). The salient features of this new opiate-receptor model are: (1) Only one receptor conformation is necessary for the binding of opiate agonists and antagonists; (2) The receptor possesses two amine-binding sites (ABS) - agonist and antagonist; (3) Agonists and antagonists interact via their nitrogen lone-pair electrons with the respective ABS; (4) In naloxone (I), for example, the antagonist effect is derived indirectly through the N-allyl group which sterically forces the N-lone-pair lobe in the direction of the antagonist ABS. According to this model, the respective conformations of naloxone, an antagonist, and oxymorphone (II), an agonist, required for binding to the receptor, are illustrated: