


Photoaddition of Vinyl Ethers to N-Methylphthalimide. Competing 2 + 2 Processes.

Paul H. Mazzocchi\*, Saeko Minamikawa and Michael Bowen  
Department of Chemistry, University of Maryland, College  
Park, Md., 20742, USA

The photochemical addition of vinyl alkyl ethers to N-methylphthalimide to give 3,4-benzo-6,7-dihydro-1-methylazepine-2,5-diones in a  $\pi^2 + \sigma^2$  process is found to compete with the  $\pi^2 + \pi^2$  Paterno-Buchi reaction in this system.

The photochemical reactions of phthalimides have been the subject of numerous recent reports. The system undergoes a variety of photoreactions common to other carbonyl compounds including Type II and photoreduction reactions.<sup>1</sup> We recently reported<sup>2</sup> the novel addition of a series of dienes to N-methylphthalimide (1) to give 6-substituted 3,4-benzo-6,7-dihydro-1-methylazepine-2,5-diones in what is formally a  $[\pi^2 + \sigma^2]$  process.

In this paper we report the reaction of vinyl ethers with 1. Irradiation of 1 and a 50-fold molar excess of ethyl vinyl ether in acetonitrile solution with a 450W Hanovia lamp gave three products which were identified as 3,4-benzo-6,7-dihydro-1-methylazepine-2,5-dione (4<sup>2</sup>, 8% yield), 3-methylene-2-methylazaindanone (7, 7% yield) identified on the basis of its NMR(CDCl<sub>3</sub>)  $\delta$ 7.40-7.90 (m, 4H), 5.19 and 4.81 (AB pattern,



$J=2\text{Hz}$ ) and 3.28 (s, 3H), which was identical with that of an authentic sample,<sup>3</sup> and the oxetane 5 (9% yield, m.p. 134-135°C) whose structure was based on the following spectral properties: NMR(CDCl<sub>3</sub>)  $\delta$ 7.23-7.78 (m, 4H), 4.59, 4.30 and 3.99 (ABX

pattern,  $J_{AB}=4.6\text{Hz}$ ,  $J_{AX}=J_{BX}=6.8\text{Hz}$ ), 3.70-3.02 (m, 6H), 3.12 (s, 3H) and 1.20 (t, 3H); IR(CHCl<sub>3</sub>) 1640 cm<sup>-1</sup>; m/e 259.<sup>4</sup>

Compound 4 arises via initial [ $\pi^2 + \sigma^2$ ] photochemical addition of 2 to 1 followed by subsequent type II cleavage of the primary photoproduct 3.<sup>2</sup> The secondary photoproduct 5 is generated by a Paterno-Buchi<sup>5</sup> reaction on 4 and this was confirmed by irradiating a mixture of 4 and 2 which afforded 5 in 47% yield.

Competing with the formation of 4 is the direct  $\pi^2 + \pi^2$  Paterno-Buchi reaction of 2 with 1 to afford 6.<sup>6</sup> Compound 6 is sufficiently unstable to the reaction conditions that it decomposes to 7,<sup>10</sup> which is observed.

Irradiation of 1 with n-butyl vinyl ether gave the corresponding (to 5) butyl oxetane, 4 and 7 in 4%, 20% and 10% yields respectively. However, when the reaction was attempted with isopropenyl ethyl ether or 1,1-dimethoxy ethylene, alkenes which we expected to be more reactive, no benzazepinediones were formed. The reasons for this selectivity are being investigated.

Acknowledgement. This research was partially supported by grants from NIH (DA 01366) and NSF (02667).

#### References

1. Y. Sato, H. Nakai, T. Mizoguchi, Y. Hatanaka and Y. Kanaoka, J. Am. Chem. Soc., 98, 2349 (1976); H. Roth and D. Hundeshogen, Arch. Pharm., 309, 58 (1976).
2. P.H. Mazzocchi, M. Bowen and N. Narain, J. Am. Chem. Soc., 99, 7063 (1977).

3. W.S. Ang and B. Halton, Aust. J. Chem., 24, 851 (1971).
4. This material gave acceptable elemental analysis (C,H,N).
5. D.R. Arnold, Advan. Photochem., 6, 301 (1968).
6. We have not observed this process with dienes<sup>2</sup> or alkenes,<sup>7</sup> however, intramolecular oxetane formation has been observed on irradiation of N-alkenylsuccinimides<sup>8</sup> but not in the corresponding N-alkenylphthalimides.<sup>9</sup>
7. P.H. Mazzocchi, S. Minamikawa and M. Bowen, J. Org. Chem., 43, 3079 (1978).
8. K. Maruyama and Y. Kubo, J. Org. Chem., 42, 3215 (1977).
9. K. Maruyama, Y. Kubo, M. Machida, K. Oda, Y. Kanaoka and K. Fukuyama, J. Org. Chem., 43, 2303 (1978).
10. Similar oxetanes have been shown to be unstable to traces of acid; N. Turro and P.A. Wriede, J. Am. Chem. Soc., 90, 6963 (1968).
11. The formation of 6 corresponds to one of two possible modes of addition to the carbonyl group in 1. We do not observe the isomeric product or decomposition products from it.

Received, 19th August, 1978