

## A NEW METHOD FOR CONSTRUCTING PYRIDINE RING: THERMOLYSIS OF OXIME O-ALLYL ETHER

Junko Koyama and Yukio Suzuta

Kobe Women's College of Pharmacy, Motoyamakita, Higashinada, Kobe, Japan

Hiroshi Irie and Itsuko Katayama

Faculty of Pharmaceutical Sciences, Kyoto University, Sakyo-ku, Kyoto, Japan

Thermolysis reaction of cyclohexanone oxime O-allyl ether under air gave 5,6,7,8-tetrahydroquinoline.

In order to extend the applicability of this reaction, several cycloalkanone oxime O-allyl ethers were subjected to thermolysis reaction: oxime O-allyl ethers of cyclopentanone, cycloheptanone, cyclooctanone, and cyclododecanone yielded cyclopenteno-, cyclohepteno-, cycloocteno-, and cyclododeceno-pyridine in reasonable yield, respectively. Treatment of O-methylallyl ether of cyclohexanone oxime under the same condition gave 3-methyl-5,6,7,8-tetrahydroquinoline as a sole product, while  $\alpha$ -methyl- and  $\gamma$ -methyl-allyl ether of cyclohexanone oxime gave a mixture of 2- and 4-methyl-5,6,7,8-tetrahydroquinoline as a result of randomization of methyl group.

Furthermore, thermolysis of oxime O-allyl ethers of di-n-propyl and di-n-butyl ketone furnished  $\alpha$ -n-propyl- $\beta$ -ethyl- and  $\alpha$ -n-butyl- $\beta$ -n-propyl-pyridine, respectively

Based on the above findings, it is suggested that this new type of reaction is a method for constructing pyridine nucleus.