

C-GLYCOSYL NUCLEOSIDE XVII. A NOVEL REACTION OF SCHIFF BASES  
WITH MERCURIC CHLORIDE IN DIMETHYL SULFOXIDE<sup>1</sup>

Haruo Ogura<sup>\*</sup> and Masakazu Sakaguchi

School of Pharmaceutical Sciences, Kitasato University,  
Shirokane, Minato-ku, Tokyo 108, Japan

Toshihiko Okamoto

Faculty of Pharmaceutical Sciences, University of Tokyo,  
Hongo, Bunkyo-ku, Tokyo 113, Japan

Kinji Gonda<sup>2</sup> and Shozo Koga

Institute of Applied Microbiology, University of Tokyo,  
Yayoi, Bunkyo-ku, Tokyo 113, Japan

Pteridine or theophylline derivatives were obtained from the reaction of Schiff bases of 5,6-diamino-1,3-dimethyluracil with HgCl<sub>2</sub> in dimethylsulfoxide via radical or ionic mechanisms which were confirmed by the studies of time dependent ESR and NMR spectra. These reactions are useful for the preparation of nucleoside analogs.

In relation to synthetic studies of nucleoside analogs, cyclization of Schiff bases of 5,6-diamino-1,3-dimethyluracil with aldehydic sugars in the presence of NBS or DDQ afforded theophylline C-nucleoside or pteridine C-nucleoside analogs, respectively.<sup>3</sup> Karrer *et al.*,<sup>4</sup> Angier *et al.*,<sup>5</sup> and Baugh *et al.*<sup>6</sup> have been reported a reaction of diaminopyrimidine and (+)-glyceraldehyde to form pteridine derivatives without proof of the intermediate.

The nucleoside binding metal ions are of prime biological importance and NMR studies on the binding have been reported by Li *et al.*<sup>7</sup> and Shimokawa *et al.*<sup>8</sup> Recently, we reported the interactions of HgCl<sub>2</sub> and CdCl<sub>2</sub> with several naturally occurring nucleosides by means of NMR studies.<sup>9</sup>

In this report we would like to show the formation of pteridine or theophylline derivatives under mild conditions, from the reactions of Schiff bases



The reaction of acetates (3a,b,c) gave theophylline nucleoside analogs (4a,b,c) in 35-40% yields under the same reaction conditions, but radical peaks could not be observed under similar conditions in the ESR spectra. The structures of 4a,b,c agreed with the theophylline nucleoside analogs which were obtained by the reaction of Schiff base acetates with NBS.<sup>3</sup>







Chart 2




Figure 1

Schiff base of 5,6-diamino-1,3-dimethyluracil with (+)-glyceraldehyde (5) was treated with  $\text{HgCl}_2$  under the same conditions to afford 1,3,7-trimethylpteridine-2,4-dione (6)<sup>12</sup> (63% yield) and 15% of bis-theophylline derivative (7).<sup>13</sup> The intermediate (B) was determined by the time dependent NMR spectra. Methylene protons at 7-position were estimated at 4.73 ppm (singlet), and methyl protons at 7-position of 6 were estimated at 2.45 ppm. (Chart 2)

A similar reaction occurred when Schiff base of DAMN with (+)-glyceraldehyde (8) was treated with  $\text{HgCl}_2$  under the same conditions, and 2,3-dicyano-5-methylpyrazine (9) was obtained in 65% yield via ionic mechanism from the ESR spectra by using the spin trapping reagents. The reaction of Schiff base of 5,6-diamino-1,3-dimethyluracil with glycolaldehyde (10) under the same conditions formed 1,3-dimethylpteridine-2,4-dione (11)<sup>14</sup> in 62% yield.

Schiff base of 5,6-diamino-1,3-dimethyluracil with benzaldehyde (12) was treated in a similar manner, the reaction proceeded under an ionic mechanism (from ESR spectra by using the spin trapping reagents) and 8-phenyltheophylline (13)<sup>15</sup> was obtained quantitatively.

**Acknowledgment** This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (187147).

#### References and Notes

1. This constitutes Part XXXI of a series entitled "Studies on Heterocyclic Compounds."

2. Present address: Department of Chemistry, Tokyo Denki University, Hatoyama-mura, Hiki-gun, Saitama 350-03, Japan.
3. H. Ogura, H. Takahashi, and M. Sakaguchi, Heterocycles, 1975, 3, 93.
4. P. Karrer and R. Schwyzer, Helv. Chim. Acta, 1948, 31, 777.
5. R. B. Angier, C. W. Waller, J. H. Botthe, J. H. Mowat, J. Semb, B. L. Hutchings, E. L. R. Stokstad, and Y. SubbaRow, J. Am. Chem. Soc., 1948, 70, 3029.
6. C. M. Baugh and E. Shaw, J. Org. Chem., 1964, 29, 3610.
7. S. M. Wang and N. C. Li, J. Am. Chem. Soc., 1968, 90, 5069; L. S. Kan and N. C. Li, ibid., 1970, 92, 281, 4823; W. R. Walker, J. M. Guo, and N. C. Li, Aust. J. Chem., 1973, 26, 2391.
8. S. Shimokawa, H. Fukui, J. Sohma, and K. Hotta, J. Am. Chem. Soc., 1973, 95, 1777.
9. K. Gonda, S. Koga, M. Sakaguchi, Y. Miyata, H. Ogura, and T. Okamoto, Yakugaku Zasshi, 1978, 98, 708.
10. a) E. C. Taylor, Jr., H. M. Loux, E. A. Falco, and G. H. Hitchings, J. Am. Chem. Soc., 1955, 77, 2243. b) F. Yoneda and S. Nishigaki, Chem. Pharm. Bull. (Tokyo), 1971, 19, 1060. c) R. Marumoto and Y. Furukawa, Chem. Pharm. Bull. (Tokyo), 1977, 25, 2974.
11. A. J. Fatiadi, "Carbohydrates in Solution," ed by R. F. Gould, American Chemical Society, 1973, p. 88.
12. G. Henseke and H. G. Patzwaltdt, Chem. Ber., 1956, 89, 2904.
13. H. Fuchs, M. Gottlieb, and W. Pfeleiderer, Chem. Ber., 1978, 111, 982.
14. A. Albert, D. J. Brown, and H. C. S. Wood, J. Chem. Soc., 1956, 2066.
15. E. C. Taylor and E. E. Garcia, J. Am. Chem. Soc., 1964, 86, 4721.

Received, 17th November, 1978