

## RING EXPANSION REACTIONS OF THIIRENE 1,1-DIOXIDES

Yasuo Yoshida, Masatoshi Uesaka, Mitsuo Komatsu, Yoshiki Ohshiro, and Toshio Agawa

Department of Petroleum Chemistry, Faculty of Engineering

Osaka University, Yamadakami, Suita, Osaka 565, Japan

Thiirenes are one of the 3-heterocyclopropenes which have been of great interest for many chemists not only because of their highly strained structures but also because of their antiaromaticity. While no thiirenes have been isolated, a few thiirene 1,1-dioxides have been prepared and studied from both structural and theoretical points of view. However chemical properties of thiirene 1,1-dioxides are not well clarified. So we studied the nucleophilic addition of metalated nitriles along with cycloadditions of 1,3-dipoles and an amino butadiene derivative, and obtained interesting results.

1) Reactions of thiirene 1,1-dioxides with metalated nitriles

2,3-Diphenylthiirene 1,1-dioxide (1a) was found to act as an ambident electrophile to afford two types of sulfur-containing cyclic products, 5-imino-2-sulfolenes, and a sulfolene-4-one, when the metalated nitriles have no  $\alpha$ -hydrogen atom. The former arose from nucleophilic attack of the aryl-substituted carbanions to the ring carbon of 1a and the latter arose from the attack of an alkyl-substituted carbanion to the sulfur atom. When nitriles bear an  $\alpha$ -hydrogen atom, an acrylonitrile derivative and vinyl sulfones were obtained. On the other hand, 2,3-diphenylthiirene 1,1-dioxide (1b) suffered only one type of nucleophilic attack to the ring carbon atom, giving the sulfinates salts.

2) Cycloaddition reactions of thiirene 1,1-dioxides with nitrilium betains and an amino butadiene derivative

Thiirene 1,1-dioxides 1a,b underwent 1,3-dipolar cycloaddition with diphenylnitrile imine and p-nitrobenzotrile benzylide, with a loss of sulfur dioxide, to give pyrazoles and pyrroles, respectively. In the latter reaction, a 4H-1,4-thiazine 1,1-dioxide derivative, corresponding to a 1:1 cycloadduct of diphenylthiirene 1,1-dioxide (1b) and the nitrile ylide, was formed as a major product along with a pyrrole derivative. However in the case of 1a no such a 1:1 adduct but only a pyrrole derivative was obtained. On the other hand, 1b was found to undergo Diels-Alder-type cycloaddition with 1-piperidino-2-methyl-1,3-pentadiene. Such a reaction seems to be the first example of Diels-Alder reaction of thiirene 1,1-dioxide.