

2-AZABICYCLO[3.2.0]HEPTANE-3,4-DIONES (6)

A NOVEL METHOD OF REGIO-CONTROLLED SYNTHESIS OF FUNCTIONALIZED
HYDROINDOLES AND ERYTHRINAN DERIVATIVES¹

Takehiro Sano^{*}, Jun Toda, Yoshie Horiguchi, and Kazue Imafuku

Showa College of Pharmaceutical Sciences, Setagaya-Ku, Tokyo 154, Japan

Yoshisuke Tsuda

Faculty of Pharmaceutical Sciences, Kanazawa University, Kanazawa 920, Japan

A regio-controlled synthesis of functionalized hydroindoles from a Δ^2 -pyrrolinedione and its application to the synthesis of erythrinan skeleton were described. The method is to construct with [2+2] photocycloaddition of a trimethylsilyloxybutadiene to a Δ^2 -pyrrolinedione followed by thermal [1,3]-rearrangement to yield a hydroindole silyl enol ether, which is isomeric with the hydroindole silyl enol ether obtained by Diels-Alder cyclization between the same substrates.

During the investigation of thermolysis of 2-azabicyclo[3.2.0]heptane-3,4-diones,²⁻⁴ we have found that the 7-*exo*-vinyl derivative (1, R=H) undergoes thermal [1,3]-rearrangement to yield hydroindole (3) via homolytic cleavage of C₁-C₇ bond.⁴ A stabilizing effect of C₇-substituent on the radical (2) wherein formed and the known fact that the electron donating substituent such as trimethylsilyloxy group at a migrating center (C₇) accelerates a thermal [1,3]-shift⁵, directly leads to the idea that a vinyl oxycyclobutane derivative (1, R=OTMS) may be a potential precursor for synthesis of functionalized hydroindoles.




Chart 1

The desired 7-vinyl-7-trimethylsilyloxy derivatives (5a-c) were readily prepared by photocycloaddition ($>300\text{ nm}$)⁶ of 2-trimethylsilyloxybutadiene (2 molar equiv.) to 1-aryl- Δ^2 -pyrroline-4,5-diones (4a-c). The reaction proceeded in regio- and stereoselective manner in every case to give the product (5) homogeneously. The structures of 5a-c were assigned from the results of analogous photocycloaddition⁷ and their NMR spectra (Table 1) except for stereochemistry at C, which is still to be established.⁸

On heating the adducts (5a-c) in boiling toluene for 3 hr, an facile [1,3]-shift occurred as expected to give the hydroindole (6a-c) in the yields shown in Table 2. The presence of the silyl enol ether moiety in 6 was easily recognized by their NMR and IR spectra. On hydrolysis with 5% HCl-THF (1:1) at room temp. for 30 min., 6 afforded the corresponding diketone (7) in quantitative yield, while on treatment with potassium fluoride (in THF, r.t., overnight), it gave the ketol (8) exclusively, as a result of an intramolecular aldol condensation of 7. They were easily distinguishable by IR spectra (Table 3) and their structures were confirmed by the following spectral evidence. In ^{13}C -NMR spectrum the ketol (8a) exhibited three $-\overset{\text{O}}{\text{C}}-$ (δ 67.6, 68.0, 88.1), one $-\overset{\text{H}}{\text{C}}-$ (δ 53.4) and one ketonic carbon (δ 207.4), while the diketone (7a) showed two $-\overset{\text{O}}{\text{C}}-$ (δ 60.2, 62.9), and two ketonic carbons (δ 204.8, 196.1), but no signal of $-\overset{\text{H}}{\text{C}}-$. The structure (8a) was further verified by the following chemical transformation.




Chart 2

Table 1. Photocycloaddition⁶ of 2-Trimethylsilyloxybutadiene to 4.

| 1,2-Adduct | | NMR(δ , CDCl ₃) | | IR(Nujol, cm ⁻¹) |
|----------------------|----------|-------------------------------------|--|------------------------------|
| mp | yield(%) | vinyl protons ^{*1} | C ₆ -H(J, Hz) ^{*2} | |
| <u>5a</u> : 176-178° | 70 | 5.05-5.57(3H) | 2.37, 3.53(14) | 1780, 1760, 1730 |
| <u>5b</u> : 141-143° | 60 | 5.32-6.37(3H) | 2.33, 3.38(13) | 1760, 1720 |
| <u>5c</u> : 169-171° | 50 | 5.03-5.53(3H) | 2.29, 3.41(13) | 1780, 1730 |

*1 Signals appeared as a typical ABM pattern.

*2 Each signal appeared as a doublet.

 Table 2. Pyrolysis of The 1,2-Adduct (5)

| Product | | NMR(δ , CDCl ₃) | | IR(Nujol, cm ⁻¹) |
|-----------------------------|------------------|-------------------------------------|--------------------|------------------------------|
| mp | yield(%) | OTMS ^{*1} | -CH= ^{*2} | |
| <u>6a</u> : — ^{*3} | 58 ^{*4} | | | |
| <u>6b</u> : 144-146° | 51 | 0.18 | 4.75 | 1765, 1720, 1645 |
| <u>6c</u> : 172-178° | 58 | 0.18 | 4.66 | 1775, 1750, 1730, 1650 |

*1 The signal appeared as a 9H singlet.

*2 The signal appeared as a 1H multiplet.

*3 The silyl enol ether of 6a was cleaved on trituration with ether to give 8a.

*4 Yield was calculated from the weight of 8a.

 Table 3. Physical Data of The Diketones (7) and The Ketols (8)

| Diketone (<u>7</u>) | | Ketol (<u>8</u>) | |
|-----------------------|------------------------------|----------------------|------------------------------|
| mp | IR(Nujol, cm ⁻¹) | mp | IR(Nujol, cm ⁻¹) |
| <u>7a</u> : 150-152° | 1770, 1740, 1720, 1710 | <u>8a</u> : 195-196° | 3200, 1770, 1725 |
| <u>7b</u> : 141-142° | 1765, 1740, 1720, 1710 | <u>8b</u> : 175-176° | 3200, 1760, 1725, 1695 |
| <u>7c</u> : 152-155° | 1770, 1745, 1725, 1700 | <u>8c</u> : 222-226° | 3480, 3330, 1755, 1730 |
| | | | 1700 |

Sodium borohydride reduction of **8a** in ethanol gave a mixture of isomeric alcohols (**9**) which, without separation, were mesylated in pyridine with methanesulfonyl chloride and the product was treated with DBU to give a compound (**10**), mp 229-231°, which was identical with the Diels-Alder adduct of **4a** and butadiene.⁹




Chart 3

It is worth noting that the silyl enol ether (**6**) (hence the diketone **7**) obtained in this reaction is isomeric with the silyl enol ether (**11**) (hence the diketone **12**) prepared by the Diels-Alder reaction between the same substrates (**4** and 2-trimethylsilyloxybutadiene).¹⁰

The above method was applied to the isoquinolinopyrrolinedione (**13**)¹⁰ which was expected to lead to erythrinan derivative. The photocycloaddition⁶ of 2-trimethylsilyloxybutadiene to **13** also proceeded in regio- and stereoselective manner to give a single 1,2-adduct (**14**)¹¹ in 90% yield, which underwent thermal [1,3]-shift in boiling toluene (3 hr) to give the erythrinan derivative (**15**)¹¹ in 64% yield. Hydrolysis of **15** with 5% HCl-THF (1:1) or KF in THF yielded the ketone (**16**), mp 283-284°, in 90% yield, identity of which with the known trioxo compound (**16**)¹² was confirmed by their direct comparisons (mp, TLC, IR and NMR). Again the ketone (**16**) was isomeric with the ketone (**17**) in the position of the carbonyl moiety in ring D of the erythrinan skeleton; the latter being prepared via the Diels-Alder reaction of **13** and 2-trimethylsilyloxybutadiene.¹⁰ 1-Methoxy-3-trimethylsilyloxybutadiene underwent cycloaddition (72%) to **13** giving the product (**18**)¹³ which was then rearranged in boiling toluene (3 hr) into the erythrinan (**19**)¹⁴ in the similar manner as above. Hydrolysis of **19** with boiling 5% HCl-THF (1:1) under reflux (1 hr) afforded the conjugated enone (**20**)¹³ (38% from **18**) and treatment of **19** with KF in THF (r.t., overnight) gave the ketol (**21**)¹³ (48% from **18**), respectively.




Chart 4

The present method for the preparation of functionalized hydroindoles and erythrinan derivatives has a great synthetic value in the points that it shows a reverse regioselectivity to Diels-Alder reaction and that the silyl enol ether whereon produced is appropriate to further manipulation.

Acknowledgement: A part of this work was supported by Grant-in-Aid for Special Project Research from the Ministry of Education, Sciences and Culture, Chemical Research in Development and Utilization of Nitrogen-Organic Resources, and Naito Research Grant, for which we are grateful.

Reference and Note

1. Dioxopyrrolines XXII. This paper also constitutes Part (5) of Syntheses of Erythrina and Related Alkaloids. Part XXI: T. Sano, J. Toda, N. Kashiwaba, Y. Tsuda, and Y. Iitaka, Heterocycles, 1981, 16, 1151.
2. T. Sano, Y. Horiguchi, and Y. Tsuda, Heterocycles, 1981, 16, 355.
3. T. Sano, Y. Horiguchi, and Y. Tsuda, Heterocycles, 1981, 16, 889.
4. T. Sano, Y. Horiguchi, S. Kambe, J. Toda, J. Taga, and Y. Tsuda, Heterocycles, 1981, 16, 893.
5. a) B. M. Trost and M. J. Bogdonowicz, J. Am. Chem. Soc., 1973, 95, 289.
b) R. W. Thies and J. E. Billigmeier, J. Am. Chem. Soc., 1974, 96, 200.
6. Irradiated in dimethoxyethane with high pressure mercury lamp (300w, pyrex filtered) at 0°C for 30 min.
7. T. Sano, Y. Horiguchi, and Y. Tsuda, Heterocycles, 1981, 16, 359.
8. The result of the thermal reaction described here is compatible with that of the *exo*-vinyl derivative (1, R=H), thus suggesting C₇-*exo*-vinyl and *endo*-OTMS configuration. However, the fact does not necessarily confirm the stereochemistry since the possibility of the thermal epimerization of C₇-substituents via the biradical intermediate (2, R=OTMS) prior to the [1,3]-shift can not be eliminated.
9. T. Sano and Y. Tsuda, Heterocycles, 1976, 4, 1361.
10. See Part XXI.
11. 14: mp 120-122°. IR(Nujol); 1770, 1720 cm⁻¹. ¹H-NMR(CDCl₃); δ 0.12(9H, s, Me₃Si-), 0.83(3H, t, J=7 Hz, -COOCH₂CH₃), 2.27 and 3.53(each 1H, d, J=14 Hz, -CH₂- on cyclobutane), 3.82 and 3.88(each 3H, s, OMe), 3.93(2H, q, J=7 Hz), 5.05-5.55(3H, m, -CH=CH₂), 6.50 and 6.63(each 1H, s, Ar-H).
15: mp 158-160°. IR(Nujol); 1765, 1740, 1705, 1650 cm⁻¹. ¹H-NMR(CDCl₃); δ 0.20(9H, s, Me₃Si-), 0.67(3H, t, J=7 Hz, -COOCH₂CH₃), 2.68(2H, d, J=5 Hz, C₄-H), 3.00(2H, bs, C₁-H), 3.61(2H, q, J=7 Hz, -COOCH₂CH₃), 3.87(6H, s, 2×OMe), 4.64(1H, d, J=5 Hz, olefinic H), 6.58 and 6.63(each 1H, s, Ar-H).
12. Y. Tsuda, Y. Sakai, and T. Sano, Heterocycles, 1981, 15, 1097.
13. 18: mp 127-128°. IR(Nujol); 1760, 1730, 1710, 1660 cm⁻¹. ¹H-NMR(CDCl₃); δ 0.06(9H, s, -SiMe₃), 0.77(3H, t, J=7 Hz, -COOCH₂CH₃), 2.26 and 3.37(each 1H, d, J=14 Hz, -CH₂- on cyclobutane), 3.34, 3.80 and 3.87(each 3H, s, -OMe), 4.35 and 6.47(each 1H, d, J=14 Hz, -CH=CH-), 6.47 and 6.58(each 1H, s, Ar-H).
20: mp 170-172°. IR(Nujol); 1765, 1730, 1710, 1680 cm⁻¹. ¹H-NMR(CDCl₃); δ 0.77(3H, t, J=8 Hz, -COOCH₂CH₃), 3.94(2H, bs, C₁-H), 3.82 and 3.88(each 3H, s, -OMe), 6.06 and 6.80(each 1H, d, J=12 Hz, -COCH=CH-), 6.53 and 6.67(each 1H, s, Ar-H).
21: mp 196-198°. IR(Nujol); 3200, 1760, 1710, 1685 cm⁻¹. ¹H-NMR(CDCl₃); δ 1.05(3H, t, J=8 Hz, -COOCH₂CH₃), 3.31(3H, s, -OMe), 3.90(6H, s, 2×OMe), 6.69 and 6.78(each 1H, s, Ar-H).
14. The silyl enol ether was too labile to be isolated in pure form and changed into 21 on direct trituration with ether.

Received, 6th May, 1981