

THE FACILE ADDITION OF SECONDARY AMINES TO WITHAFERIN A. A FURTHER EXAMPLE OF ALUMINA-CATALYZED ADDITION OF AMINES TO  $\alpha, \beta$ -UNSATURATED KETONES

S. William Pelletier\*, Gulilat Gebeyehu and Naresh V. Mody  
Institute for Natural Products Research and The Department of Chemistry,  
The University of Georgia, Athens, Georgia 30602, USA

**Abstract**—The alumina-catalyzed addition of various secondary amines to withaferin A (1) is facile and efficient. The addition is illustrated by the preparation in excellent yield of 3-aziridinyl-2,3-dihydrowithaferin A (2), a compound with significant *in vivo* antitumor activity.

The introduction of an amine functionality in the steroid nucleus is known to alter biological activity significantly.<sup>1</sup> Synthetic and naturally-occurring 3-aminosteroids are known to possess important biological activities, such as CNS, anaesthetic and neuromuscular blocking.<sup>2</sup> For a study of the relationship of structure with antitumor activity, we required a simple and efficient method of introducing an amine functionality at the C(3) position of withaferin A (1). As earlier investigators observed<sup>3,5</sup> that alcohols add to the enone moiety of withaferin A with ease (especially in the presence of a base such as DBU), we anticipated that amines, and in particular secondary amines, would add in a similar fashion to give the required amino adduct. However, treatment of withaferin A (1) with aziridine or diethylamine resulted in the formation of a complex mixture of products, and the reaction did not proceed to completion within two days. Hence we took advantage of our recent discovery of the use of alumina as a catalyst for the addition of secondary amines to exocyclic  $\alpha, \beta$ -unsaturated ketones.<sup>6</sup>

In this paper we describe an alumina-catalyzed addition of secondary amines to the enone moiety of withaferin A, a reaction which is facile and efficient. The results of addition of nine secondary amines to withaferin A are presented in Table I.













Table I: Addition of Secondary Amines to Withaferin A

Compound	R	Reaction Time	%Yield†
2		1 hr	85
3		24 hr	65
4		1 hr	70
5		24 hr	68
6		5 hr	67
7		3 hr	86 ‡
8		24 hr	62*
9		48 hr	57*
10		24 hr	67*

†Yield is after flash chromatography.<sup>7</sup> All compounds were fully characterized by mass, proton & carbon-13 NMR spectra.

‡The material was triturated with  $\text{CHCl}_3$ /hexane before chromatography.

\*Yield is based on recovered withaferin A (54, 70, and 85% conversion for compounds 8, 9, and 10, respectively)

The structures of addition products were determined by  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral analysis. The values and assignments of the chemical signals of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of the products are given in Tables II and III. The  $^{13}\text{C}$  chemical shifts assignments were based on previous studies reported from our laboratory<sup>3</sup> and others.<sup>8</sup>

The following is a representative experimental procedure: 500 mg of withaferin A (1) was treated with 5 ml of aziridine<sup>9</sup> and 500 mg of  $\text{Al}_2\text{O}_3$  (activity-III)<sup>10</sup> in 120 ml of toluene at 25°C. The reaction mixture was stirred under nitrogen until the starting material was consumed (TLC, 1 hr). The solid that was obtained after filtration and evaporation was subjected to flash chromatography<sup>7</sup> [ $\text{SiO}_2$ , 40-63  $\mu\text{m}$ ; 3% EtOH in EtOAc] to give 463 mg of 3-aziridinyl-2,3-dihydrowithaferin A (2). Crystallization from EtOAc/hexane gave compound 2 as white cubes, mp 205.5 -207.5°C (corr.).

Table II.  $^{13}\text{C}$  Chemical Shifts and Assignments of Compounds 2-10<sup>†</sup>

	2	3	4	5	6	7	8	9	10
C(1)	210.6	211.5	211.5	211.3	211.3	211.1	211.1	210.3	211.0
C(2)	39.8	36.2	38.7	36.4	37.2	36.5	37.3	36.6	37.9
C(3)	68.9	65.7	64.5	63.9	64.4	63.9	59.6	55.8	63.2
C(4)	77.1	78.0	75.8	73.3	73.2	74.0	73.8	74.3	74.0
C(5)	65.6	65.7	65.8	64.6	63.7	65.2	64.6	64.7	65.0
C(6)	61.1	60.2	61.4	59.9	59.3	61.0	60.0	59.7	60.0
C(7)	29.8	29.6	29.8	29.9	29.9	29.9	29.8	29.8	29.8
C(8)	31.1	31.1	31.2	31.1	31.2	31.4	31.2	31.3	31.3
C(9)	42.7	42.9	43.2	42.8	42.8	43.2	43.3	43.0	42.9
C(10)	50.5	50.4	50.2	49.9	49.9	50.4	49.8	50.1	50.0
C(11)	21.7	21.5	22.0	21.5	21.5	22.0	21.2	21.3	21.5
C(12)	27.3	27.2	27.2	27.3	27.3	27.3	27.2	27.2	27.2
C(13)	42.7	42.6	42.6	42.8	42.7	42.8	42.7	42.7	42.7
C(14)	56.0	56.1	55.9	56.1	56.1	56.2	56.0	56.1	56.1
C(15)	24.3	24.3	24.3	24.3	24.3	24.4	24.2	24.2	24.2
C(16)	39.2	39.2	39.3	39.2	39.2	39.3	39.1	39.1	39.1
C(17)	52.0	51.9	52.0	52.0	52.0	52.1	51.9	51.9	52.0
C(18)	11.6	11.6	11.6	11.7	11.6	11.8	11.6	11.6	11.6
C(19)	15.9	15.5	16.3	15.7	15.8	16.2	15.4	15.5	15.7
C(20)	38.8	38.8	38.7	38.9	38.8	38.9	38.8	38.8	38.8
C(21)	13.4	13.3	13.3	13.4	13.4	13.4	13.3	13.4	13.3
C(22)	78.7	78.7	78.7	78.7	78.7	78.7	78.6	78.7	78.4
C(23)	29.4	29.6	29.3	29.0	29.4	29.2	28.9	28.9	29.0
C(24)	153.1	153.3	153.3	152.9	152.9	152.9	153.2	152.9	153.2
C(25)	125.7	125.7	125.7	125.8	125.8	125.9	125.7	125.7	125.7
C(26)	167.0	167.0	166.9	167.0	167.0	167.0	166.9	166.9	167.0
C(27)	57.3	57.1	57.1	57.4	57.4	57.4	57.1	57.3	57.2
C(28)	20.1	20.0	20.0	20.0	20.0	20.0	20.0	19.9	20.0
a	26.8	65.8	51.6	50.5	51.1	50.2	42.7	48.0	37.0
b	26.4	40.9	23.3	24.5	28.9	67.2	13.1	28.3	58.4
c	-	16.9	-	26.3	26.7	-	-	129.8	138.7
d	-	-	-	-	-	-	-	124.6	127.1
e	-	-	-	-	-	-	-	118.3	128.5
f	-	-	-	-	-	-	-	127.4	128.6
g	-	-	-	-	-	-	-	107.3	-
h	-	-	-	-	-	-	-	150.2	-

<sup>†</sup> Chemical shifts are recorded in ppm downfield from TMS. The spectra were taken in  $\text{CDCl}_3$  solution 15.03 MHz in the Fourier mode using a JEOL FX-60 spectrometer in conjunction with a JEC-980 computer.

Table III. The 90 MHz Proton NMR Shifts of Compounds 2-10<sup>a</sup>

	2	3	4	5	6	7	8	9	10
C(3)-H	2.75(m)	2.80-3.06(m)	2.87-3.05(m)	2.87(m)	2.35-2.73(m)	2.87(m)	3.10(m)	3.30(m)	2.60-2.85(m)
C(4)-H	3.41(bs)	3.33(bs)	3.52(d, J=4Hz)	3.51(d, J=5Hz)	3.10-3.50(m)	3.65(m)	3.45(d, J=5Hz)	3.56(d, J=5Hz)	3.5(bs)
C(6)-H	3.28(bs)	3.20(s)	3.20(s)	3.25(s)	b	3.25(s)	3.25(s)	3.26(s)	3.20(s)
C(18)-H <sub>3</sub>	0.69(s)	0.68(s)	0.69(s)	0.67(s)	0.66(s)	0.69(s)	0.68(s)	0.67(s)	0.68(s)
C(19)-H <sub>3</sub>	1.30(s)	1.25(s)	1.34(s)	1.28(s)	1.28(s)	1.35(s)	1.25(s)	1.33(s)	1.33(s)
C(21)-H <sub>3</sub>	0.98(d, J=6Hz)	0.98(d, J=6Hz)	0.97(d, J=6Hz)	0.98(d, J=6Hz)	0.96(d, J=6Hz)	0.98(d, J=6Hz)	c	0.96(d, J=6Hz)	0.97(d, J=6Hz)
C(22)-H	4.48(m)	4.45(m)	4.45(m)	4.45(m)	4.40(m)	4.51(m)	4.42(m)	4.66(m)	4.40(m)
C(27)-H <sub>2</sub>	4.38(s)	4.38(s)	4.30(s)	4.30(s)	4.30(s)	4.38(s)	4.30(s)	4.33(s)	4.25(s)
C(28)-H <sub>3</sub>	2.05(s)	2.03(s)	2.05(s)	2.03(s)	2.03(s)	2.03(s)	2.03(s)	2.03(s)	2.02(s)
R	1.7(m)	1.09(d, J=2Hz)	2.65-2.8(m) 1.69(m)	2.63(m) 2.44(m) 1.46(bm)	2.35-2.73(m) 1.53(bs)	3.65(bs) 2.55(bs)	2.52(q, J=7Hz) 0.98(t, J=7Hz)	2.83(m) 6.57-6.96(m)	2.18(s) 3.0(bs) 7.15(bs)
		1.32(d, J=2Hz) 1.9(bs)							

<sup>a</sup> Chemical shifts are recorded in ppm downfield from TMS. The spectra were taken in CDCl<sub>3</sub> on the Varian EM-390 NMR spectrometer.

<sup>b</sup> The chemical shift is part of the multiplet at 6.3.10-3.50.

<sup>c</sup> The chemical shift was covered by the triplet at 6 0.98.

The above results demonstrate that the alumina-catalyzed addition of secondary amines to enone systems is general and compatible with various sensitive functional groups.

A preliminary study of 3-aziridinyl-2,3-dihydrowithaferin A (2) has shown promising activity against murine P-388 lymphocytic leukemia (T/C 149 at 80 mg/kg).<sup>11</sup> The results of tests of compound 2 and other amino adducts against various tumor systems will be reported in due course.

Acknowledgement: This work was supported by Grant CA 29520 (formerly CA 24180) from the National Cancer Institute.

#### References and Notes

1. W. R. Buchell, Adv. Steroid Biochem. and Pharm., **3**, 39 (1972).
2. G. Gjerstad and A. Modak, Quart. J. of Crude Drug Research, **8**, 1141 (1968).
3. D. Lavie, E. Glotter, and Y. Shiro, J. Chem. Soc., 7515 (1965).
4. S. M. Kupchan, W. K. Anderson, P. Bollinger, R. W. Daskotch, R. M. Smith, J.A.S. Renauld, H. K. Schnoes, A. L. Burlingame, and D. H. Smith, J. Org. Chem., **34**, 3858 (1969).
5. (a) S. W. Pelletier, N. V. Mody, J. Nowacki, and J. Bhattacharyya, J. Nat. Prod., **42**, 512 (1979), (b) S. W. Pelletier, G. Gebeyehu, J. Nowacki, and N. V. Mody, Heterocycles, **15**, 317 (1981).
6. S. W. Pelletier, A. P. Venkov, J. Finer-Moore, and N. V. Mody, Tetrahedron Lett., **21**, 809 (1980).
7. W. C. Still, M. Kahn, and A. Mitra, J. Org. Chem., **43**, 2923 (1978).
8. H. E. Gottlieb and I. Kirson, Org. Mag. Reson., **16**, 20 (1981).
9. K. Campbell, A. H. Sommers, and B. K. Campbell, "Organic Synthesis", John Wiley, New York, 1955, Coll. Vol. 3, p. 148.
10. E. Merck neutral alumina (90 active 70-230 mesh) for column chromatography was adjusted to activity III by addition of an appropriate amount of water.
11. We thank the National Cancer Institute for the test results.

Received, 5th October, 1981