

THE SYNTHESIS AND CHARACTERISATIONS OF RETINOIDAL
4-YLIDENEBUTENOLIDES

Masayoshi Ito,* Tsutomu Iwata, and Kiyoshi Tsukida
Kobe Women's College of Pharmacy, Motoyamakitamachi,
Higashinada, Kobe 658, Japan

Abstract — Retinoidal 4-ylidenebutenolides [(5) and (18)]
have been synthesised and their spectral characterisations have
been described.

4-Ylidenebutenolides (1) occur widely in nature and many of them are known to possess a wide range of biological activities. The chemistry of natural 4-ylidenebutenolides has been well reviewed by Pattenden¹ and Yamamoto.² Although a number of methods for the synthesis of 4-ylidenebutenolides have been presented, there has been so far reported no synthetic pathway for natural 4-ylidenebutenolides (2) displaying extended conjugation at C-2 position such as peridinin (3)³ and tetrenolin (4).⁴ As an extension of the synthetic work⁵ to develop the new antitumour retinoids, we have been interested in the preparation of the retinoidal 4-ylidenebutenolide (5).⁶ In this communication, we describe the first synthesis of the retinoidal 4-ylidenebutenolides (5) and (18) and their spectral characterisations.




Fig. 1

The aldol condensation (piperidine/MeOH) between β -cyclocitral (6)⁷ and pyruvic aldehyde dimethylacetal (7) gave the *trans* acetal-dienone (8) [δ^8 7.59 (d,J=16,7-H), 6.46(d,J=16,8-H)] in 60% yield which was condensed with diethyl methoxycarbonylmethylphosphonate in the presence of *n*-butyl-lithium to afford a mixture (9) of the β -ionylidene ester derivatives [9-*cis* : 9-*trans*⁹ = ca. 1 : 7 by HPLC] in 94% yield. The mixture was treated with 15% H₂SO₄ in MeOH to provide a methoxy-lactone (10) in 68% yield. The structure of this product was confirmed by its spectral data [λ 258, 319 nm; ν 1755, 1612 cm⁻¹; δ 6.90(d,J=16,7-H), 6.32 (d,J=16,8-H), 6.04(s,11'-H), 5.91(s,10-H), 3.50(s,OMe)]. Hydrolysis of the methoxy-lactone (10) with 30% H₂SO₄ in dioxan and subsequent PCC-oxidation of the resulting hydroxy-lactone (11) led to the unstable conjugated anhydride (12) [ν 1840, 1765 cm⁻¹] in ca. 60% yield which, without purification, was condensed with the phosphorane (13)¹⁰ in dry benzene under argon to yield a mixture of the conjugated ylidenebutenolides (14)(10%), (15)(24.5%), and (16)(8.2%). The isomers (14), (15), and (16) were clearly separated by preparative TLC [SiO₂/benzene and *n*-hexane:ether(3:2)] and fully characterised by their spectral data (Table 1) respectively. The regioselectivity of this condensation reaction is in favour of carbanion attack at the less hindered carbonyl function (C-11) in the anhydride (10). The two isomers (14) and (15) were interconverted by heating or TLC treatment.¹¹ The lactonic carbonyl absorptions of the compounds (14) and (15) were shifted to the higher frequency by the effect of exocyclic double bond in the 4-ylidenebutenolide structure (2) compared with those [1740-1760 cm⁻¹] of the normal α,β -unsaturated γ -lactone ring. In the NMR data of the compounds (14) and (15), the proton signals at C-10 and C-12 were in good agreement with those at the corresponding positions in the model compounds (19) and (20).^{12, 13} Therefore, extension of conjugation at the C-2 position in the 4-ylidenebutenolide ring (2) had little effect on the chemical shifts at C-10 and C-12 positions in the ylidenebutenolides (14) and (15). Stereochemistry around the newly formed Δ^{11} -double bond was determined as shown in the Fig. 2 from both the chemical shift difference of C-13-methyl signals and that of C-10-olefinic proton signals in the two isomers (14) and (15). The Wittig reaction of the keto-butenolide (14) with the phosphorane (17)¹⁴ in a sealed bottle under argon produced a mixture of the retinoidal 4-ylidenebutenolides (5)(25.1%) and (18)(14.0%) which was separated by preparative TLC [SiO₂/benzene]. The same reaction of the isomeric keto-




Fig. 2

Table 1 Spectral data of the keto-butenolides (14)-(16)

		(14)	(15)	(16)	
UV	$\lambda_{\text{max}}^{\text{EtOH}}$	330 nm	335 nm	293 nm	
IR	ν^{CHCl_3}	1790	1787	1785	
		1665	1686	1665	
		1620 cm^{-1}	1602 cm^{-1}	1643 1610 cm^{-1}	
NMR	δ^{CDCl_3} ppm	10-H	7.08 (s)	7.93 (s)	6.32 (s)
		12-H	5.51 (s)	6.14 (s)	-
		12'-H	-	-	5.72 (s)
		13-CH ₃	2.56 (s)	2.33 (s)	-
		13'-CH ₃	-	-	2.59 (s)

Table 2 Spectral data of the retinoidal 4-ylidenebutenolides (5) and (18)

		(5)	(18)	
UV	$\lambda_{\text{max}}^{\text{EtOH}}$	356 nm	356 nm	
IR	ν^{CHCl_3}	1770	1775	
		1765	1760	
		1712	1700	
		1600 cm^{-1}	1598 cm^{-1}	
NMR	δ^{CDCl_3} ppm	1-gemCH ₃	1.07 (s)	1.07 (s)
		5-CH ₃	1.77 (s)	1.77 (s)
		13-CH ₃	2.55 (s)	2.32 (s)
		CO ₂ CH ₃	3.73 (s)	3.72 (s)
		14-H	5.57 (s)	5.77 (s)
		12-H	6.04 (s)	7.32 (s)
		8-H	6.22 (d, J=16)	6.23 (d, J=16)
		10-H	7.00 (s)	7.09 (s)
		7-H	7.38 (d, J=16)	7.37 (d, J=16)

butenolide (15) also led to a mixture of (5)(16.8%) and (18)(9.5%). Characteristic spectral data of the retinoidal 4-ylidenebutenolides (5) and (18) are summarised in Table 2. Stereochemistry of the Δ^{13} -double bond was determined from the follow-

ing NMR data. The methyl signal at C-13 in the 13-*trans* isomer (5) is extremely deshielded by the effect of both enolic oxygen in the butenolide ring and the carbonyl group of ester. The proton signal at C-12 in the 13-*cis* isomer (18) is also deshielded by the anisotropic effect of the carbonyl group of ester. The biological activities of the retinoidal 4-ylidenebutenolides prepared here are now under investigation. The present synthetic route could be expected to extend to the studies towards the synthesis of the 4-ylidenebutenolide (2).

REFERENCES AND FOOTNOTE

1. G. Pattenden, Fortschr. Chem. Org. Naturst., 1978, 35, 133.
2. M. Yamamoto, J. Synth. Org. Chem. Japan, 1981, 39, 25.
3. J. E. Johansen, S. Liaaen-Jensen, and G. Borch, Marine Natural Products Chemistry, edited by D. J. Faulkner and W. H. Fenical, Plenum Press, p. 225 (1977) and references therein.
4. G. G. Gallo, C. Coronelli, A. Vigevani, and G. C. Lancini, Tetrahedron, 1969, 25, 5677.
5. M. Ito, M. Ohno, E. Takano, Y. Oda, and K. Tsukida, Heterocycles, 1979, 12, 505.
6. We have employed the numbering system used in the retinoids and carotenoids fields.
7. N. Müller and W. Hoffmann, Synthesis, 1975, 781.
8. NMR spectra were determined on a 200 MHz superconducting FT-NMR spectrometer using CDCl₃ solutions.
9. Geometrical isomerism is expressed on the basis of the retinoidal polyene chain.
10. F. Ramirez and S. Dershowitz, J. Org. Chem., 1957, 22, 41.
11. Heating or TLC treatment led to an equilibrium mixture (ca. 1 : 1) of (14) and (15).
12. C. F. Ingham, R. A. Massy-Westropp, G. D. Reynolds, and W. D. Thorpe, Aust. J. Chem., 1975, 28, 2499.
13. T. Kitamura, Y. Kawakami, T. Imagawa, and M. Kawanishi, Tetrahedron, 1980, 36, 1183.
14. O. Isler, H. Gutmann, M. Montavon, R. Rüegg, G. Ryser, and P. Zeller, Helv. Chim. Acta, 1957, 40, 1242.

Received, 18th March, 1982