

A NEW AND SIMPLE SYNTHESIS OF 1,2-DIAZAAZULENE

Kimiaki Imafuku* and Akiko Shimazu


Department of Chemistry, Faculty of Science,

Kumamoto University, Kurokami, Kumamoto 860, Japan


Abstract — 3-Methyl-8-methylamino-1,2-diazaazulene was synthesized in 24% yield from 3-acetyltropolone via 2-acetyl-7-methylaminotroponone and its hydrazone.

1,2-Diazaazulene is of interest for their relationship with 1,3-diazaazulenes. However, little is known about the chemistry of 1,2-diazaazulene derivatives, except for two reports by Matsumoto¹ and by Treibs.² These communications gave no details of their preparation and properties.

Recently, we found that 3-acetyltropolone (**1**) reacted with amines at room temperature to afford regioselectively 2-acetyl-7-(substituted amino)tropones.³ 2-Acetyl-7-methylaminotroponone (**2**) was readily available from the reaction of **1** with methylamine in 90% yield. In this communication, we wish to describe a new and convenient synthesis of 1,2-diazaazulene from 3-acetyltropolone (**1**).



A mixture of 2-acetyl-7-methylaminotropone (2) and hydrazine hydrate in methanol was refluxed for 2 h to afford 2-acetyl-7-methylaminotropone hydrazone (3) (66%): mp 133-135°C; Ir (CHCl₃) 3320 and 1595 cm⁻¹; Nmr (CDCl₃) δ 2.09 (3H, s, CH₃), 3.00 (3H, d, J = 6.0 Hz, N-CH₃), 5.25 (2H, br, NH₂), and 6.3-7.7 (5H, m, NH + tropone). Found: C, 62.81; H, 6.77; N, 22.14%. Calcd for C₁₀H₁₃N₃O: C, 62.80; H, 6.85; N, 21.98%. A solution of 3 in acetic acid was heated on a water bath for 1 h. After the removal of the solvent, the residue was chromatographed on a Wakogel B-10 plate to give 3-methyl-8-methylamino-1,2-diazaazulene (4) (40%): mp 174-176°C; Ir (CHCl₃) 3320, 1620, and 1580 cm⁻¹; Uv (CH₃OH) 233 (log ε 4.21), 255 (4.33), 309 (3.87), 366 (3.58), and 437 nm (3.67); Nmr (CDCl₃) δ 2.73 (3H, s, C-CH₃), 3.36 (3H, s, N-CH₃), 6.6-7.1 (2H, m), 7.55 (1H, br, NH), and 7.5-8.1 (2H, m). Found: C, 69.25; H, 6.27; N, 24.14%. Calcd for C₁₀H₁₁N₃: C, 69.34; H, 6.40; N, 24.26%. The overall yield of the 1,2-diazaazulene (4) from 1 was 24%.



Although the compound (4) has two tautomeric forms such as A and B, it seems that the structure A is reasonable from the following spectral data. The Uv spectrum of 4 is shown in Fig. 1 together with those of 3-methyl-1,8-dihydrocycloheptapyrazol-8-one (5) and its hydrazone (6),⁴ which are corresponding to the tautomeric form (B). The absorption maxima of 4 are almost consistent with Treibs' data² and were observed at fairly longer wavelengths than those of 5 and 6. Furthermore, the absorption bands in visible region showed a shift towards long wavelength region in comparison with those of 1,3-diazaazulene.^{5,6} In Nmr spectrum of 4, the signals in the seven-membered ring protons were observed at lower magnetic field than those of the compounds (5 and 6). The values of chemical shifts are comparable to the values in 1,3-diazaazulenes.⁷

Studies on reactions of the 1,2-diazaazulene (4) are now in progress.




Fig. 1. Uv and Visible Spectra

— 4, — — 5, — · — 6,
 - - - 1,3-Diazaazulene

REFERENCES

1. S. Matsumoto, Sci. Repts. Tohoku Univ., **I**, 1958, 42, 222.
2. W. Treibs, Chimia, 1972, 26, 627.
3. K. Imafuku and A. Shimazu, Synthesis, 1982, in press.
4. A. Yamane, M. Nagayoshi, K. Imafuku, and H. Matsumura, Bull. Chem. Soc. Jpn., 1979, 52, 1972.
5. T. Nozoe, T. Mukai, and I. Murata, J. Am. Chem. Soc., 1954, 76, 3352.
6. I. Murata, Tohoku Daigaku Hisui-yoeki Kagaku Kenkyusho Hokoku, 1960, 2, 99.
7. R. Cabrino, B. Ricciarelli, and F. Pietra, Tetrahedron Lett., 1974, 3069.

Received, 23rd August, 1982