

## SYNTHESIS OF FURANS AND BUTENOLIDES USING ACYLBUTENE DITHIOACETALS

Yoshio Negishi, Renji Okazaki, and Naoki Inamoto

Department of Chemistry, Faculty of Science, The University of Tokyo,  
Hongo, Tokyo 113, Japan

Reaction of acylketene dithioacetals(1) with dimethylsulfonium methylide gave 2,2-dimethylthio-2,5-dihydrofurans(2) in high yields. Mild acidic hydrolysis or treatment with Florisil of 2 afforded in good yields 2-methylthiofurans(3), severer acidic hydrolysis of which led to the formation of butenolides(4). The 2-methylthiofurans(3) were converted to various substituted furans by following reactions: A: Alkylation of 5-position by treatment with butyllithium followed by addition of alkyl halide

B: Removal of the 2-methylthio group with Raney-Ni(W-7) deactivated by acetone

C: Substitution of the 2-methylthio group with nickel-catalyzed Grignard crosscoupling

The alkylation(A) of 3 followed by hydrolysis gave 5-alkylbutenolides(5).

Two naturally occurring, isomeric furans, perillene(6) and rosefuran(7) were synthesized by the above reactions from 1,1-dimethylthio-3-oxo-1-butene(1,  $R^1=Me$ ,  $R^2=H$ ) and 1-bromo-3-methyl-2-butene.

