


A NEW SYNTHESIS OF 4H-1,4-THIAZINES AND N-ALKYL-
 4H-1,4-BENZOTHAZINES

G.Liso, G.Trapani, A.Reho, A.Latrofa, P.Marchini
 Dipartimento Farmaco-Chimico - Facoltà di Farmacia
 Via Amendola 173, 70126 Bari, Italy

We have recently reported that the reaction of 2,2'-dithiodianiline with electron-deficient alkynes constitutes a new synthetic approach to N-unsubstituted 4H-1,4-benzothiazines (J.Chem.Soc., Perkin Trans.I,1983,567).

The results of further investigation carried out in order to study the scope and limitations of this process with respect to 2,2'-diaminodisulfides are now reported. The use of the aliphatic disulfide **1** and the N-alkylated disulfide **2** allows the synthesis of 4H-1,4-thiazines **3** and N-alkyl 4H-1,4-benzothiazines **4** respectively, in satisfactory to good yields.



	R	R'	X
(3,4)a	$\overline{\text{CH}(\text{CH}_2)_5\text{CH}_2}$	H	OC_2H_5
(3,4)b	$\overline{\text{CH}(\text{CH}_2)_3\text{CH}_2}$	COOCH_3	OCH_3
(3,4)c	$\overline{\text{CH}(\text{CH}_2)_5\text{CH}_2}$	C_6H_5	CH_3
(4)d	$\overline{\text{CH}(\text{CH}_2)_5\text{CH}_2}$	H	$(\text{CH}_2)_2\text{CH}_3$