


BIS HETEROANNULATION 6. THE FIRST EXAMPLE OF A DIELS-ALDER REACTION INVOLVING A THIAZOLE RING WITH AN ACETYLENIC DIENO-PHILE. GEOMETRICAL CONTROL OF REACTION PATHWAY

Peter A. Jacobi, * Kim T. Weiss and (in part) Melissa Egbertson
Hall-Atwater Laboratories, Wesleyan University, Middletown, Connecticut
USA 06457

Abstract — Acetylenic thiazoles of proper design have been shown to undergo an intramolecular Diels-Alder reaction leading directly to fused-ring thio-
phene derivatives.


We have recently reported that acetylenic oxazoles of type 1 undergo a facile intramolecular Diels-Alder reaction, leading, via intermediate 2, to highly substituted furans of type 3 ("Bis Heteroannulation", Scheme 1).¹ This methodology has been successfully applied to the synthesis



Scheme 1

of a variety of furanosesquiterpenes,¹ and we have been intrigued with the possibility that thiazoles of type 4 might exhibit similar reactivity. Such a transformation could be of considerable practical importance since it is obvious, for example, that a β -substituted thiophene can be considered as a latent "isopreneoid" unit.² Furthermore, substitution patterns of the type generalized in 7 are both exceedingly common in nature and sometimes difficult to obtain using presently available techniques.³


A major difficulty with this approach is the lack of reactivity of thiazoles in Diels-Alder reactions.⁴ To the best of our knowledge there has been only one report of a reaction of this class,⁵ and even in this case the primary adduct was not isolated. This lack of reactivity is presumably due to the greater aromaticity of thiazoles relative to oxazoles,⁴ as well as to the high nucleophilicity of the thiazole ring.^{6,7} 2,4-Dimethylthiazole (8), for example, reacts with dimethyl acetylenedicarboxylate to give adducts 9 and 10 as the only isolated products (Scheme 2).⁷ We



Scheme 2


have now found, however, that intramolecular variants of this reaction are possible if proper geometrical restraints are imposed.

Thiazoles 11a-d were prepared following standard literature procedures,^{1,8} and they were subjected to a range of cyclization conditions (Scheme 3). Not surprisingly, both 11a and 11b



Scheme 3

reacted exclusively to give products derived from intermediates 12a and 12b (path a).⁷ In each case models clearly indicate that the 7-exo-dig transition state required for path a is highly favored,⁹ and no trace of thiophenes 13a or 13b could be detected in the crude reaction mixtures. Thus, for example, after brief heating (refluxing ethylbenzene, 3 h) all 11a was consumed and a single major product was evident by TLC ($R_f = 0.60$, 20% acetone/hexane, silica gel). This product was isolated as a highly unstable crystalline solid, mp 90 - 92° C (yield 55 - 60% yield), and assigned structure 14 on the basis of mechanistic considerations⁷ as well as the following



observations.¹⁰ First, elemental analysis and mass spectral data showed that 14 was isomeric with 11a, thereby ruling out the possibility of an intramolecular Diels-Alder reaction with loss of acetonitrile.¹ Second, the infrared spectrum (CHCl_3) of 14 was devoid of absorptions in the regions expected for hydroxyl or acetylenic bond stretching, and all attempts at preparing an acetate derivative failed. The major IR peaks (1690 , 1605 cm^{-1}), however, agreed precisely with those expected for a β -aminoacrylate system (cf. compound 15, IR 1690 , 1595 cm^{-1})¹¹. Third, the ^{13}C NMR data was in full accord with structure 14 and in particular indicated the absence of an aromatic thiazole ring.¹⁰ Absorptions for C-2 and C-3, however, were readily apparent at 137.5 and 114.7 ppm, and of particular importance the assigned values for C-5, C-15, and C-16 (159.2 , 84.6 and 167.9 ppm, respectively) are in excellent agreement with those found in the model system 15 (164.1 , 77.3 and 169.7 ppm, respectively, for C-11, C-14 and C-15).¹¹ These absorptions would appear to be highly characteristic of the β -aminoacrylate system. In addition, the position of absorption for C-10 (110.3 ppm) is in total agreement with the closely related system 16¹² (95.1 ppm for C-6) after correcting for standard incremental substituent effects.¹² Fourth, the ^1H NMR spectrum of 14, including exhaustive decoupling experiments, is in full accord with the assignments made above.¹⁰ In particular, the E-configuration at C₅ - C₁₅ is based on the logical expectation of a facile equilibration to the thermodynamically most stable isomer, as well as the strongly deshielding influence of the ester functionality on H-6 (5.91 ppm). Finally, it is interesting to note that although 14 is undoubtedly an unusual structure, Dreiding models of this compound are virtually free of strain.

In contrast to the results presented above neither 11c nor 11d are electronically disposed to undergo the 7-exo-dig process, and the alternative 8-endo-dig cyclization is geometrically impossible. Therefore, we believed, these materials should be more amenable to undergoing an intramolecular Diels-Alder reaction. Our initial efforts in this area, however, were unrewarding. Thus, thiazole 11c suffered mainly slow decomposition to polar products over a period of several days in refluxing ethylbenzene (~ 138° C), and all attempts at catalyzing this process were either ineffective (BF₃ · Et₂O, Hg(OAc)₂, TsOH, ZnCl₂), or led to addition reactions at the acetylenic triple bond (AlCl₃).¹³ Also, higher temperatures led to rapid decomposition which could not be controlled by the addition of various radical scavengers, including, for example, 3-tert-butyl-4-hydroxy-5-methylphenyl sulfide and 4,4'-butylidene-bis-6-tert-butyl-m-cresol.¹⁴ With thiazole 11d, however, the additional activating influence of a methoxyl group apparently facilitates the Diels-Alder process. Thus, in the presence of a catalytic amount of methylene blue,¹⁵ 11d gave a 60% yield of thiophene 13d after 3 days at reflux in degassed mesitylene (165°). That 13d was actually in hand was unequivocally established by all of the usual criteria,¹⁶ as well as the fact that its ¹H NMR spectrum was virtually superimposable with that of the previously described furan analog [3-methoxymethyl-4-keto-6-methyl-4,5,6,7-tetrahydrobenzofuran].¹⁷ This example, then, provides the first conclusive evidence that thiazoles can undergo the Diels-Alder reaction with acetylenic dienophiles in exactly analogous fashion as their oxazole counterparts.

In closing, we might only add that the intramolecular nature of these reactions should be even more favorable in multicyclic systems such as 4 (A, B = cycloalkyl) since the reacting centers are held more rigidly in proximity to each other. Therefore, we believe, reactions of this type might find considerable applicability in the synthesis of naturally occurring terpenes.

ACKNOWLEDGMENT Financial support of this work by the National Science Foundation (Grant No. CHE-8108984) is gratefully acknowledged. The Varian XL-200 spectrometer used in this work was financed in part by the National Science Foundation (Grant No. CHE-7908593), the Dreyfus Foundation, and Wesleyan University.

REFERENCES AND NOTES

1. P.A. Jacobi, and T.A. Craig, *J. Am. Chem. Soc.*, 1978, 100, 7748. (b) P.A. Jacobi, D.G. Walker and I. Odeh, *J. Org. Chem.*, 1981, 46, 2065. (c) P.A. Jacobi and D.G. Walker, *J. Am. Chem. Soc.*, 1981, 103, 4611. (d) P.A. Jacobi, T.A. Craig and B.A. Arrick, *J. Am. Chem. Soc.*, submitted. (e) P.A. Jacobi, D.G. Walker and R.F. Frechette, *J. Am. Chem. Soc.*, submitted.
2. (a) A.I. Meyers, "Heterocycles in Organic Synthesis", John Wiley and Sons, Inc., New York, 1974. (b) G.R. Pettit and E.E. van Tamelen, *Org. React.*, 1962, 12, 356.

3. (a) T.K. Devon and A.I. Scott, "Handbook of Naturally Occurring Compounds", Academic Press, New York, 1972, Vol. II. For leading synthetic references in this area see (b) T. C. T. Chang and M. Rosenblum, J. Org. Chem., 1981, 46, 4103; and (c) S. F. Martin and T. Chou, J. Org. Chem., 1978, 43, 1027.
4. (a) J. V. Metzger, Ed., "Thiazole and its Derivatives", John Wiley and Sons, Inc., New York, 1979. (b) D. N. Reinhoudt, Adv. Heterocycl. Chem., 1977, 21, 253.
5. Takeda Chemical Industries, Ltd., Fr. Pat. 1,400,843; Chem. Abst., 1965, 63, 9922.
6. R. M. Acheson, M. W. Foxton and G. R. Miller, J. Chem. Soc., 1965, 3200.
7. (a) P. J. Abbott, R. M. Acheson, U. Eisner, D. J. Watkin and J. R. Carruthers, J. C. S. Perkin I, 1976, 1269. (b) D. H. Reid, F. S. Skelton and W. Bonthron, Tetrahedron Letters, 1964, 1797.
8. D. S. Tartell, H. P. Hirschler and R. B. Carlin, J. Am. Chem. Soc., 1950, 72, 3138; see also Ref. 3.
9. J. E. Baldwin, J. C. S. Chem. Comm., 1976, 738.
10. Physical and chemical data for compound 14: Rf 0.60 (20% acetone/hexane, silica gel); Mass spectrum, M^+ m/e 311; IR($CHCl_3$) 1690, 1605 cm^{-1} , no hydroxyl or acetylenic absorptions; 1H NMR($CDCl_3$) δ 0.95 (d, 3H, J = 6 Hz, collapsing to a singlet upon irradiation at 1.96), 1.28 (t, 3H, J = 7 Hz, collapsing to a singlet upon irradiation at 3.86), 1.46 - 1.80 (br m, 2H), 1.96 (m, 1H, H-8), 2.12 (s, 3H), 2.24 (m, 2H, H-9's), 3.66 (s, 3H, -OMe), 3.86 (q, 2H, J = 7 Hz, collapsing to a singlet upon irradiation at 1.28), 5.29 (d, 1H, J = 0.5 Hz, collapsing to a sharp singlet upon irradiation at 5.91, H-15), 5.91 (br s, 1H, collapsing to a doublet, J = 3 Hz, upon irradiation at 1.74 and sharpening upon irradiation at 1.60, H-6); ^{13}C NMR($CDCl_3$) 11.8 (q, C-14), 14.9 (q, C-13), 20.8 (q, C-11), 25.1 (d, C-8), 33.5 (t, C-7), 46.6 (t, C-9), 50.7 (q, C-17), 69.5 (t, C-12), 81.4 (d, C-6), 84.6 (d, C-15), 110.3 (s, C-10), 114.7 (s, C-3), 137.5 (s, C-2), 159.2 (s, C-5), 167.9 (s, C-16). The analytical sample crystallized from ether/pet ether in the form of colorless needles, mp 90 - 92° C, which, however rapidly decomposed at ambient temperatures and were unsuitable for X-ray analysis: Anal. calcd. for $C_{15}H_{21}NO_4S$: C, 57.84, H, 6.79, N, 4.52; Found: C, 57.59, H, 6.87, N, 4.23.
11. D. J. Hart, J. Org. Chem., 1981, 46, 367.
12. (a) J. Steele and R. J. Stoodley, J. C. S. Chem. Comm., 1982, 1215. (b) R. M. Silverstein, G. C. Bassler and T. C. Morrill, "Spectrometric Identification of Organic Compounds", John Wiley and Sons, New York, 1981.
13. A. Medici, P. Pedrini and M. Fogagnolo, J. C. S. Chem. Comm., 1980, 1077.
14. Y. Kishi, M. Aratani, H. Tanino, T. Fukuyama, T. Goto, S. Inoue, S. Sugiura and H. Kakoi, J. C. S. Chem. Comm., 1972, 64.
15. (a) R. H. Schlessinger and A. Lopes, J. Org. Chem., 1981, 46, 5252. (b) D. F. Taber, C. Campbell, B. P. Gunn and I. -C. Chiu, Tetrahedron Letters, 1981, 5141.

16. Physical and chemical data for compounds 11d and 13d: (a) 11d: pale yellow oil, Rf 0.45 (30% acetone/hexane, silica gel); Mass spectrum, M^+ m/e 295; IR(CHCl₃) 2205, 1715, 1565 cm⁻¹; ¹H NMR(CDCl₃) δ 1.01 (d, 3H, J = 6 Hz), 1.35 (t, 3H, J = 7 Hz), 2.20 (s, 3H), 2.57 (m, 3H), 2.76 (m, 2H), 3.39 (s, 3H), 4.02 (q, 2H, J = 7 Hz), 4.25 (s, 2H); Anal. calcd for C₁₅H₂₁NO₃S: C, 60.99, H, 7.17, N, 4.74, S, 10.85; Found: C, 60.70, H, 7.40, N, 4.44, S, 10.60. (b) 13d: colorless oil, Rf 0.50 (30% acetone/hexane, silica gel); Mass spectrum, M^+ m/e 254; IR(CHCl₃) 1660, 1555, 1500 cm⁻¹; ¹H NMR(CDCl₃) δ 1.14 (d, 3H, J = 6 Hz, collapsing to a singlet upon irradiation at 2.42, Me-6), 1.42 (t, 3H, J = 7 Hz, collapsing to a singlet upon irradiation at 4.13, -OCH₂CH₃), 2.30 (dd, 1H, J = 14, 12Hz, collapsing to a doublet, J = 12 Hz, upon irradiation at 2.58, H-7), 2.42 (m, 1H, sharpens upon irradiation at 1.14, H-6), 2.58 (m, 2H, simplifies upon irradiation at either 2.30 or 2.95, H-5' and H-7'), 2.95 (dd, 1H, J = 15, 3 Hz, collapsing to a doublet, J = 3 Hz, upon irradiation at 2.58, H-5), 3.41 (s, 3H, -CH₂OCH₃), 4.13 (q, 2H, J = 7 Hz, collapsing to a singlet upon irradiation at 1.42, -OCH₂CH₃), 4.60 (s, 2H, -CH₂OCH₃). Anal. Calcd. for C₁₃H₁₈O₃S: C, 61.39, H, 7.13; Found: C, 61.60, H, 7.25.
17. I. A. Odeh, MA thesis, Department of Chemistry, Wesleyan University, Middletown, Connecticut, 06457, 1981. See also Ref. 1b.

Received, 30th November, 1983