


THE STRUCTURE OF GRACILINE, A NEW DITERPENOID ALKALOID FROM
DELPHINIUM GRACILE

Antonio G. González,* Gabriel de la Fuente, Matías Reina, and
Isabel Timón


Instituto de Productos Naturales Orgánicos, CSIC, La Laguna;
Instituto Universitario de Química Orgánica, Universidad de
La Laguna, Tenerife, Canary Islands, Spain

Abstract - Besides hetisinone, 13-acetylhetisinone, cardio-
petamine, and atisinium chloride, a new diterpenoid alkaloid
graciline was isolated from Delphinium gracile, and its
structure elucidated. The ^{13}C -NMR chemical shifts for graciline,
cardiopetaline and cardiopetalidine are presented.

We report here the structure elucidation of graciline (**1**), a new lycoctonine-
type diterpenoid alkaloid, isolated from Delphinium gracile DC., collected in
Cadiz, Spain.



1 R=H
2 R=Ac



3 R=H R₁=OCH₃
4 R=R₁=
5 R=OH R₁=H

Graciline had mp 98-100°C. Its high resolution MS was characteristic of alkaloids
with lycoctonine skeleton¹: M⁺ 361.2258 amu (3%), C₂₁H₃₁NO₄ (calc. 361.2253), M⁺-CH₃
(14%), M⁺-C₃H₄O (100%), and |M⁺-C₃H₄O|-CH₃ (13%). The ^1H -NMR (CDCl₃) gave signals
at δ 0.89 (3H, s, CH₃), 1.11 (3H, t, J = 7Hz, N-CH₂-CH₃), and 4.07 (1H, t, J = 4.5Hz,
C-14 β H). The loss of a molecule of acrolein in the MS¹, the IR absorptions at 978
and 875 cm⁻¹², and the ^1H -NMR signals at δ 3.73 (1H, m, W_{1/2} = 7Hz, C-1 β H) and 3.85

(1H, s, C-19H), pointed out the presence of the C-1-C-19 inner ether in the new base. Treatment of graciline (1) with acetic anhydride in pyridine afforded a monoacetate (2) as a resin, M^+ 403. IR (CHCl_3), 1725 and 1255 cm^{-1} (acetate). Its $^1\text{H-NMR}$ spectrum exhibited a one-proton triplet centered at δ 4.76 ($J = 4.5\text{Hz}$), indicating that the C-14 α OH group was acetylated.

TABLE 1

Carbon	1	3	4	5	Carbon	1	3	4	5
1	89.3	72.4	72.3	72.7	12	30.6	29.3	32.8	32.0
2	28.9	29.6	29.7	29.6	13	35.5	44.1	34.9	34.8
3	22.8	31.3	31.3	31.9	14	74.6	75.6	76.3	75.8
4	38.1	32.9	32.9	33.5	15	27.5	42.5	32.6	26.7
5	36.4	45.1	46.3	47.5	16	23.0	82.4	25.1	24.9
6	33.1	25.2	25.6	34.0	17	64.5	63.0	63.0	64.0
7	87.1	46.7	46.7	87.2	18	19.8	27.6	27.6	27.4
8	76.2	74.3	77.2	78.4	19	68.7	60.3	60.4	59.4
9	46.8	46.7	46.9	48.0	20	48.0	48.3	48.3	50.5
10	45.3	40.4	44.1	43.8	21	14.3	13.0	13.0	13.5
11	48.0	48.9	49.0	50.2	16'		56.2		

^{13}C Chemical shifts in ppm downfield from TMS and assignments using deuteriochloroform as solvent.

The $^{13}\text{C-NMR}$ spectrum of graciline (1) showed twenty signals for twenty-one carbon atoms in the molecule, and the assignments were made by comparison with the spectra of karakoline (3)³, cardiopetaline (4), cardiopetalidine (5)⁴, and other published $^{13}\text{C-NMR}$ data for related diterpenoid alkaloids. The chemical shift pattern in graciline (1) was very close to that of cardiopetalidine (5). As in the case of 18-methoxygadesine⁵, the appearance of a new doublet at 68.7 ppm, the disappearance of a triplet at \sim 60 ppm, the downfield shift of C-1, and the β or γ effects observed on the resonances of C-3, C-4, C-5 and C-18 in the spectrum of graciline (1) with respect to that of the other alkaloids considered, afforded evidence for the presence of a C-1-C-19 inner ether in (1).

On the other hand, it is interesting to note the α and β substituent effect of \sim 58 ppm and \sim 9 ppm, respectively, produced upon replacement of C-16 β H by OCH_3 ,

and the upfield shift of C-15 by ~ 5.5 ppm in graciline (1) and cardiopetalidine (5) with respect to cardiopetaline (4) due to the γ effect of C-7OH.

The oxidation of cardiopetalidine (5) with KMnO_4 ⁶ led to graciline (1) in 83% yield (mp, IR, ¹H-NMR and MS identical), and therefore its structure was definitively established.

We have also isolated hetisinone⁷, 13-acetylhetisinone⁸, cardiopetamine⁹, and atisinium chloride¹⁰ from this plant, all of which were identified by comparison with authentic samples.

ACKNOWLEDGEMENT This work was partially financed by a grant from the Comisión Asesora de Investigación Científica y Técnica of Spain.

REFERENCES

1. S.W. Pelletier and S.W. Page, "Chemistry of Diterpene Alkaloids" in "The Alkaloids", ed. by J.E. Saxton, The Chem. Soc., London, 1973, vol. 3, p. 235.
2. R. Auct, D. Clayton, and L. Marion, Can J. Chem., 1957, 35, 397.
3. Yang Tsung-re, Hao Xiao-jiang, and Chow Jun, Acta Botanica Yunnanica, 1979 1, 41.
4. A.G. González, G. de la Fuente, M. Reina, V. Zabel, and W.H. Watson, Tetrahedron Letters, 1980, 21, 1155.
5. A.G. González, G. de la Fuente, and O. Munguía, Heterocycles, 1983, 20, 409.
6. O. Achmatowicz, Y. Tsuda, and L. Marion, Can. J. Chem., 1965, 43, 2336.
7. R.T. Aplin, M.H. Benn, S.W. Pelletier, J. Solo, S.A. Telang, and H. Wright, Can. J. Chem., 1968, 46, 2635.
8. A.G. González, G. de la Fuente, and M. Reina, Anal. Quim., 1981, 77C, 171.
9. A.G. González, G. de la Fuente, M. Reina, P.G. Jones, and P.R. Raithby, Tetrahedron Letters, 1983, 24, 3765.
10. S.W. Pelletier, R. Aneja, and K.W. Gopinath, Phytochemistry, 1968, 7, 625.

Received, 11th November, 1983