

THE CHEMISTRY OF FUROXANS

Wanda Sliwa and Antoni Thomas^x

Department of Organic Chemistry

Pedagogical University, 42-201 Częstochowa, POLAND

Abstract - Syntheses, chemical reactivity, as well as physical and biological properties of furoxans are presented.

I. INTRODUCTION

The present review, dealing with 1,2,5-oxadiazole-1-oxides, i.e. furoxans is a continuation of our former papers concerning oxa-, thia- and selenadiazoles¹⁻⁴.


The chemistry of furoxans is intensively developing⁵⁻⁷; because of the big amount of papers concerning benzofuroxans, useful synthons of biologically active quinoxaline-di-N-oxides^{8,9}, this topic is not included here.

Presenting syntheses and properties of furoxans, we divide them into two groups - symmetrically and unsymmetrically 3,4-disubstituted.

II. SYNTHESSES

SYMMETRICALLY 3,4-DISUBSTITUTED FUROXANS

Among numerous syntheses of furoxans there ought to be mentioned chemical oxidation reactions of dioximes¹⁰⁻¹³; the anodic oxidation being a convenient modification of this method¹⁴.




R = Me 33%

R = C₆H₅ 68%

^x Deceased



The following procedure also results in furoxans¹⁵:




R = NHR¹, R¹ = C₁₋₆ alkyl, C₄₋₇ cycloalkyl

R = NR²R³, R², R³ = C₁₋₄ alkyl, and other


The furoxan ring can be formed in the reaction of terminal acetylenes with NOF¹⁶:




1
35%



2
39%



3
44%




4
78%

The presence of α -acetoxy group in the starting acetylene is necessary for the reaction to proceed. The process begins with the attack of nitronium ion, present in the reaction mixture, on the acetylenic moiety. The dimerisation of the formed nitroacetylene and the subsequent rearrangement gives way to 1 - 4.

Structures of 1 - 4 can be confirmed by their thermal reversion into two molecules of β -acetoxy- α -oxonitrile oxide, reacting as 1,3-dipole with dipolarophiles¹⁷ /described in the following part, dealing with the chemical reactivity of furoxans/.

For the synthesis of isoxazoles and dihydroisoxazoles often the cycloaddition of nitrile oxides to substituted alkenes or alkynes is used. In this procedure, as the side reaction, the dimerisation of $\text{RCN}^{\oplus}-\text{O}^{\ominus}$, resulting in furoxan, takes place; for this purpose the use of certain nitrile oxides in cycloaddition reactions is limited¹⁸.


For instance, in the following reaction, instead of the expected 1,3-cycloadduct, only 3,4-diphenylfuroxan could be obtained¹⁹.




UNSYMMETRICALLY 3,4-DISUBSTITUTED FUROXANS


In the oxidation reaction of dioximes, the structure of the resulting furoxans depends on the configuration of dioximes, as well as on the nature of the oxidizing agent²⁰.

The oxidation of antichloromethylglyoxime 5 with dinitrogen tetroxide gives way to the mixture of two isomers, 6 and 7, in the ratio of 2,4 : 1²¹.




However, when ceric ion was used as the oxidizing agent, the major product was 7²¹. This fact can be explained by the following oxidation mechanism, where the attack of Ce^{4+} on the oxime group adjacent to the methyl is favoured.







The oxidation reaction of anti-chlorophenylglyoxime results in the mixture of two isomeric furoxans 8 and 9, undergoing thermal isomerisation at the temperature of 100°C ²².




The mixture of syn- and anti-oximes 10, formed in the reaction of sorbic acid with sodium nitrite yields furoxan 11 in the following procedure²³:



Among bicyclic furoxans, trimethylenefuroxan ought to be mentioned; this compound is obtained by oximation of cyclopentene-1,2-dione and subsequent treatment of dioxime with sodium hypochlorite²⁴:




The strained furoxans 12 - 14 can be synthesized from nitrooximes 15 and 16 in the following procedure²⁵:



In the case of 16, the mixture of isomers 13 and 14, decomposing explosively when heated to 80°C , is formed.

Pyrido[2,3-c]furoxan 17 can be obtained in following procedures, the aim of these reactions being the use of 17 as the synthon of pyridopyrazine-1,4-dioxides, in the analogy to benzofuroxans^{26,27}.




III. CHEMICAL REACTIVITY

SYMMETRICALLY 3,4-DISUBSTITUTED FUROXANS


In numerous studies of the thermolysis of alkyl- and arylfuroxans there was found, that at the lower temperature range /100 to 300°C , depending on the substituent/, the ring cleavage leading to nitrile oxides, takes place.

Nitrile oxides in the presence of dipolarophiles give 1,3-dipolar cycloadducts and in the absence of dipolarophiles rearrange to isocyanates²⁸.


However at very high temperatures /500 - 600°C , under flash vacuum pyrolysis conditions/ the formed nitrile oxides can be isolated and characterized¹⁸.




Furoxans generally are readily available, therefore their use as nitrile oxide precursors is of interest in the synthesis of heterocycles. In this regard the reaction of 18 with a series of dipolarophiles, giving rise to isoxarolines, was studied¹⁸.




In a similar way, fused furoxans heated with dipolarophiles yield bicycloadducts, for instance²⁴:



Readily accessible bicyclic furoxans, e.g. 19 give rise to bis-nitrile oxides and di-isocyanates, difficult to obtain on other routes. Thus, thermolysis of 19 in the presence of tetradec-1-ene and decanonitrile, used as dipolarophiles, resulted in cycloadducts 21 and 22, respectively²⁸.




Thermolysis of **19** in the absence of dipolarophiles, with hexadecane yielded diisocyanate **23**, and with decan-1-ol **23** was isolated as the carbamate **24**²⁸.




Similar results were obtained in the case of 3,4-diphenylfuroxan²⁸.

When furoxans are thermolyzed under flash vacuum pyrolysis conditions nitrile oxides can be isolated and identified, for instance dianisylfuroxan gives way to anisonitrile oxide²⁹:




On this way 3,4-dimethylfuroxan yields the unstable acetonitrile oxide.




So, the FVP technique permits the spectroscopic examination and identification of shortlived nitrile oxides, and on the other hand, provides a convenient synthetic route from furoxans to isoxazolines.

The following cycloadducts of nitrile oxides, resulting in the FVP of furoxans with alkenes, were obtained²⁹:




R	R ¹	yield(%)
4-MeOC ₆ H ₄	Bu	75
Me	Bu	79
Et	Bu	95
C ₆ H ₅	Bu	97
4-MeOC ₆ H ₄	C ₁₂ H ₂₅	86


Among other cycloaddition reactions of nitrile oxides, formed in situ from furoxans one can mention the thermolysis of 25 in the presence of dimethyl maleate^{30,31}:



The by-product 27 results from the dimerisation of the oxirane 28, formed by the decomposition of cycloadduct 26 taking place under the reaction conditions.




Also diacyl furoxans may serve as the source of nitrile oxides, trapped as 1,3-cycloadducts¹⁷:




The reaction with nitroenamine provides a convenient route in the synthesis of 4-nitroisoxazoles¹⁷.

However, in the treatment of 3,4-dibenzoylfuroxan 29 with DMAD, instead of the expected isoxazole 30 the adduct 31 is formed¹⁷.




The mechanism of this reaction can be presented as follows:




Furoxan 29 is in the thermal equilibrium with the ring - opened isomer 32. The intramolecular transfer of the benzoyl group in 32 gives rise to 33, which undergoes cycloaddition with dipolarophiles to yield 31.

Other mechanism, a nitrene - like addition is also possible³², however a proof for the described mechanism was obtained by heating 29 in toluene in the absence of dipolarophile. Under these conditions 29 gives 34, resulting from the dimerisation of the nitrile oxide 33¹⁷.



Furoxans can be considered as cyclic 1,3-dipolar nitrones, and as the first example of the nitrono - type 1,3-dipolar cycloaddition of these species the reaction of diethyl furoxandicarboxylate with olefins, giving way to bicyclic compounds 35, was reported ³³. In the first step the 1,3-dipolar cycloadduct 36 is formed, which eliminates ethyl cyanoformate by a retro-1,3-dipolar cycloaddition to give 37; the next 1,3-dipolar cycloaddition of 37 with another molecule of olefin results in the end product 35. The stereochemistry of this reaction is described.





R
 n-C₆H₁₃
 n-C₇H₁₅
 n-C₈H₁₇
 n-C₁₀H₁₇
 CH₂-C₆H₅ and other

Furoxans treated with reducing agents, e.g. triethyl phosphite, can be converted into the appropriate furazans ^{13,34}.


3,4-Diphenylfuroxan was found to be inert towards nucleophiles; the reaction with piperidine, even under reflux, did not take place ^{35,36}.

In the investigation of 3,4-dicyanofuroxan 38, the explosive properties of its mixtures with hydrazine have been observed. The reaction of 38 with hydrazine and hydroxylamine gives way to 39 and 40, respectively ³⁷.







Moderate heat converted the oxazino-furoxan 40 into 42, via the tautomerisation of 40 to 41 and the subsequent elimination of imidogen ³⁷.




Pyrimidofuroxans undergo nucleophilic addition at C₇, for instance ³⁸:



In the study of aminative ring opening reactions there was observed that pyrimidofuroxan 43 treated with primary or secondary amines yields aminonitrosopyrimidines via furoxan ring opening ³⁹:




On the other hand, in such reactions of 44 the pyrimidine ring opening takes place ³⁹:






UNSYMMETRICALLY 3,4-DISUBSTITUTED FUROXANS


Unsymmetrical diarylfuroxans undergo oxidation with ozone to give aromatic acids⁴⁰.




Studying the oxidation reaction of isomeric 45 and 46, the following results were obtained: in the oxidation of 45, besides the expected 47 also 48, 49 and 50 were formed, while in the case of 46 the sole product was 51^{41,42}.



Furoxans 52 and 53 treated with sodium methoxide afford 54 and 55, respectively. The thermal isomerisation of 54 to 55 can be performed by its refluxing in benzene⁴².



The reactivity of furoxans has been largely investigated having in view their biological activity. For instance, the thermolysis of 56 resulted in its isomerisation, and the aminolysis of 57 gave rise to 58⁴³:




IV. PHYSICAL PROPERTIES

The ¹³C NMR spectroscopy is often used in the structure elucidation, e.g. in the case of 3,4-dimethylfuroxan⁴⁴ or norbornene furoxans²⁵. This technique is also very helpful in the identification of isomeric pairs of furoxans:

	Z	Y	ref.
	Me, Et	H	29
	C ₆ H ₅	H	20
	Me	Cl	21
	C ₆ H ₅	Cl	22
	C ₆ H ₅	NO ₂	41

Numerous papers are dealing with X-ray crystallographic analysis of furoxans:


	Z	R	ref.
	NO ₂	H	45
	HOOC-C _H (H)-C _H (H)-	H	23
	C ₆ H ₅	NO ₂	42

as well as of bicyclic systems:


		ref.
	n = 3	24
	n = 4	46
	n = 6	47

and of norbornene furoxans ²⁵.

Among investigations of isomeric pairs of furoxans by X-ray crystallographic analysis the following ones ought to be presented:


	Z	ref.
	-C(=O)-NH ₂	48
	-C(=O)-NMe ₂	49,50
	-NH-C(=O)-O-CHMe ₂	51
	-SO ₂ C ₆ H ₅	52

Some furoxans have explosive properties, e.g. dicyanofuroxans ^{37,53}, potassium salt 59⁵⁴, trimethylene furoxan²⁴ and dicyclopentadiene furoxan ²⁵.




V BIOLOGICAL ACTIVITY


Among biologically active furoxans the following ones show antihypertensive properties ^{15,34,43}:



R = OH, alkoxy, amino



R = C₁₋₄ alkyl, C₅₋₇ cycloalkyl



Z

(CH₂)₄

(CH₂)₅

(CH₂)₂O(CH₂)₂

(CH₂)₂NMe(CH₂)₂

and other

REFERENCES

1. W. Sliwa and A. Thomas, Wiad. Chem., 1981, 35, 373.
2. A. Thomas and W. Sliwa, Heterocycles, 1983, 20, 1043.
3. W. Sliwa and A. Thomas, Heterocycles, 1983, 20, 71.
4. W. Sliwa, Heterocycles, 1984, 22, 1571.
5. A. Gasco and A.J. Boulton, Adv. Heterocyclic Chem., 1981, 251.
6. L.I. Klmelnitskii, S.S. Novikov and T.L. Godovikova, Khimiya furoxanov, Nauka, Moskva, 1983.
7. R.M. Paton, Comprehensive Heterocyclic Chemistry, Vol. 6, A.R. Katritzky, Ch. W. Rees, Ed., Pergamon Press, 1984.
8. P. Benko, D. Bozsing and I. Farkas, Hung. Teljes HU 23.243 /1982/; Chem. Abstr., 1983, 98, 198273f.
9. C.H. Issidorides and M.J. Haddadin, US 4.343.942 /1982/; Chem. Abstr., 1983, 98, 4563g.

10. K.L. Stuart, Heterocycles, 1975, 3, 651.
11. I. Tabakovič, M. Trkovnik and D. Galijaš, J. Electroanalyt. Chem. /Amsterdam/, 1978, 86, 241.
12. S. Spyroudis and A. Varvoglis, Synthesis, 1975, 445.
13. D.N. Nicolaides, K.E. Litinas and I. Naidou, Chem. Chron., 1982, 11, 73; Chem. Abstr., 1983, 98, 89268n.
14. N. Henning, Th. Dassler and W. Jugelt, Zeitschr. f. Chemie, 1982, 22, 25.
15. K. Schoenafinger, R. Beyerle, A. Mogilev, H. Bohn, M. Just, P.A. Martorana and R.E. Nitz, Eur. Pat. Appl. EF 54.872 /1982/; Chem. Abstr., 1982, 97, 216189u.
16. D.R. Brittelli and G.A. Boswell, Jr., J. Org. Chem., 1981, 46, 312.
17. D.R. Brittelli and G.A. Boswell, Jr., J. Org. Chem., 1981, 46, 316.
18. R.A. Whitney and E.S. Nicholas, Tetrahedron Letters, 1981, 22, 3371.
19. W.P. Iwchin, W.F. Smirnov and A.G. Chakryriyanova, Zh. Org. Khim., 1983, 19, 1319.
20. L.I. Khmel'nitskii, T.I. Godovikova, N.A. Rulyeva, B.N. Hasanov and S.S. Novikov, Izv. Akad. Nauk SSSR, Ser. Khim., 1979, 2295.
21. R. Calvino, A. Gasco, E. Menziani and A. Serafino, J. Heterocyclic Chem., 1983, 20, 783.
22. R. Calvino, R. Fruttero, A. Gasco, V. Mortarini and S. Aime, J. Heterocyclic Chem., 1982, 19, 427.
23. T. Osawa, Y. Kito, H. Namiki and K. Tsuji, Tetrahedron Letters, 1979, 4399.
24. J.F. Barnes, M.J. Barrow, M.M. Harding, R.M. Paton, P.L. Ashcroft, J. Crosby and C.J. Joyce, J. Chem. Res. /S/, 1979, 314.
25. J.F. Barnes, M.J. Barrow, M.M. Harding, R.M. Paton, A. Sillitoe, P.L. Ashcroft, R. Bradbury, J. Crosby, C.J. Joyce, D.R. Holmes and J. Milner, J. Chem. Soc., Perkin Trans. I, 1983, 293.
26. D. Binder, C. R. Noe, J. Nussbaumer and B.C. Prager, Monatsh. Chem., 1980, 111, 407.
27. D. Binder, A. Georgopoulos, C.R. Noe, J. Nussbaumer, B.C. Prager and F. Turnovsky, Arzneim. Forsch., 1982, 32, 10.
28. J.A. Chapman, J. Crosby, C.A. Cummings, R.A.C. Rennie and R.M. Paton, J. Chem. Soc., Chem. Commun., 1976, 240.

29. W.R. Mitchell and R.M. Paton, Tetrahedron Letters, 1979, 2443.
30. G. Westphal and G. Karge, Zeitschr. f. Chemie, 1982, 22, 138.
31. G. Westphal and M. Karge, Zeitschr. f. Chemie, 1982, 22, 217.
32. M. Altaf-ur-Rahman, A.J. Boulton and D. Middleton, Tetrahedron Letters, 1972, 3469.
33. T. Shimizu, Y. Hayashi and K. Teramura, J. Org. Chem., 1983, 48, 3053.
34. K. Schoenafinger, R. Beyerle, P. Martorana and R.E. Nitz, Ger. Offen. DE 3.134.849 /1983/; Chem. Abstr., 1983, 98, 198249c.
35. W.S. Latham, O. Meth-Cohn and H. Suschitzky, J. Chem. Soc., Perkin Trans. I, 1976, 2216.
36. N.A. Muffary, M.J. Haddadin, C.H. Issidorides, J.W. Mc Farland and J.D. Johnston, J. Chem. Soc., Perkin Trans. I, 1972, 965.
37. J.H. Boyer and T.P. Pillai, Heterocycles, 1982, 19, 1063.
38. G. Tennant and Ch.W. Yacomeni, J. Chem. Soc., Chem. Commun., 1982, 60.
39. G. Tennant and G.M. Wallace, J. Chem. Soc., Chem. Commun., 1982, 267.
40. J.H. Boyer and Ch. Huang, Pol. J. Chem., 1981, 55, 1233.
41. N.N. Makhova, I.W. Owchinnikov, B.N. Hasanov and L.I. Khmel'nitskii, Izv. Akad. Nauk SSSR, Ser. Khim., 1982, 646.
42. R. Calvino, A. Gasco, A. Serafino and D. Viterbo, J. Chem. Soc., Perkin Trans. II, 1981, 1240.
43. K. Schoenafinger, R. Beyerle, A. Mogilev, H. Bohn, M. Just, P.A. Martorana and R.E. Nitz, Eur. Pat. Appl. EP 54.873 /1982/; Chem. Abstr., 1983, 98, 107296b.
44. F.A.L. Anet and I. Yavari, Org. Magn. Resonance, 1976, 8, 158.
45. A.F. Cameron and A.A. Freer, Acta Cryst., 1974, B30, 354.
46. M.J. Barrow, Acta Cryst., 1982, B38, 308.
47. M.M. Harding and R.M. Paton, Acta Cryst., 1982, B38, 1395.
48. M. Calleri, G. Chiari, A. Chiesi Villa, A. Gaetani Manfredotti, C. Guastini and D. Viterbo, Acta Cryst., 1975, B31, 2384.
49. M. Calleri, D. Viterbo, A. Gaetani Manfredotti and C. Guastini, Cryst. Struct. Commun., 1974, 3, 269.
50. A. Chiesi Villa, C. Guastini, M. Calleri and G. Chiari, Cryst. Struct. Commun., 1974, 3, 265.

51. M. Calleri, G. Chiari, A. Chiesi Villa, A. Gaetani Manfredotti, C. Guastini and D. Viterbo, Acta Cryst., 1977, B33, 479.
52. A. Chiesi Villa, A. Gaetani Manfredotti, C. Guastini and D. Viterbo, Acta Cryst., 1976, B32, 1032.
53. R.H. Homewood, V.J. Krukonis and R.C. Loszewski, US 3.832.249 /1974/; Chem. Abstr., 1975, 82, 113795z.
54. A. Gasco, V. Mortarini, R. Calvino and A. Serafino, Tetrahedron Letters, 1974, 627.

Received, 31st July, 1984