

AROMATIC VERSIONS OF CLAISEN REARRANGEMENT OF LACTONIC SILYL  
ENOLATES — A NOVEL APPROACH TO FURANOSESQUITERPENE FRAMEWORK

Hideo Nemoto<sup>a</sup>, Eiki Shitara<sup>a</sup>, Keiichiro Fukumoto\*<sup>a</sup>, and Tetsuji Kametani<sup>b</sup>

a Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980, Japan


b Institute of Medicinal Chemistry, Hoshi University, Ebara 2-4-41, Shinagawa-ku, Tokyo 142, Japan

**Abstract** — A new route to the decalin systems (14) and (15) is described. The key step for this synthesis is the Claisen rearrangement of furfuryl lactonic silyl enolates (8) and (9) derived from the furfuryl lactones (6) and (7), respectively.

In the preceding paper<sup>1</sup>, we described the aromatic version of an Ireland's allylic ester silyl enolate rearrangement and could show that the rearrangement occurred under remarkably mild conditions. The most attractive feature of the Ireland's rearrangement seems that a relatively accessible acyl oxygen bond is used to generate a more difficultly available carbon-carbon bond. Especially in the case of lactone, cycloalkenes are readily available<sup>2,3</sup>. If this could be the case for our aromatic version of the Ireland's rearrangement, functionalized furanocycloalkenes could be generated. On the other hand, the decalin systems fused with a furan ring are an important basic skeleton of eudesmane or eremophilane types of sesquiterpenes<sup>4</sup> and numerous investigations directed toward a synthesis of this type of compounds have been cumulated up to this time<sup>5,6</sup>. In these concepts, we have undertaken the investigation to explore a new synthetic entry to furanodecalin systems.

Firstly, the furfuryl lactones (6) and (7), key intermediates for generating furfuryl lactonic silyl enolates (8) and (9), were prepared as follows. The ketonic ester (1) [ $m/z$  184 ( $M^+$ )], prepared in 60 % yield by successive treatment (pyrrolidine, benzene, reflux, 12 h; methyl acrylate, dioxane, reflux, 1.5 h) of cyclohexanone, reacted with 2-lithio-5-methylfuran<sup>7</sup> (THF,  $-78^\circ\text{C}$ , 2 h) to give the furfuryl ester (2) and then the furfuryl carboxylic acid (3), obtained by the hydrolysis (LiOH, THF- $\text{H}_2\text{O}$ , room temperature, 12 h) of 2, was lactonized (silica gel,  $\text{CH}_2\text{Cl}_2$ , room temperature, 12 h) to furnish the furfuryl lactone (6) as a stereoisomeric mixture [ $m/e$  234 ( $M^+$ );  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ ) 1730  $\text{cm}^{-1}$ ] in 35 % overall yield based on 1. The furfuryl lactone (7) [ $m/e$  220 ( $M^+$ );  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ ) 1720  $\text{cm}^{-1}$ ] was also prepared in 58 % overall yield starting from the ketonic ester (1) by using 2-lithiofuran<sup>8</sup>, instead of 2-lithio-5-methylfuran under the same conditions described for the synthesis of the furfuryl lactone (6), via the furfuryl ester (4) and the furfuryl carboxylic acid (5). Silylation (TBSCl, LDA, HMPA-THF,  $-78^\circ\text{C}$ , 1 h) of 6 and 7 gave the furfuryl lactonic silyl enolates (8) and (9), respective-

ly, which were subjected to the rearrangement (toluene, reflux, 12 h) and then the hydrolysis (10 % HCl, THF, room temperature, 5 min) of the resulting silyl esters (12) and (13) to afford the carboxylic acids (14) [ $m/z$  234 ( $M^+$ ); max ( $\text{CHCl}_3$ )  $1710\text{ cm}^{-1}$ ; ( $\text{CCl}_4$ ) 2.16, 2.20 (3H, each s, Me) and 5.77, 5.83 (1H, each br s, ArH)] and (15) [ $m/z$  220 ( $M^+$ );  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ )  $1720\text{ cm}^{-1}$ ;  $\delta$  ( $\text{CCl}_4$ ) 6.26 (1 H, m, ArH) and 7.06 (1 H, m, ArH)] in 28 % and 55 % overall yields, respectively as stereoisomeric mixtures. In these reactions, it is obvious that [3.3] sigmatropic rearrangement and hydrogen shift were occurred successively giving firstly 10 and 11 and then aromatized compounds (12) and (13) respectively. Finally, the decarboxylation of 15 was conducted under Mukaiyama's conditions<sup>9</sup> [2-dipyridyl disulfide),  $\text{Ph}_3\text{P}$ ,  $\text{CH}_3\text{CN}$ , room temperature, 0.5 h; Zn, cat.  $\text{NiCl}_2$ , DMF,  $100^\circ\text{C}$ , 1 h] to afford the tetrahydronaphthofuran (18) [ $m/z$  172 ( $M^+$ );  $\delta$  ( $\text{CCl}_4$ ) 6.55 (1H, d,  $J=2$  Hz, ArH), 6.76(1H, d,  $J=8$  Hz ArH), 7.16(1H, d,  $J=8$  Hz ArH), and 7.42(H, d,  $J=2$  Hz, ArH). UV<sub>max</sub>  $\lambda$  246, 276 and 285 nm]<sup>10</sup> in 54% yield, which was derived by the oxidation of the initially expected compound (17) via the thioester (16).



Thus, we could disclose a new route for the synthesis of furanosesquiterpene framework, although the stereochemistry of the synthetic intermediates remained unknown.

## REFERENCES AND NOTES

1. H. Nemoto, E. Shitara, K. Fukumoto, and T. Kametani, Heterocycles, 1985, **23**, 549.
2. S. Danishefsky, R. L. Funk, and J. F. Kerwin, Jr. J. Am. Chem. Soc., 1980, **102**, 6881.
3. S. Danishefsky and K. Tsuzuki, J. Am. Chem. Soc., 1980, **102**, 6891.
4. T. K. Devon and A. I. Sott, "Handbook of Naturally Occurring Compounds," Vol. II. Terpenes, Academic Press, New York, 1972.
5. "The Total Synthesis of Natural Products", Ed. by J. Apsimon, Vol. 2, John Wiley & Sons, New York, 1973.
6. "Terpenoids and Steroids", Ed. by J. R. Hanson (Specialist Periodical Reports), The Royal Society of Chemistry, London, Vol. 1-12.
7. G. Büchi and H. Wüest, J. Org. Chem., 1966, **31**, 977.
8. V. Ramanathan and R. Levine, J. Org. Chem., 1962, **27**, 1216.
9. M. Araki, S. Sakata, H. Takei, and T. Mukaiyama, Bull. Chem. Soc. Jpn., 1974, **47**, 1977, T. Goto, M. Onaka, and T. Mukaiyama, Chem. Lett., 1980, 709.
10. Characteristic in benzofuran chromophore; A. I. Scott, "International Series of Monographs on Organic Chemistry", Vol. 7, "Interpretation of the Ultraviolet Spectra of Natural Products", Ed. by G. Villars, Pergamon Press, New York, 135(1964).

Received, 15th April, 1985