

THE SYNTHESIS OF 5- AND 7-ACETYLINDOLE DERIVATIVES. I.
THE PHOTOCHEMICAL REARRANGEMENT OF 1-ACETYLINDOLINE

Masao Akagi* and Kazuko Ozaki

Osaka University of Pharmaceutical Sciences,
2-10-65, Kawai, Matsubara, Osaka 580, Japan

Abstract- 5- and 7-Acetyl substituted indole derivatives have been prepared via intramolecular photorearrangement of acyl radical and intermolecular electrophilic acylation of 1-acetylindoline.

The photochemistry of the enamide system and its useful application have been well investigated;¹ simple enamides generally undergo a [1,3]-acyl radical shift to afford vinylogous amides. Meanwhile, photoisomerization of 1-acylindoles of an enamide system have been described to merely provide 2-,3-,4- and 6-acylindoles,² and then neither of 5- and 7-acylindoles formed.

We wish to report here that the photoisomerization of 1-acetylindoline (1) to 5- and 7-acetylindolines or corresponding indoles.³

Irradiation of 1-acetylindoline (1) in acetone with a 60-W low pressure mercury lamp in nitrogen atmosphere for 24 h afforded 5- and 7-acetylindolines (2 and 3, in 12% and 11% yields, respectively), together with their corresponding N-acetates (4 and 5) and indole (6). Formation of 2 and 3 is interpreted in terms of a normal rearrangement in analogy with the photo-Fries rearrangement of acetanilides.^{4,5}

However, 1,5-diacetylindoline (4, in 2.4% yield) and 1,7-diacetylindoline (5, in 2.5% yield) are abnormal products.^{5,6} The structure of these compounds were therefore confirmed by following treatments and spectral data⁷ (Table I and II).





Table I. ^1H -NMR Chemical Shifts (ppm) of 1-5

Compounds	1	2	3	4	5
C(2)- <u>H</u>	4.44(2H) (t, J=9Hz)	4.26(2H) (t, J=9Hz)	4.24(2H) (t, J=9Hz)	4.40(2H) (t, J=9Hz)	4.63(2H) (t, J=9Hz)
C(3)- <u>H</u>	3.46(2H) (t, J=9Hz)	3.58(2H) (t, J=9Hz)	3.54(2H) (t, J=9Hz)	3.45(2H) (t, J=9Hz)	3.60(2H) (t, J=9Hz)
C(4)- <u>H</u>	7.44(3H) 4,5,6-H (m)	8.25(1H) (d, J=2Hz)	7.88(1H) (d, J=9Hz)	8.08(1H) (d, J=2Hz)	7.46(1H) (d, J=9Hz)
C(5)- <u>H</u>		----	7.83(1H) (t, J=9Hz)	----	7.60(2H) 5,6-H(m)
C(6)- <u>H</u>		8.21(1H) (dd, J=9; 2Hz)	8.17(1H) (dd, J=9; 2Hz)	8.10(1H) (dd, J=9; 2Hz)	
C(7)- <u>H</u>	8.10(1H) (d, J=9Hz)	7.75(1H) (d, J=9Hz)	----	8.34(1H) (d, J=9Hz)	----
NCOMe	2.79(s)	----	----	2.61(s)	2.30(s)
CCOMe	----	2.84(s)	2.88(s)	2.80(s)	2.66(s)

Run at 300 MHz in CF_3COOD solution.Table II. ^{13}C -NMR Chemical Shifts (ppm) of 1-5

Compounds	1	2	3	4	5
C(2)	48.70	47.18	46.62	49.18	49.51
C(3)	27.93	28.79	27.90	27.51	29.03
C(4)	124.51	125.79	115.75	124.51	124.99
C(5)	123.53	156.29	115.27	146.96	124.37
C(6)	127.43	127.85	128.14	129.51	126.45
C(7)	116.87	106.98	154.04	116.02	137.52
C(3a)	131.11	128.87	128.50	131.68	131.56
C(7a)	142.89	130.35	131.54	132.74	133.76
NCOMe	168.68	-----	-----	169.33	168.29
NCOMe	24.19	-----	-----	24.31	23.52
CCOMe	-----	196.47	199.09	196.95	200.41
CCOMe	-----	26.13	26.56	26.50	28.83

Run at 75.44 MHz in CDCl_3 solution.

The compounds 2 and 3 were easily converted to 4 and 5 by treatment with acetic anhydride. Hydrolysis of 4 and 5 with methanolic hydrochloride readily gave 2 and 3, respectively. The structure of 2, 3, 4 and 5 were assigned by comparing those uv and nmr spectra with those of the analogous ketoamines (II and III)⁵ and ketoamides (IV and V).⁶

It is generally believed that the photo-Fries rearrangement is an intramolecular reaction.^{8,9} However, occurrence of 4 and 5 in our case clearly indicated that a part of the reaction proceed in an intermolecular fashion. Therefore, the reaction mechanism could be interpreted as follows. Thus, irradiation of 1 produced acetyl and indolyl radicals and subsequent recombination of them in a solvent cage⁸ afforded 2 and 3. Further intermolecular reaction of 2 and 3 with an another acetyl radical, which was generated either by the other molecule (1) or by the Norrish type I fission of acetone used as a solvent, afforded 4 and 5.

Further confirmation of the structures 4 and 5 was carried out by comparing these uv and nmr spectra with those of the following Friedel-Crafts products.

The acylation (the condition, 1 : AlCl₃ : AcCl = 1 : 3 : 3)¹⁰ of 1 afforded 4 and 5 in the ratio of 93 : 7 in 80.8% yield.

In addition, the irradiation of 2 and 3 in acetone solution for 70 h afforded 5-acetylidole (7) and 7-acetylidole (8)¹¹ in 90% and 95% yields, respectively, and then formation of 4 and 5 was not detected by acetyl radical from acetone.

The compounds 2 and 3 were also converted by means of Pd-C dehydrogenation to 7 and 8 in 51% and 76% yields,^{12,13} respectively.

ACKNOWLEDGEMENT

We would like to express our thanks to Miss N.Nabae and Miss K.Takeda for the measurments of nmr and ms spectra, and Mrs. Y.Tsukamoto for microanalyses.

REFERENCES AND NOTES

- 1.G.Lenz, Synthesis, 1978, 489.
- 2.a) S.Naruto and O.Yonemitsu, Tetrahedron Letters, 1971, 2297.
- b) M.Somei and M.Natsume, Tetrahedron Letters, 1973, 27, 2451.
- c) Y.Ban, K.Yoshida, J.Goto and T.Oishi, J.Am.Chem.Soc., 1981, 103, 6990.
- d) Y.Ban, K.Yoshida, J.Goto, T.Oishi and E.Takeda, Tetrahedron, 1983, 39, 3657.
- 3.Presented in part at the 35th Meeting of Kinki Branch, Pharmaceutical Society of Japan, Kyoto, Absts. Papers P.43, 1985.

- 4.a) D.Elad, Tetrahedron Letters, 1971, 873.
 b) H.Kobsa, J.Org.Chem., 1962, 27, 2293.
- 5.The photoisomerization of 2-methylacetanilide (mp 112-114°C, M^+ :m/z 149) afforded 2-methyl-4-acetylaniline (II)(mp 108-109°C, M^+ :m/z 149), which was acylated with acetic anhydride to 2-methyl-4-acetylacetanilide (IV)(mp 153-154°C, M^+ :m/z 191).
- 6.2-Acetyl-4-methylaniline (III)(mp 35°C, M^+ :m/z 149) and 2-acetyl-4-methylacetanilide (V)(mp 123-124°C, M^+ :m/z 191) were formed simultaneously by photochemical rearrangement of 4-methylacetanilide (mp 149-150°C, M^+ :m/z 149) in acetone.
- 7.Compound 1:mp 104-105°C (ether), M^+ :m/z 161 ($C_{10}H_{11}ON$), 188 (base peak), uv λ EtOH/max. 206, 252, 281, 290nm., ir ν Nujol/max. 1645, 1600 cm^{-1} .
 Compound 2:mp 76-77°C (ether), M^+ :m/z 161 ($C_{10}H_{11}ON$), 146 (base peak), 118. uv λ EtOH/max. 206, 244, 337nm., ir ν Nujol/max. 3270, 1635, 1580 cm^{-1} .
 Compound 3:mp 85-86°C (ether), M^+ :m/z 161 ($C_{10}H_{11}ON$) (base peak), 146, 118. uv λ EtOH/max. 234, 259, 380nm., ir ν Nujol/max. 3380, 1630, 1610, 1575 cm^{-1} .
 Compound 4:mp 140-141°C (EtOH), M^+ :m/z 203 ($C_{12}H_{13}O_2N$) (base peak), 188, 161, 146, 118., uv λ EtOH/max. 206, 312nm., ir ν Nujol/max. 1685, 1645, 1590, 1580 cm^{-1} .
 Compound 5:mp 126-127°C (EtOH), M^+ :m/z 203 ($C_{12}H_{13}O_2N$), 161 (base peak), 146, 118. uv λ EtOH/max. 226, 256, 304nm., ir ν Nujol/max. 1680, 1640, 1585 cm^{-1} .
 Compound 6:mp 52-53°C (n-Hexane), M^+ :m/z 117 (C_8H_7N),
 uv (EtOH) characteristic indole chromophore.
- 8.a) H.Shizuka, Bull.Chem.Soc.Japan, 1969, 42, 52, 57.
 b) H.Shizuka, M.Kato, T.Ochiai, K.Matsui and T.Morita, ibid., 1970, 43, 67.
- 9.a) J.C.Anderson and C.B.Reese, Proc.Chem.Soc., 1960, 217.
 b) J.C.Anderson and C.B.Reese, J.Chem.Soc., 1963, 1781.
 c) H.Shizuka and I.Tanaka, Bull.Chem.Soc.Japan, 1969, 41, 2343.
- 10.Y.Murakami, M.Tani, K.Tanaka and Y.Yokoyama, Heterocycles, 1984, 22, 241.
- 11.Indoline was easily dehydrogenated by irradiation in acetone solution to indole.
- 12.Compounds 4 and 5 were not converted to indole derivatives neither by Pd-C dehydrogenation reaction nor by irradiation in acetone solution.
- 13.Compound 7:bp $2_{155}^{\circ}C$, uv λ EtOH/max. 205, 250, 297nm., ir ν film/max. 3300, 1650, 1600 cm^{-1} . 1H -nmr($CDCl_3$) δ (ppm) 2.66(s, CH_3), 7.26(d, J=3 Hz, C_2 -H), 6.38(d, J=3 Hz, C_3 -H), 8.32(d, J=2 Hz, C_4 -H), 7.85(dd, J=8 ; 2 Hz, C_6 -H), 7.39(d, J=8 Hz, C_7 -H), 9.24(NH).
 Compound 8:mp.65-66°C (n-Hexane), uv λ EtOH/max. 203, 225, 236, 240, 330nm. ir ν Nujol/max. 3370, 1640, 1580 cm^{-1} . 1H -nmr($CDCl_3$) δ (ppm) 2.70(s, CH_3), 7.32(d, J=3 Hz, C_2 -H), 6.58(d, J=3Hz, C_3 -H), 7.77(d, J=8Hz, C_4 -H), 7.16(t, J=8Hz, C_5 -H), 7.89(d, J=8Hz, C_6 -H).

Received, 16th September, 1986