


FOUR NEW DIMERIC MONOTERPENE ALKALOIDS FROM SCAEVOLA RACEMIGERA
DÄNIKER (GOODENIACEAE)

Alexios-Leandros Skaltsounis, François Tillequin, and Michel Koch
Département de Pharmacognosie de l'Université René Descartes,
U.A. au C.N.R.S. n°484, Faculté des Sciences Pharmaceutiques et
Biologiques, 4 avenue de l'Observatoire F. 75006 Paris, France
Jacques Pusset and Gerard Chauvière
Laboratoire des Plantes Médicinales du C.N.R.S., Montravel, Nouméa,
New Caledonia


Abstract - Four new dimeric monoterpene alkaloids (1-4) have been isolated from the aerial parts of Scaevola racemigera Däniker. Their structures have been elucidated by spectroscopic studies and chemical correlations.

Scaevola racemigera Däniker (Goodeniaceae) is a tall shrub indigenous to the coastal regions of New Caledonia^{1,2}. Various monoterpene alkaloids had been previously isolated from the aerial parts of this species³. A thorough search for the more polar constituents of the alkaloid extract has now led to the isolation of four novel dimeric monoterpene alkaloids named scaevodimerines A, B, C and D. The present paper describes their structure determination.


Scaevodimerine A (1) has been obtained as a colourless amorphous solid, $[\alpha]_D^{20} +31^\circ$ (MeOH, c = 1) (contents : 0.02 % from the dried plant material⁴). Its empirical formula has been determined by high resolution mass spectroscopy as $C_{21}H_{25}N_3O_4$ (Found : 383.1831 ; Calcd. : 383.1845). The uv spectrum exhibited characteristic absorptions at $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ) : 214(4.06), 237(4.00), 269(4.30) and 335(3.70) associated with a pyridine-dihydropyridine chromophore^{5,6}. The ir spectrum displayed bands at $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 1715 and 1665 typical to a conjugated carboxylic ester. The ¹H nmr spectrum (Table I) exhibited all the characteristic signals of a tetrahydrocantlyline unit³. The chemical shift of H-6 (5.38 ppm instead of 4.06 ppm for tetrahydrocantlyline itself) gave evidence for an esterifi-




1




2




3



4



5



6

cation by a carboxylic acid at the 6-position³. Additional signals on the nmr spectrum permitted depicting the structure of the esterifying unit as 1,2-dihydro-1-methyl-2,7-naphthyridine-4-carboxylic acid. Of particular interest were the coupling constants between Me-1', H-1', NH-2' and H-3' which provided evidence for the 1'-position of the methyl group. Methanolysis of scaevodimerine A (0.25 N MeONa in MeOH/reflux under Ar/72 h) led in 40 % yield to an equimolecular mixture of 1,2-dihydro 4-carbomethoxy-1-methyl-2,7-naphthyridine (5) characterized by its ¹H nmr spectrum (Table I) and of tetrahydrocantleyine identical with the natural product (uv, ir, sm, ¹H nmr, $[\alpha]_D^{20}$)³. This experiment gave evidence for the absolute configuration at C-4a, C-6, C-7 and C-7a. Unfortunately, the absolute configuration at C-1' on the 1,2-dihydro 2,7-naphthyridine moiety of the molecule remains unknown since 5 spontaneously aromatizes by air-oxidation at room temperature within a few hours to the corresponding achiral naphthyridine 6. The uv spectrum of this latter compound exhibited absorptions at $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ) : 222(3.80) and 290(3.70) typical to a 2,7-naphthyridine chromophore⁷. The ir spectrum of 6 displayed characteristic bands^{8,9} at ν_{\max}^{KBr} cm⁻¹ : 1725 (ester, c = O), 1605 and 845 (2,7-naphthyridine and its ¹H-nmr spectrum (Table I) was very close to that described for 2,7-naphthyridine itself¹⁰, except for the chemical shifts of H-3 and H-5 strongly deshielded by the carbomethoxy group at C-4.

Scaevodimerine B (2) has also been obtained as a colourless amorphous solid, $[\alpha]_D^{20} +39^\circ$ (MeOH, c = 1) (contents : 0.05 % from the dried plant material). The empirical formula could be established by high resolution mass spectrometry as C₂₁H₂₃N₃O₄ (Found : 381.1700 ; Calcd. : 381.1688). Its general spectral features were closely related to those of 1. Nevertheless, differences could be noted on its uv spectrum whose typical absorptions : $\lambda_{\max}^{\text{EtOH}}$ nm (log ϵ) : 222(3.99), 283(4.03) were only associated with a pyridine-derivated chromophore. Its ¹H nmr spectrum (Table I) exhibited the characteristic signals of tetrahydrocantleyine unit esterified at the 6-position³ and those previously encountered in the spectrum of naphthyridine 6 with the exception of that associated with the COOMe group. Methanolysis of scaevodimerine B (1N MeONa in MeOH/reflux/2h) led to tetrahydrocantleyine and to 4-carbomethoxy-1-methyl-2,7-naphthyridine 6 in almost quantitative yield. This experiment provided evidence for both stereochemistry on the tetrahydrocantleyine moiety of the molecule and substitution pattern on the 2,7-naphthyridine unit. The structure of scaevodimerine B could thus be depicted as 2. In a similar way, the structures of scaevodimerine C (3), $[\alpha]_D^{20} -18^\circ$ (MeOH, c = 1),





TABLE I

¹H N.M.R. spectra of scaevodimerines A, B, C, D and of compounds 5 and 6 (270 MHz, δ ppm/TMS, J in Hz).

	<u>1</u> ^{a,c}	<u>2</u> ^a	<u>3</u> ^{a,c}	<u>4</u> ^a	<u>5</u> ^b	<u>6</u> ^b
H-1a	2.89 ddd : 13,7,2	2.82 ddd : 13,7,2	8.58	-	-	-
H-1b	3.26 ddd 13,5,2	3.33 ddd : 13,5,2	s	-	-	-
NH-2	4.58 br.s. D ₂ O exch.	4.48 br.s. D ₂ O exch.	-	7.44 br.s. D ₂ O exch.	-	-
H-3	7.53 d 7, tr. to s by D ₂ O	7.53 d 7, tr. to s by D ₂ O	9.07 s	7.22 d, 6, tr. to s by D ₂ O	-	-
COOMe-4	3.67 s	3.68 s	3.96 s	3.73 s	-	-
H-4a	5.11 td 9,7	3.22 td 9,7	-	3.59 ddd 11,9,7	-	-
H-5a	1.71 ddd 15,9,5	1.87 ddd 15,9,5	3.53 m	1.89 ddd 11,9,7	-	-
H-5b	2.34 ddd 15,7,2	2.32 ddd 15,7,2	3.53 m	2.64 m	-	-
H-6	5.36 ddd 5,4,2	5.56 ddd 5,4,2	5.80 td 5,5	5.55 td 4,2	-	-
H-7	2.00 m	2.15 m	3.53 m	2.64 m	-	-
Me-7	1.04 d 7	1.14 d 7	1.38 d 7	1.34 d 7	-	-
H-7a	2.00 m	2.15 m	-	2.82 dd 11,8	-	-
H-1'	4.96 qd 7,2 tr. to q by D ₂ O	-	1.89 qd 7,2, tr. to q by D ₂ O	-	4.74 qd 7,2, tr. to q by D ₂ O	-
Me-1'	1.49 d 7	3.13 s	1.48 d 7	3.13 s	1.27 d 7	3.03 s
NH-2'	7.52 br.s. D ₂ O exch.	-	7.06 br.s. D ₂ O exch.	-	8.24 br.s. D ₂ O exch.	-
H-3'	7.94 d 7, tr. to s by D ₂ O	9.63 s	7.71 d 7, tr. to s by D ₂ O	9.67 s	7.73 d 7, tr. to s by D ₂ O	9.71 s
COOMe-4'	-	-	-	-	3.60 s	3.90 s
H-5'	8.20 d 5	8.78 d 5	8.20 d 6	8.81 d 5	7.96 d 5	8.57 d 5
H-6'	8.57 d 5	8.82 d 5	8.38 d 6	8.84 d 5	8.20 d 5	8.82 d 5
H-8'	8.11 s	9.20 s	8.10 s	9.27 s	8.08 s	9.07 s

a : in CDCl₃ ; b : in CD₃SOCD₃ ; c : recorded at 65°C

Scheme 1



$C_{21}H_{21}N_3O_4$ (h.r. ms ; Found : 379.1521 ; Calcd. : 379.1532), (contents : 0.01 % from the dried plant material) and scaevodimerine D (4), $[\alpha]_D^{20} +48^\circ$ (MeOH, $c = 0.2$), $C_{21}H_{21}N_3O_5$ (h.r. ms ; Found : 395.1465 ; Calcd. : 395.1481), (contents : 0.005 % from the dried plant material) could be determined as 3 and 4. Of particular interest were their 1H nmr spectra (Table I) which exhibited signals typical to cantleyine^{11,12} and 1,2-dihydro-1-methyl-2,7-naphthyridine -4-carboxylic ester units for 3 and to strychnovoline¹³ and 1-methyl-2,7-naphthyridine -4-carboxylic ester units for 4.

Scaevodimerines are the first dimeric monoterpene alkaloids and seem related to dimeric iridoid derivatives^{14,15}. Their structures involve a 2,7-naphthyridine skeleton which has been previously seldom encountered in higher plants belonging to the families Oleaceae^{16,17} Scrofulariaceae⁹ and Valerianaceae¹⁸. This naphthyridine unit arises most probably from ammonia condensation with a secologanin-derived dialdehyde^{19,23} followed by rearrangements^{24,25} shown in Scheme 1. Since the alkaloids described in this paper have been extracted by standard means using ammonia as a base, a second alkaloid extraction avoiding the use of a nitrogen-containing alkaline agent and a study of the iridoid contents of the plant seem necessary to understand the exact biogenetic origin and the chemotaxonomic significance of the naphthyridine derivatives.

REFERENCES AND NOTES

1. Part CVIII in the series "Plantes de Nouvelle-Calédonie".
Part CVII, see "Epi-3 antirhine, un nouvel alcaloïde extrait de Guettarda heterosepala", C. Kan-Fan, M.H. Brillanceau and H.P. Husson, J. Nat. Prod., submitted for publication.
2. A.U. Däniker, Mitt. Bot. Mus. Univ. Zürich, 1933, 142, 475.
3. A.-L. Skaltsounis, S. Michel, F. Tillequin, M. Koch, J. Puset and G. Chauvière, Helv. Chim. Acta, 1985, 68, 1679.
4. The plant material has been collected at the Mont Kaala (New Caledonia) in 1982. Herbarium samples (PUCH 250) are held in the herbaria of the Centre ORSTOM de Nouméa.
5. A.W. Sangster and K.L. Stuart, Chem. Rev., 1965, 65, 69 and references cited therein.
6. U. Eisner and J. Kuthan, Chem. Rev., 1972, 72, 1 and references cited therein.
7. N. Ikekawa, Chem. Pharm. Bull., 1958, 6, 269.

8. N. Ikekawa, Chem. Pharm. Bull., 1958, 6, 401.
9. K.J. Harkiss and D. Swift, Tetrahedron Lett., 1970, 4775.
10. W.W. Paudler and S.J. Cornrich, J. Heterocyclic Chem., 1970, 7, 419.
11. N.K. Hart, S.R. Johns and J.A. Lamberton, Aust. J. Chem., 1969, 22, 1283.
12. T. Sévenet, B.C. Das, J. Parello and P. Potier, Bull. Soc. Chim. France, 1970, 3120.
13. S. Michel, A.-L. Skaltsounis, F. Tillequin, M. Koch and L. Aké-Assi, J. Nat. Prod., 1985, 48, 86.
14. G.A. Cordell, in R.H.F. Manske, "The Alkaloids", Academic Press, New York, 1977, 16, 431.
15. J.L. El-Naggar and J.L. Beal, J. Nat. Prod., 1980, 43, 649 and references cited therein.
16. N.K. Hart, S.R. Johns and J.A. Lamberton, Aust. J. Chem., 1968, 21, 1321.
17. H. Ripperger, Phytochemistry, 1978, 17, 1069.
18. M.M. Janot, J. Guilhem, O. Contz, G. Venera and E. Cionga, Ann. Pharm. Franç., 1979, 37, 413.
19. R.T. Brown, C.L. Chapple, D.M. Duckworth and R. Platt, J. Chem. Soc. Perkin Trans. I, 1976, 160.
20. G. Kinast and L.F. Tietze, Chem. Ber., 1976, 109, 3640.
21. F.A. MacKellar, R.C. Kelly, E.E. van Tamelen and C. Dorschel, J. Am. Chem. Soc., 1973, 95, 7155.
22. H.C. Beyerman, L.A. van Dijk, J. Levisalles, A. Melera and W.L.C. Veer, Bull. Soc. Chim. France, 1961, 1812.
23. R.T. Brown and C.L. Chapple, Tetrahedron Lett., 1976, 787.
24. J. Stöckigt, H.P. Husson, C. Kan-Fan and M.H. Zenk., J. Chem. Soc. Chem. Comm., 1977, 164.
25. H.P. Husson, C. Kan-Fan, T. Sévenet and J.P. Vidal, Tetrahedron Lett., 1977, 1889.

Received, 14th October, 1986